

Efficiency of Large Double Auctions

Martin W. Cripps and Jeroen M. Swinkels*
John M. Olin School of Business
Washington University in St. Louis

October, 2003

Abstract

We consider large double auctions with private values. Values need be neither symmetric nor independent. Multiple units may be owned or desired. Participation may be stochastic. We introduce a very mild notion of “a little independence.” We prove that *all* non-trivial equilibria which satisfy this notion are asymptotically efficient. For any $\alpha > 0$, inefficiency disappears at rate $1/n^{2-\alpha}$.

1 Introduction

Many market settings are approximated by a double auction. Standard examples are the London gold market, and the order books maintained by NYSE specialists. These auctions typically have many traders on each side of the market.

More importantly, large double auctions are in some sense the “right” model for micro-foundations of price formation in competitive markets. Like a competitive market, a large double auction has many traders. However, unlike the standard competitive model, traders are strategic. Hence, if traders asymptotically ignore their effect on price this is a result, not an assumption. And, there is an explicit mechanism translating individual behaviors into prices. So, one of the thorniest problems of the standard Walrasian model – how does the market get to equilibrium if everyone is a price taker – is explicitly addressed. Finally, double auctions are a better setting for thinking about price formation than one-sided auctions, both because they are often a better match to reality, and especially because they capture the essential problems of trade better than a one-sided auction. A large one-sided auction allows one to ask if traded units end up in the right hands. But, it does not address whether the correct number of units trade in the first place.

*We thank, without implicating, George Mailath, Andy McLennan, Rich McLean, John Nachbar, Phil Reny, and Mark Satterthwaite. We also thank the Boeing Center for Technology, Information, and Manufacturing for financial support.

In a seminal paper, Rustichini, Satterthwaite and Williams (1994, henceforth RSW) consider a double auction in which n buyers and sellers draw private values *iid*. They show that symmetric, increasing, differentiable equilibria in this setting are in the limit efficient and that convergence is fast, of order $1/n^2$. This is especially attractive in light of experimental evidence on efficiency in double auctions with only a moderate number of players.¹

In independent work, Fudenberg, Mobius and Seidel (2003) extend RSW to a setting in which a one dimensional state is sampled and values are then drawn *iid* from a density that depends on the state, but has non-shifting support and uniform lower bound across states.² They also show existence of a pure increasing symmetric equilibrium when the number of players is large.³

These results are useful in thinking about how auctions approximate competitive equilibria. However, there are several dimensions along which they could be strengthened.

1. The proof technique depends heavily on symmetric distributions of values.
2. Even in the symmetric setting, there is no guarantee of uniqueness. So, while well behaved symmetric equilibria are asymptotically efficient, there may be other (possibly asymmetric) equilibria as well. In particular, there is always the no-trade equilibrium in which all buyers make an offer of zero, and all sellers make an offer higher than any possible valuation. Results before this paper do not rule out other intermediate trade equilibria.
3. While one may be willing to rule out the asymmetric equilibria on *a priori* grounds in the symmetric case, selecting the “good” equilibria is much harder if the initial setting is itself asymmetric.
4. Imposing symmetry on values and bids assumes away half the problem. Objects that trade automatically move from and to the right people, and so the only question is whether the volume of trade is right. Without symmetry, it may also occur that, for example, a low valued buyer wins an object when a higher valued buyer does not.
5. Finally, these papers consider only single unit demands and supplies.

We present a model and results addressing all these points. We consider a generalized private value double auction setting.⁴ Players can be highly asymmetric, and demand or supply multiple units. Beyond the assumption of private

¹Satterthwaite and Williams (2002) establish that in the *iid* setting, this rate is fastest among all mechanisms. Important precursors to RSW include Chatterjee and Samuelson (1983), Wilson (1985), Gresik and Satterthwaite (1989), and Satterthwaite and Williams, (1989).

²Our model will encompass this case. See Example 4 below.

³Jackson and Swinkels (2001) shows existence of non-trivial equilibria in double auctions. FMS shows that in the setting they consider, one of these equilibria is pure and increasing.

⁴A beautiful paper by Perry and Reny (2003) extends the previous work on information aggregation in large one sided common value auctions (Wilson 1977, Milgrom 1979, Pesendorfer and Swinkels 1997, etc.) to the double auction setting. A symmetric single unit demand and supply setting is maintained. Using a discrete bid space to get existence, they show that in a

values, there are only three assumptions with any bite. First, while individual values need be neither full support or even non-degenerate, we require that any given interval in the support of values is eventually hit in expectation by many players. We term this condition *no asymptotic gaps (NAG)*. Analogously, we will require there to be *no asymptotic atoms (NAA)*: it cannot be the case that a positive limiting fraction of players are expected to pile up in the same arbitrarily small interval.

Most critically, we drastically relax independence. We require only that a “little” independence across players persists as the number of players grows. A sequence of distributions over player values satisfies z -independence, $z \in (0, 1]$ if the probability of any given event on player i 's values changes by factor bounded between z and $1/z$ when one conditions on the values of the remaining players, where z holds uniformly in the number of players. 1-independence is the standard notion of independence, while two perfectly correlated random variables do not satisfy z -independence for any $z > 0$.

An interpretation of z -independence is that each player has at least a small idiosyncratic component to his valuation, one that cannot be precisely predicted no matter how much one knows about the values of other players. As such, this is a fairly weak condition, admitting very broad classes of distributions.

Because values can be highly correlated (positively, negatively or otherwise) under z -independence, even in the limit the allocation and price setting problem will generally be non-trivial.

There is always a no-trade equilibrium in a double auction setting. Jackson and Swinkels (2001, henceforth JS) show that there is at least one non-trivial equilibrium as well. Our major result is simply stated:

As the number of players grows, every non-trivial equilibrium of the double auction setting converges to the Walrasian outcome. Inefficiency disappears at rate $1/n^{2-\alpha}$ for any $\alpha > 0$.

Asymptotic efficiency implies asymptotic uniqueness and pureness: over relevant ranges, bids must be arbitrarily close to value. Thus, as n grows large, there are precisely two types of equilibria of private value double auctions:

1. equilibria involving no trade
2. equilibria in which a near efficient level of trade occurs, at a price near the competitive one.

With single unit demands and supplies, our proof works because in each outcome of a double auction, there is at most one buyer who is both currently winning an object and who would have raised the price had he bid more (the lowest winning buyer). So, while many buyers might have raised price by bidding more, only one would care that he did so. This is symmetric for sellers

one dimensional affiliated setting, the equilibrium price converges to the rational expectations equilibrium value. So, Perry and Reny generalizes RSW in the direction of non-private values while retaining most of other restrictions, while we generalize RSW in most other directions, while, critically, retaining private values.

considering lowering their bids. So, the expected *relevant* impact on price from increased bids by buyers is already order $1/n$. And, with lots of bidders, even if an increase in bid increases price, it should do so by an amount related to $1/n$, since this should be the expected distance to the next bid. But then, since the expected impact on price is order $1/n^2$, it must be that bidding honestly almost never wins an extra object, and so those objects that are traded must be allocated very efficiently.

The focus then turns to showing that the right number of objects trade, or, equivalently, that the *competitive gap* defining the range of market clearing prices grows small. This turns out to be much the hardest part of the paper (especially with a rate). In the symmetric case, one can appeal to the first order conditions of players near a discontinuity in bids. Here, things are much more difficult, as without symmetric increasing strategies, (a) the very concept of a “gap” becomes more complicated (b) it is hard to identify which player types might bid near a gap, and (c) players can have very different beliefs about the likelihoods of the events involved. We show that the only way to have a significant competitive gap without violating the efficiency already shown for those objects traded is for the market to essentially become deterministic, with a given set of buyers and sellers always trading. But then, any member of either of these groups can favorably influence the price without losing the chance to trade.

The efficiency result generalizes to multiple unit demands as long as *NAG* continues to hold for the first unit of demand and supply for each player. If this holds, we can reformulate the arguments just outlined applied only to the highest bid by each buyer and lowest bid by each seller to show small price impacts of honest bidding. From there to (fast) efficiency for all units involves a careful tracking of incentives, but is otherwise straightforward.

We begin by setting up the basic single unit demand and supply model. We then introduce z -independence. Analysis of efficiency for the large double auction with single unit demands and supplies follows. Then, we generalize to auctions with multiple unit demands and supplies. We conclude with some thoughts on extensions. All proofs are relegated to an appendix.

2 The Model

We begin with the structure of a given double auction \mathcal{A} . A finite set N of players is divided into subsets N_S and N_B . Players in N_S are potential sellers, each with one unit to sell. Players in N_B are potential buyers, each desiring a single unit.

Each $i \in N$ has valuation v_i . For sellers, this might be either a production cost or a value in use. For $i \in N_B$, we assume $v_i \in [0, 1)$. For $i \in N_S$, we assume $v_i \in (0, 1]$. A buyer with value 0 or a seller with value of 1 will never trade. Because of this, there is no loss of generality in assuming an equal number of buyers and sellers. Let $n \equiv |N_s| = |N_B|$. Because one can “park” extra buyers at 0 and extra sellers at 1, the model also allows a stochastic number of buyers

and sellers. The vector $v \equiv \{v_i\}_{i \in N}$ is drawn according to a probability measure P on $[0, 1]^n \times (0, 1]^n$. The marginal of P onto v_i is P_i .

Each player i observes his value and then submits a bid $b_i \in [0, 1]$. Trade is determined by crossing the demand and supply curves constructed from the submitted buy and sell bids.⁵ Call the (random) range of possible market clearing prices the *competitive gap*, $cg \equiv [\underline{cg}, \overline{cg}]$. If we let $b^{(i)}$ denote the i^{th} highest bid, then a little time with the appropriate figure shows that $cg = [b^{n+1}, b^n]$.

Assumption 1 *Trade takes place at price*

$$p = \hat{p}(cg, \overline{cg})$$

where \hat{p} is differentiable, takes values in $[\underline{cg}, \overline{cg}]$, and has derivatives bounded by 0 and 1.⁶

Imagine that the bidder who submitted \overline{cg} raises his bid substantially. As long as his bid continues to define \overline{cg} , he raises the price at rate at most 1. As soon as he passes the next bid up, he ceases to affect price. Let $\overline{ug} = b^{n-1}$ be this next bid, and define the *upper supporting gap* as $ug \equiv [\overline{cg}, \overline{sg}]$. Then, the maximum effect on the price is $|ug|$. Similarly, let $\underline{lg} = b^{n+1}$, and define the *lower supporting gap* as $lg \equiv [\underline{lg}, \underline{cg}]$. So, cg determines the amount of choice there is in setting a market price, while lg and ug determine how closely “supported” this range is.

Each player i has a *vNM* utility function u_i . No particular structure on risk preferences is required, but we do require each u_i to be increasing and have slope bounded from 0 and ∞ .⁷

2.1 Equilibrium

A set of distributional strategies $\{\mu_i\}_{i \in N}$ (Milgrom and Weber, 1982) is an *equilibrium* if it is a Bayesian Nash equilibrium in which buyers never bid above v_i , and sellers never bid below v_i . The equilibrium is *non-trivial* if there is a positive probability of trade.

We show that non-trivial equilibria are asymptotically efficient. This, of course, is a better result if such equilibria exist! Under slightly stronger conditions than we use here, JS show that this is indeed the case.⁸

2.2 Sequences of Auctions

Consider a sequence of such auctions $\{\mathcal{A}^n\}$, where n tends to infinity. We need three conditions that apply across n . First, while individual values need not

⁵If tied buy and sell bids allow more than one level of trade, the largest is chosen.

⁶This of course includes the standard k double auction.

⁷In the proofs, we assume risk neutrality. Dealing with *vNM* utility functions with slope bounded from 0 and ∞ involves scaling potential gains down by some factor from the risk neutral case, and potential losses up. This merely introduces notation.

⁸The two key assumptions are mutual absolute continuity of P with respect to $\Pi_i P_i$, and atomless P_i . Neither assumption plays any further role in the development here.

have full support (and may, in fact, be atomic), we require that as n grows large, each subinterval is hit with non-vanishing probability.

Assumption 2 (No Asymptotic Gaps) *There is $w > 0$ such that for all n , and for all intervals $I \subseteq (0, 1)$ of length $1/n$ or greater,*

$$\sum_{i \in N_B} P_i[I] \geq wn |I|$$

and

$$\sum_{i \in N_S} P_i[I] \geq wn |I|.$$

Note that P, N_B, N_S etc. all vary from one \mathcal{A}^n to another. We suppress this in our notation as convenient.

Our second assumption is similar:

Assumption 3 (No Asymptotic Atoms) *There is $W < \infty$ such that for all n , and for all intervals $I \subseteq (0, 1)$ of length $1/n$ or greater,*

$$\sum_{i \in N_B} P_i[I] \leq Wn |I|$$

and

$$\sum_{i \in N_S} P_i[I] \leq Wn |I|.$$

That is, not too many values fall in any given interval. These conditions hold only on $(0, 1)$, allowing a positive mass of buyers with value 0 or sellers with value 1, consistent with our earlier discussion of “parking” extra players.

Example 1 *Let sellers $i \in \{1, \dots, n\}$ have $v_i \equiv i/n$ and similarly for buyers. NAG and NAA are satisfied for $w = W = 1$. So, individual values need neither have full support nor be non-atomic.*

Example 2 *Each P_i is continuous with density bounded by w and W .*

Each of these two assumption has an analog in RSW. NAA is needed for a rate of convergence result, but not for convergence itself.

3 Z-Independence

Our final condition is the most important. We wish to relax independence considerably while still requiring “some persistent independence” as the population grows.

We require that knowledge about the values of players other than i provides at most a finite likelihood ratio on the values of player i , independent of how many other players there are.

Definition 1 *The sequence of probability measures $\{P^n\}$ satisfies z -independence, $z \in (0, 1]$, if for all n , for all $i \in N$, for any positive probability event F_{-i} involving only v_{-i} and any positive probability event F_i involving only v_i ,*

$$z \Pr(F_i) \leq \Pr(F_i|F_{-i}) \leq \frac{1}{z} \Pr(F_i). \quad (1)$$

That is, there is still some idiosyncrasy in each v_i even as the market becomes large.⁹

For fixed n , z -independence is slightly stronger than mutual absolute continuity (consider a uniform and a triangular distribution on $[0, 1]$) but weaker than having a continuous Radon-Nikodym derivative bounded from 0 and ∞ . The real content of z -independence is in the uniformity of z across n .

Assumption 4 (z -independence) *There exists $z > 0$ such that $\{P^n\}$ is z -independent.*

3.1 Examples

Example 3 *With probability $1/2$, values are drawn iid uniform $[0, 1]$, and with probability $1/2$, x is drawn uniformly from $[1/n, 1 - 1/n]$, and values are drawn iid uniform $[x - 1/n, x + 1/n]$. For each n , P^n is absolutely continuous with respect to $\prod P_i$ (and, the example is easily modified such that the Radon-Nikodym derivative is continuous as well). But, as $n \rightarrow \infty$, seeing the values of two randomly selected players within $2/n$ of each other makes it arbitrarily likely that all remaining players will also have such a value.*

We would like this example to be ruled out by our notion of “some persistent independence.” A first thought might be to require that no matter what we know about one subset of the players’ values, beliefs about the rest of the players’ values are updated by at most a finite ratio. This turns out to be much too strong.

Example 4 *Nature chooses $x \in \{L, H\}$ equiprobably. If L is drawn, values are drawn iid according to density $f(v) = 1/2 + v$. If H is drawn, values are drawn iid according to density $f(v) = 3/2 - v$.*

A finite likelihood ratio condition fails if both events involve large numbers of players. For example, let F_O be the event that less than 50% of the odd numbered buyers have value below $1/2$, and let F_E be the event that less than 50% of the even numbered buyers have value below $1/2$. Then, as $n \rightarrow \infty$, $\Pr(F_O|F_E) \rightarrow 1$, while $\Pr(F_O|F_E^C) \rightarrow 0$. Since for each n , F_E and F_E^C have the same size, this also means that the Radon-Nikodym derivative satisfies no uniform bound across n .

⁹A contemporaneous paper by Peters and Severinov (2002) uses a similar condition (in a different model) in a finite type setting.

This example exhibits a great deal of independence despite the fact that likelihood ratios and Radon-Nikodym derivatives diverge. We would like to admit it.

Note that Example 3 fails z -independence for any $z > 0$, as v_i is, $1/2$ of the time, arbitrarily closely predicted by v_{-i} . However Example 4 satisfies $.5$ -independence; all one can extract from v_{-i} is information about whether x is L or H , which changes the density on v_i from 1 to something between $1/2$ and $3/2$.

Example 4 generalizes to any process in which a state is sampled and then, conditional on the state, values v_i are drawn independently from measures with non-moving support V_i according to densities uniformly bounded (across states and n) away from zero and infinity. So our setting encompasses Fudenberg et al. (2003) (and more importantly, non-symmetric analogues to their model). However, even with symmetry, z -independence admits many distributions which cannot be generated in this way.

Example 5 *There are 2 players. The density on values is 2 on $[0, \frac{1}{2}] \times [0, \frac{1}{2}]$, and $2/3$ elsewhere. This satisfies $\frac{2}{3}$ -independence. One can generate this using states and conditionally-independent values. In state θ_1 (which occurs 1 time in 3) values are uniform on $[0, \frac{1}{2}] \times [0, \frac{1}{2}]$, while in state θ_2 they are uniform on $[0, 1] \times [0, 1]$. But, one cannot do so without shifting supports.*

Postlewaite and Schmeidler (1986) define *non-exclusivity* as a situation where the information of $n - 1$ players is enough to predict the relevant state of the economy. A variety of follow-on papers relax this to hold only asymptotically.¹⁰ On first view, z -independence is antithetical to non-exclusivity, since no matter how much is known about the rest of the players, the value of player i remains uncertain. However, note that non-exclusivity refers to information about the underlying state, not to the signals players realize conditional on those states. In Example 4, v_{-i} is asymptotically fully informative about L vs. H , while of bounded informativeness about v_i . Hence Example 4 can satisfy both conditions.

Example 6 *Nature draws v_1 uniformly from $[0, 1]$ (this person is a “fashion leader”), and then draws subsequent players iid according to a density with support $[0, 1]$ but concentrated around v_1 .*

Since the impact of an early draw on later draws does not vanish, z -independence does not imply weak mixing. It is also easy to construct sequences satisfying weak mixing under which successive draws are arbitrarily correlated, violating z -independence.

Example 7 *A parameter x is chosen from $[0, 1]$. Values are drawn conditionally independently according to $f(\cdot|x)$, where $f(\cdot|x)$ satisfies MLRP in x . As long as $f(\cdot|0)/f(\cdot|1)$ is uniformly bounded, z -independence is satisfied for*

¹⁰A good entry point is McLean and Postlewaite (2002).

$z = \min_x f(x|0)/f(x|1)$. Choose a subset of the players, and replace v_i by $1 - v_i$. This measure continues to satisfy z -independence, but is obviously not affiliated. So, affiliation has essentially nothing to do with the issues at hand.

We close this subsection with an example illustrating the surprising degree of correlation z -independence can imply. Define $[x]$ as the largest integer smaller than x .

Example 8 For $m \leq n$, let $\zeta_B(m) = \binom{n}{m} (.5)^n$ be the probability of m heads from n flips of a fair coin.¹¹

Now, for some $0 < a < 1/2$, generate ζ_C from ζ_B by first defining $\zeta'_C(m) = \zeta_B(m)a^{|m-[n/2]|}$, and then defining ζ_C from ζ'_C by normalizing. Informally one makes each outcome successively further away from $[n/2]$ more unlikely by a factor of a . Choose m according to ζ_C , choose each subset of coins of size m with equal probability, and make the coins in the subset heads, and the remainder tails. When a is small, drawing exactly $[n/2]$ heads by this process becomes very probable.¹² For $a = .1$, e.g., there is an 80% chance of exactly $[n/2]$ heads regardless of n .¹³

This process satisfies z -independence! If there are m' heads among all but coin i , the probability that i is heads is $\Pr(m = m' + 1)/\Pr(m = m' + 1)$. By construction, this is either a or $1/a$.

Thus, z -independence does not imply limit “noise.” So the techniques in Mailath and Postlewaite (1990), Al-Najjar and Smorodinsky (1997), and Swinkels (2001) do not apply.

3.2 A Preliminary Lemma

Our first lemma shows that if values are z -independent then so too are bids. The intuition for this is that b_i is a garbling of v_i .¹⁴ It also describes the implications of z -independence for groups of players.

¹¹The example can easily be extended from coins to values in the standard domain.

¹²Note that

$$\begin{aligned} \sum_{m=0}^r \zeta'_C &= \sum_{m=0}^r \zeta_B(m)a^{|m-r/2|} \leq \zeta_B(r/2) \sum_{m=0}^r a^{|m-r/2|} \\ &\leq \zeta_B(r/2) \left(1 + 2 \sum_{i=1}^{\infty} a^i\right) = \zeta_B(r/2) \left(1 + \frac{2a}{1-a}\right) \end{aligned}$$

Thus,

$$\zeta_C(n/2) \geq \frac{\zeta_B(n/2)}{\zeta_B(n/2) \left(1 + \frac{2a}{1-a}\right)} = \frac{1}{\left(1 + \frac{2a}{1-a}\right)}.$$

As $a \rightarrow 0$, this tends to 1.

¹³For $a = .1$, the previous expression is equal to $\frac{1}{\left(1 + \frac{2(.1)}{1-.1}\right)} = .81$.

¹⁴A related lemma appears in JS.

Throughout the paper, for any non-empty $K \subset N$, when we write F_K (respectively $F_i, F_{-i}, F_{N \setminus K}$), we mean an arbitrary positive probability event involving only the values or bids of the players in K ($\{i\}, N \setminus i, N \setminus K$).

Lemma 1 *Fix a non-empty $K \subset N$. Let $a = \min\{|K|, |N \setminus K|\}$. Then for all F_K , and $F_{N \setminus K}$,*

$$z^{-a} \Pr(F_K) \geq \Pr(F_K | F_{N \setminus K}) \geq z^a \Pr(F_K). \quad (2)$$

Let X_K be a random variable that depends only on the values/bids of the players in K . Then:

$$z^{-a} E(X_K) \geq E(X_K | F_{N \setminus K}) \geq z^a E(X_K). \quad (3)$$

When a is large, these bounds are weak; for arbitrary events involving many players, likelihood ratios can explode.

3.3 Large Deviations

Given $K \subset N$ and events $\{F_i\}_{i \in K}$ let Q_K be the number of F_i that are true. Notice that $E(Q_K) = \sum_{i \in K} \Pr(F_i)$. Let us stochastically bound Q_K . Note first that for each i ,

$$\Pr(F_i | F_{-i}) \geq z \Pr(F_i).$$

Note also that

$$\Pr(F_i^c | F_{-i}) \leq \frac{1}{z} \Pr(F_i^c) = \frac{1}{z} (1 - \Pr(F_i))$$

and so

$$\Pr(F_i | F_{-i}) \geq 1 - \frac{1}{z} (1 - \Pr(F_i)).$$

Thus,

$$\Pr(F_i | F_{-i}) \geq \underline{p}_i \equiv \max \left\{ z \Pr(F_i), 1 - \frac{1}{z} (1 - \Pr(F_i)) \right\}.$$

Since this is true for all F_{-i} we show that Q_K first order stochastically dominates $|K|$ independent coins with parameters \underline{p}_i .

Similarly, $|K|$ independent coins with parameters

$$\bar{p}_i \equiv \min \left\{ \frac{1}{z} \Pr(F_i), 1 - z(1 - \Pr(F_i)) \right\}$$

stochastically dominate Q_K .

Sets of independent coins are well understood. We can apply the theory of large deviations to obtain:

Lemma 2 *For all $K \subset N$ and $F_{N \setminus K}$,*

$$\Pr \left(Q_K < \frac{z}{3} E(Q_K) \mid F_{N \setminus K} \right) \leq e^{-.3zE(Q_K)} \quad (4)$$

$$\Pr \left(Q_K > \frac{3}{z} E(Q_K) \mid F_{N \setminus K} \right) \leq e^{-E(Q_K)} \quad (5)$$

This casts light on Example 8. Under z -independence, probabilities that start in the interior of $(0, 1)$ cannot be moved too far toward or away from the boundaries. But, probabilities can be moved around essentially arbitrarily within $(\underline{p}_i, \bar{p}_i)$.

A useful implication of Lemma 2 is that the probability of at least one success is not drastically affected by $F_{N \setminus K}$.

Corollary 1

$$\Pr(Q_K \geq 1 \mid F_{N \setminus K}) \geq (1 - e^{-z}) \Pr(Q_K \geq 1). \quad (6)$$

3.4 Normal Realizations

We prove convergence at rate $1/n^{2-\alpha}$ for any given $\alpha > 0$. It is convenient to fix α now. We will need various fudge factors along the way. Choose $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ so that

$$\alpha > \alpha_1 > \alpha_2 > \alpha_3 > \alpha_4 > 2\alpha/3$$

Let $w' \equiv \frac{z}{6}w$ and $W' \equiv \frac{6}{z}W$.

Definition 2 *A realization is normal if every interval $I \subseteq (0, 1)$ of length $1/n^{1-\alpha/3}$ or greater has between $w'n |I|$ and $W'n |I|$ buyers (respectively sellers) with value in that interval.*

Let \mathcal{N} be the event that the realization is normal.

Say that a statement is true for n sufficiently large (n SL) to mean that there exists an n^* depending only on the parameters such that the statement is true for all $n > n^*$. Then, a key implication of Lemma 2 is

Lemma 3 *For all n SL, $\Pr(\mathcal{N}) \geq 1 - 1/n^4$.*

Together with NAG and NAA, Lemma 3 implies that the limiting realized true demand and supply curves are unlikely to have either vertical or flat sections (except at 0 for buyers and 1 for sellers).¹⁵

4 Analysis of the Double Auction

4.1 Summing Deviations

Fix an equilibrium μ of \mathcal{A}^n . Consider buyer i 's distributional strategy μ_i . A *deviation* for i is a measurable mapping d_i from $[0, 1]^2$ to $[0, 1]$. First i draws

¹⁵We show that percentage efficiency losses are asymptotically less than $1/n^{2-\alpha}$. As for all rate of convergence results, this does not say anything about small n . The construction underlying normality in particular only holds for n pretty large. We use normality to sidestep a set of statistical issues related to the generality of our set-up, especially non-symmetry. There seems to be nothing in the underlying *incentives* being exploited that precludes much faster convergence, and our expectation would be that actual convergence is indeed very fast. RSW supplement their rate result (where the constant is again large) with numerically solved examples. Such solutions are beyond our ability in this setting.

v_i and b_i according to μ_i , but then she modifies her chosen bid according to d_i . Consider d_i for which $b_i \leq d_i(b_i, v_i) \leq v_i \forall b_i, v_i$. That is, i sometimes raises her bid, but not beyond her true value (since μ_i did not involve i bidding more than her true value, this is coherent).

In any given realization, d_i may have benefit \hat{B}_i in that i wins when he otherwise would not have, or may have cost \hat{C}_i that i pays more when he would have already won. To formalize this, let p be price under μ , and p_d the price when i uses d_i . Let W_i be the event that i wins with d_i , but not without. Then

$$B_i \equiv E(\hat{B}_i) = \Pr(W_i)E(v_i - p_d|W_i).$$

Let $O_i \subset W_i^c$ be the event that i wins without d_i . Then

$$C_i \equiv E(\hat{C}_i) = \Pr(O_i)E(p_d - p|O_i).$$

Since μ_i is a best response, $B_i \leq C_i$. So, given such a d_i for each buyer,

$$\sum_{N_B} B_i \leq \sum_{N_B} C_i.$$

Each d_i is unilateral. But, there is nothing wrong with summing the incentive constraints implied.

Consider $\sum_{N_B} C_i$. *Ex-post*, $\hat{C}_i > 0$ only if (a) trade was occurring and (b) the original b_i was equal to $\overline{c\bar{g}}$, and uniquely so. When $b_i > \overline{c\bar{g}}$ (or is tied at $\overline{c\bar{g}}$), increasing b_i does not affect p . If $b_i < \overline{c\bar{g}}$, increasing b_i may increase p , but as i was not originally winning, she is unhurt. So, there is at most one i with $\hat{C}_i > 0$.¹⁶ And, as discussed above, for this i , $\hat{C}_i \leq |ug|$. Thus,

$$\sum_{N_B} C_i \leq \Pr(T)E(|ug||T).$$

For sellers, the same analysis applies if bids are lowered, but not below value. We have thus established:

Lemma 4 *For any set $\{d_i\}_{i \in N_B}$, for which $d_i(b_i, v_i) \in [b_i, v_i]$ for all (b_i, v_i)*

$$\sum_{N_B} B_i \leq \Pr(T)E(|ug||T). \quad (7)$$

For any set $\{d_i\}_{i \in N_S}$, for which $d_i(b_i, v_i) \in [v_i, b_i]$ for all (b_i, v_i)

$$\sum_{N_S} B_i \leq \Pr(T)E(|lg||T). \quad (8)$$

While easy to prove, this bound is powerful. Independent of the number of bidders, the total benefit to players of bidding more aggressively in terms of making new trades must be small in equilibrium.

¹⁶If $\overline{c\bar{g}}$ is a seller's bid, no buyer is hurt by d_i .

4.2 The Probability of Trade is Bounded from Zero

An important first step is to show that non-trivial equilibria are not “almost trivial” in the sense that trade becomes increasingly rare as n grows. For each n , choose a non-trivial equilibrium of \mathcal{A}^n . Let V be the number of objects traded and T be the event that $V \neq 0$.

Our first lemma is technical.

Lemma 5 *Along any subsequence, if $\frac{E(V|T)}{n} \not\rightarrow 0$, then $\Pr(T) \rightarrow 1$.*

Intuitively, if many players trade given T , then many players must occasionally be bidding in a fairly aggressive way. But then, by z -independence, at least a fraction of them will be doing so almost all the time. The proof is more complicated because T is linked to all player’s actions, and so z -independence does not immediately apply.

Using Lemma 5, we can show:

Proposition 1 *There is $\gamma > 0$ such that for all n SL, and all non-trivial equilibria,*

$$\Pr(T) \geq \gamma.$$

For intuition, think about a situation where *in aggregate* buyers only make a “serious” offer with some probability δ close to 0, and symmetrically for sellers (clearly, if there is a non-vanishing probability of a serious offer on either side, trade will not disappear). Trade occurs at most 2δ of the time, since trade requires a serious offer from at least one side. Hence, by Lemma 4, the total costs to buyers (or sellers) of making more generous offers is like (has the same order as) δ . But, from z -independence, the probability of a serious offer from one side but not the other is like $(1 - \delta)\delta \cong \delta$. When there is a serious offer on one side but not the other, a number of bidders on the other side that grows like n would have benefited by deviating to trade at the serious offer. The gains are thus like $n\delta$, while costs are like δ . This is a contradiction.

4.3 Small Supporting Gaps

We show next that the upper and lower supporting gaps shrink quickly. This proceeds in two steps. First, we show that $E(|ug|)$ (respectively $E(|lg|)$) is like $1/n$. The idea is most easily seen if for each n , ug has constant length v . By Lemma 3, a number of buyers proportional to nv will have v_i in the top half of ug . At most one of these buyers is winning an object (they are not bidding above \overline{ug} , as bids are below value, and only one bid below \overline{ug} is filled). By raising b_i to $v - v/4$ all but this player (acting unilaterally) would win an extra object and earn at least $v/4$. So, $\sum B_i \geq nv^2$ (up to some constants). But by Lemma 4, $\sum C_i \leq v$, since the one person who is hurt raises the price by at most v . Thus,

$$nv^2 \leq \sum_{N_B} B_i \leq \sum_{N_B} C_i \leq v,$$

from which $v \leq \frac{1}{n}$. The actual proof has to count for the fact that $|ug|$ is stochastic, as are the number of bidders in any given interval. Formally:

Lemma 6 *For n SL and all x ,*

$$E(|ug|) \leq \frac{1}{n^{1-\alpha_4}}, \quad E(|lg|) \leq \frac{1}{n^{1-\alpha_4}}$$

Fix x , and consider $\Pr(|ug| \geq x)$. Consider again buyers raising b_i to $v - x/4$. When $|ug| \geq x$, then as above, a number of buyers like nx makes gains $x/4$, and so $\sum B_i = nx^2$ (again ignoring constants). And, $\sum C_i \leq E(|ug|) \leq 1/n$ from the first step. So

$$\Pr(|ug| \geq x)nx^2 \leq \frac{1}{n},$$

from which $\Pr(|ug| \geq x) \leq \frac{1}{n^2x^2}$. Formally

Lemma 7 *For n SL and all x ,*

$$\Pr(|ug| \geq x) \leq \frac{1}{n^{2-\alpha_3}x^2}, \quad \Pr(|lg| \geq x) \leq \frac{1}{n^{2-\alpha_3}x^2}.$$

For $x \geq 0$, let $L_B(x)$ be those buyers with values above $\bar{c}g + x$ that do not receive an object, and let $l_B(x) \equiv \#L_B(x)$. Similarly let $L_S(x)$ be those sellers with values below $\underline{c}g - x$ who do not sell, and let $l_S(x) \equiv \#L_S(x)$. Let

$$SL_B(x) \equiv \sum_{i \in L_B(x)} v_i - \bar{c}g, \quad SL_S(x) \equiv \sum_{i \in L_S(x)} \underline{c}g - v_i.$$

For buyers, this is the loss in consumer surplus compared with being able to price take at $\bar{c}g$, and analogously for sellers. Lemma 6 implies that both the number of such players and the associated loss is small. The intuition again comes from considering players bidding closer to their values.

Lemma 8 *For n SL and for all x ,*

$$E(l_B(x)) \leq \frac{1}{xn^{1-\alpha_4}}, \quad E(l_S(x)) \leq \frac{1}{xn^{1-\alpha_4}}.$$

Further

$$E(SL_B(1/n)) \leq \frac{1}{n^{1-\alpha_3}}, \quad E(SL_S(1/n)) \leq \frac{1}{n^{1-\alpha_3}}.$$

4.4 Small Competitive Gaps

Let us now turn to the competitive gap. Our key lemma:

Lemma 9 *For n SL and for all x ,*

$$\Pr(|cg| \geq x) \leq \frac{1}{n^{2-\alpha_2}x^2}.$$

To see the intuition for Lemma 9, consider first a fixed interval $I = (\underline{I}, \bar{I})$ such that n prespecified bidders always bid above \bar{I} (up) and the rest always bid below \underline{I} (down). Then, the competitive gap will always include I . And, since the probability of trade is bounded away from 0, the set of up bidders must contain a buyer, and the set of down bidders a seller. But then, by bidding $\underline{I} + \varepsilon$, any up buyer can still trade and force the price near the bottom of \underline{I} , while by bidding $\bar{I} - \varepsilon$, any down seller can still trade and force the price near \bar{I} , contradicting equilibrium.

If this situation arises only in the limit then the buyer or seller occasionally loses a trade by bidding more aggressively, but this becomes unlikely. Finally (because this is what we will really need), imagine that I shrinks as n grows. Then, as the gain from affecting the market price shrinks, we must be careful that the loss from lost trades shrinks as well. To do this, pick a buyer whose value is not too much above \bar{I} , so that his value of trade was quite small, and a seller whose value was not too much below \underline{I} . The efficiency of the allocation among buyers and sellers (Lemma 8) lets us do this.

We show that if Lemma 9 fails, then the limit is as described. For intuition, assume there is some interval I of length x such that nobody ever bids in I , and such that $\Pr(I \subseteq cg)$ does not fall quickly. Let p_i be the probability that i bids up, and q_i the probability of down. Order the players so that p_i is increasing. Run along them stopping at the player i where one counts $n-1$ ups. For $I \subseteq cg$, we need to hit exactly one more up in the rest of the sequence. If one hits no more ups, $I \subseteq ug$, while if one hits 2 more ups, $I \subseteq lg$, either of which is rare by Lemma 7. But, we argue, the only way to make 1 more up likely, but neither 0 nor 2 more ups likely is for the next player to have p_{i+1} nearly 1, and for the remaining players to *in aggregate* have almost no chance of even one up. Essentially, if p_{i+1} is not near one, then, since p_i is decreasing, the probability on who is the n^{th} up is “spread out”. But then, z -independence makes it likely that one also over or undershoots by 1. And, given that the next player is likely to hit, there must rarely be any more hits in the remaining population. Running through the players in reverse order and counting downs, when one hits $n-1$ downs, the next one must almost certainly play down, and then there must almost never be any more downs. Since both of these are true at once, in aggregate, the first n bidders almost always bid up and the remaining down. Hence, $\Pr(I \subseteq cg) \rightarrow 1$.

The proof is long: cg can move around, sometimes including one interval and sometimes another, players might bid not only above or below any given I , but sometimes within it, and one must be careful not to double count the ways in which a population “one player away” from creating a long cg might end up creating a long supporting gap.

4.5 Efficiency

We are now ready for our main theorem:

Theorem 1 *All non-trivial equilibria of the single unit demand/supply double auction are asymptotically efficient. Uniformly across non-trivial equilibria, efficiency losses go to zero faster than $1/n^{1-\alpha}$ for any given $\alpha > 0$. The fraction of expected surplus lost compared to a Walrasian market thus shrinks as $1/n^{2-\alpha}$.*

For intuition, note that in Section 4.3 we showed that the efficiency loss from failing to trade objects between sellers with value below \underline{cg} and buyers with values above \overline{cg} is small (of order $1/n$). So, the only efficiency losses to worry about are from pairs of buyers and sellers both having value in cg . The loss from missing such a trade is at most $|cg|$. And, using NAA, the number of such buyers and sellers is like $|cg|n$. So, the deadweight loss triangle from too little trade has area $|cg|^2 n$. But, from Lemma 9, $\Pr(|cg| \geq x) \leq \frac{1}{n^{2-\alpha} x^2}$, and so the expected loss here is like $1/n$ as well. Finally, from NAG, expected feasible surplus grows like n , and so proportional losses are like $1/n^2$. A formal accounting of efficiency losses is subsumed by the proof of the multiple unit case, and so omitted in the appendix.

4.6 Asymptotic Uniqueness of Equilibrium

In the space of allocations, *all* non-trivial equilibria converge to the Walrasian outcome. Over “relevant” ranges bids must thus converge to true values. So, if in the limit, the Walrasian price is either p_1 or $p_2 > p_1$, then, players with value near p_1 or p_2 must bid close to value. But it is difficult to show that, for example, a player with value well above p_2 must bid near value. A rate of convergence result for bids is thus cumbersome. Intuitively, over relevant ranges convergence should be order $1/n$.

5 Multiple-Unit Demands and Supplies

Assume now that each player has demand or supply for at most m units, for some fixed m . For buyers, let v_{ih} , $h \in \{1, \dots, m\}$, be i 's incremental value for unit h .¹⁷ For sellers, let v_{ih} be the incremental cost of unit h . We assume v_{ih} is non-increasing in h for buyers and non-decreasing for sellers. Bids are (non-increasing for buyers, non-decreasing for sellers) m -vectors. JS applies to show existence of equilibria in this setting, subject to the same strengthenings as before.

We assume the following version of NAG.

Assumption 5 (No Asymptotic Gaps*) *There is $w > 0$ such that for all n , and for all intervals $I \subseteq (0, 1)$ of length $1/n$ or greater,*

$$\sum_{i \in N_B} P_i[v_{i1} \in I] \geq wn |I|$$

¹⁷As before, we include atoms for buyers at 0 and sellers at 1. So, this does not imply that buyers have positive value for all m units or that sellers are want or are able to sell m units.

and

$$\sum_{i \in N_S} P_i[v_{i1} \in I] \geq wn |I|.$$

That is, when n is large, there are many buyers whose *highest* value might fall in any given interval, and many sellers whose *lowest* cost might fall into any given interval.¹⁸

As before, z -independence applies only across players, and does not restrict the relationship of the different values of any given player. NAA is assumed to apply to all values, not just the first. So, not too many v_{ih} fall in any given interval.

Theorem 2 *With NAG*, Theorem 1 continues to hold even with multiple-unit demands and supplies.*

Most of the incentive arguments rely only on the highest value unit of demand for buyers and lowest cost unit for sellers. The proof proceeds in two steps. Define \overline{ug} as the m^{th} bid up from \overline{cg} , and ug as $(\overline{cg}, \overline{ug})$. In the appendix, we show that Lemma 7 continues to hold for this definition of ug . The modification to the intuition is very small: when ug is long, there are many buyers with highest value in the top half of ug . But, only m of them can be winning a first object. Given this, Lemma 9 is easily extended as well. Instead of sorting players into those who play “up” and “down”, sort them into those who make 0 up bids, 1 up bid, etc. This is notationally intensive but straightforward and hence omitted.

Finally, we must show that since $|ug|$, $|cg|$ and $|lg|$ shrink quickly, inefficiency in the market disappears as $1/n$. A proof of this is in the appendix. To see the issues involved, note that for the single unit case (and for the first unit of demand in the multiple unit case), a buyer’s impact on the price is small for two reasons. First, he is unlikely to be pivotal. Second, even if he is pivotal, he doesn’t affect the price much, since the next bid up is likely to be close. We exploit both of these forces in showing Lemma 7 and Lemma 9 and their adaptations here.

For units of demand after their first, many buyers can simultaneously be in the position that in raising bids other than their first, they pay more for units they were already winning. To get around this, consider the deviation to honest bidding. In any given realization, let x be $\overline{ug} - \underline{cg}$. This is the maximum impact of i raising his m bids on price. If $v_{ih} < \underline{cg}$, then the deviation is irrelevant.

If $\underline{cg} \leq v_{ih} \leq \underline{cg} + 2mx$, then i may not benefit very much from any new unit won by raising b_{ih} , and may hurt himself by raising the price by as much as x on each of $m - 1$ units already being won. But, critically, because of NAA, the number of v_{ih} in $(\underline{cg}, \underline{cg} + 2mx)$ is only like nx (as always, ignoring constants). So, the expected cost to bidders from this case is like $E(nx^2)$. But, the modified versions of Lemma 7 and Lemma 9 give that $E(nx^2)$ is like $1/n$.

¹⁸There are less restrictive ways in which one might generalize NAG. For example, if each buyer’s first value is uniform $[3, 4]$, and their second value is uniform $[0, v_{i1}]$ then there are many buyer values in each range. An example in Section 5.1 of Swinkels (2001), suggests that this is not strong enough to guarantee efficiency.

And, the expected efficiency loss from such players not winning also falls like $1/n$.

Consider objects with v_{ih} above $\underline{cg} + 2mx$ where i is already winning an h^{th} object. As before, only one of the associated bids can be \overline{cg} . So, the sum of costs in terms of raising these bids is at most x . And, $E(x) \leq 1/n$ as well.

The remaining objects have v_{ih} above $\underline{cg} + 2mx$ but are not winning. But, then the deviation to v wins an extra object at price at most $\underline{cg} + x$, and raises the price by at most x on $m - 1$ units, for a net profit of $v_{ih} - \underline{cg} - mx$. The efficiency loss from i not winning object h is at most $v_{ih} - \underline{cg}$, which, given that $v_{ih} - \underline{cg} > 2mx$, is at most twice $v_{ih} - \underline{cg} - x$. So, on these objects, bidder's profits from the deviation are at least half of the efficiency loss on these units. Since costs from raising bids on other units are insignificant, it follows that the efficiency loss on these units is small since otherwise bidders will in aggregate have a profitable deviation. As the efficiency loss on other units is also small, we are done.

6 Extensions

6.1 One-sided Uniform-price Auctions

Swinkels (2001) considers large one-sided auctions with independent values and a little bit of “noise.” An example is if there is a small independent probability that each player sleeps through the auction. In the uniform price case, it is shown that with the noise, the impact that any given player has on the price grows small in expectation. But then, since “honest” bidding has a small effect on the price paid, it must also have little benefit in winning extra objects. This implies asymptotic efficiency (without a rate of convergence).

An easy extension to the arguments here shows that a one sided uniform price auction with z -independent values converges to efficiency at rate $1/n^{2-\alpha}$, even without noise. This paper thus significantly generalizes Swinkels (2001) for the uniform price case. The key is that here we think of “cost” as the impact on price *in circumstances where the player affecting the price cares*. This is a simpler object to bound, allowing both the greater generality, and fast convergence.¹⁹

6.2 Weaker Information Assumptions

We can weaken the information assumptions considerably. There is no problem if most players have considerably more knowledge about each other's values than z -independence allows. What counts (for convergence, rates are more delicate) is that from the point of view of a non-vanishing fraction of players, there are “lots” of players who he cannot predict precisely, and that NAG applies to this set of players.

¹⁹The stronger notion of vanishing impact is needed to prove results for discriminatory auctions, which are also analyzed in that paper.

6.3 Non-private Values

We can also weaken the assumption of private values somewhat. Assume that an ε fraction of the players have private values, and the remainder some sort of common. The arguments above show that over relevant ranges, the players with private values bid close to value. NAG implies that their bids are then closely packed almost surely. Thus, the impact of bids on price disappears for all players. But then, common value types should bid nearly “honestly” (their bid should nearly equal the expected value of object conditional on being pivotal). Working out such a model is left to future work.

References

- Al-Najjar, Nabil I., and Rann Smorodinsky (1997) “Pivotal Players and the Characterization of Influence,” *Journal of Economic Theory* **92** (2), 318-342.
- Chatterjee, Kalin, and William Samuelson (1983) “Bargaining Under Asymmetric Information,” *Operations Research*, **31** 835-851.
- Fudenberg, Drew, Markus Mobius, and Adam Szeidl (2003) “Existence of Equilibrium in Large Double Auctions,” mimeo, Harvard University.
- Gresik, Tom, and Mark Satterthwaite (1989) “The Rate of Which a Simple Market Becomes Efficient as the Number of Traders Increases: An Asymptotic Result for Optimal Trading Mechanisms,” *Journal of Economic Theory*, **48** 304–332.
- Jackson, Matthew O., Leo Simon, Jeroen M. Swinkels, and William Zame (2002) “Communication and Equilibrium in Discontinuous Games of Incomplete Information,” *Econometrica* **70** (5), 1711–1740.
- Mailath, George and Andrew Postlewaite (1990) “Asymmetric Information Bargaining Problems with Many Agents,” *Review of Economic Studies*, **57** 351–367.
- Jackson, Matthew O. and Jeroen M. Swinkels (2001) “Existence of Equilibrium in Single and Double Private Value Auctions,” mimeo John M. Olin School of Business.
- McLean, Richard, and Andrew Postlewaite (2002) “Informational Size and Incentive Compatibility,” *Econometrica* **70**, 2421-2454.
- Milgrom, Paul, (1979) “A Convergence Theorem for Competitive Bidding with Differential Information,” *Econometrica* **47**, 670-688.
- Milgrom, Paul and Robert J. Weber (1985) “Distributional Strategies for Games with Incomplete Information,” *Mathematics of Operations Research*, **10**, 619-632.
- Perry, Motty, and Philip J. Reny (2003) “Toward a Strategic Foundation for Rational Expectations Equilibrium,” mimeo, University of Chicago.
- Peters, Mike and Sergei Severinov (2002) “Internet Auctions with Many Traders,” Mimeo, UBC.

- Pesendorfer, Wolfgang and Jeroen M. Swinkels (1997) “The Loser’s Curse and Information Aggregation in Common Value Auctions,” *Econometrica* **65**, 1247-1282.
- Postlewaite, Andrew, and David Schmeidler (1986) “Implementation in Differential Information Economies,” *Journal of Economic Theory* **39**, 14-33.
- Rustichini, Aldo, Mark A. Satterthwaite and Steven R. Williams (1994) “Convergence to Efficiency in a Simple Market with Incomplete Information,” *Econometrica* **62** (1), 1041–1063.
- Satterthwaite, Mark A. and Steven R. Williams (1989) “The Rate of Convergence to Efficiency in the Buyer’s Bid Double Auction as the Market Becomes Large,” *Review of Economic Studies* **56**, 477-498.
- Satterthwaite, Mark A. and Steven R. Williams (2002) “The Optimality of a Simple Market Mechanism,” *Econometrica* **70** (5), 1841–1864.
- Shiryayev, Albert N. (1996) *Probability*, Springer-Verlag, Berlin, Heidelberg.
- Swinkels, Jeroen M. (2001) “Efficiency of Large Private Value Auctions,” *Econometrica* **69** (1), 37–68.
- Wilson, Robert (1977) “A Bidding Model of Perfect Competition,” *Review of Economic Studies* **66**, 511-518.
- Wilson, Robert (1985) “Incentive Efficiency of Double Auctions,” *Econometrica* **53**, 1101-1115.

7 Appendix

7.1 Proofs for Section 3.2

Proof of Lemma 1 Wlog, let $K = \{1, 2, \dots, |K|\}$. Let P_K and $P_{N \setminus K}$ be the marginals of P on K and $N \setminus K$ respectively, and let $P_{K \times N \setminus K}$ be the associated product measure. Fix a rectangular event $F_K = F_1 \cap F_2 \cap \dots \cap F_{|K|}$, where F_i only involves v_i .

$$\begin{aligned}
 \Pr(F_K | F_{N \setminus K}) &= \prod_{i=1}^{|K|} \Pr(F_i | F_{i+1} \cap \dots \cap F_{|K|} \cap F_{N \setminus K}) \\
 &\leq z^{-|K|} \prod_{i=1}^{|K|} \Pr(F_i | F_{i+1} \cap \dots \cap F_{|K|}) \text{ (using } z\text{-independence)} \\
 &= z^{-|K|} \Pr(F_K) = z^{-|K|} P_K(F_K).
 \end{aligned}$$

Analogously,

$$\Pr(F_K | F_{N \setminus K}) \geq z^{|K|} P_K(F_K).$$

These inequalities extend to any F_K in the product σ -algebra, as such a set is the limit of a countable union of rectangles. Thus

$$z^{-|K|} \Pr(F_K) \Pr(F_{N \setminus K}) \geq \Pr(F_K \cap F_{N \setminus K}) \geq z^{|K|} \Pr(F_K) \Pr(F_{N \setminus K}) \quad (9)$$

or equivalently

$$z^{-|K|}P_{K \times N \setminus K} \geq P \geq z^{|K|}P_{K \times N \setminus K}. \quad (10)$$

Let F_K and $F_{N \setminus K}$ be events about values *and* bids. Then

$$\begin{aligned} \Pr(F_K \cap F_{N \setminus K}) &= \int_{[0,1]^{|N|}} \Pr(F_K|v_K) \Pr(F_{N \setminus K}|v_{N \setminus K}) dP \\ &\leq z^{-|K|} \int_{[0,1]^{|N|}} \Pr(F_K|v_K) \Pr(F_{N \setminus K}|v_{N \setminus K}) dP_{K \times N \setminus K} \\ &= z^{-|K|} \int_{[0,1]^{|K|}} \Pr(F_K|v_K) dP_K \int_{[0,1]^{|N|-|K|}} \Pr(F_{N \setminus K}|v_{N \setminus K}) dP_{N \setminus K} \\ &= z^{-|K|} \Pr(F_K) \Pr(F_{N \setminus K}). \end{aligned}$$

The first integral is defined by the players' distributional strategies. The second line uses (10). The third line applies Fubini's Theorem. The final line integrates. Similarly $\Pr(F_K \cap F_{N \setminus K}) \geq z^{|K|} \Pr(F_K) \Pr(F_{N \setminus K})$ so (9) holds for all events.

Similarly, for rectangular events $F_{N \setminus K}$,

$$z^{-|N \setminus K|} \Pr(F_K) \Pr(F_{N \setminus K}) \geq \Pr(F_K \cap F_{N \setminus K}) \geq z^{|N \setminus K|} \Pr(F_K) \Pr(F_{N \setminus K}). \quad (11)$$

Combining,

$$z^{-a} \Pr(F_K) \Pr(F_{N \setminus K}) \geq \Pr(F_K \cap F_{N \setminus K}) \geq z^a \Pr(F_K) \Pr(F_{N \setminus K}) \quad (12)$$

Dividing through by $\Pr(F_{N \setminus K})$ gives (2).

Let X_K be a step function with values x^α on a finite partition $\{F^\alpha\}_{\alpha \in A}$ where each F^α is an event on bids and values in K . By the definition of conditional expectation $E(X_K|F_{N \setminus K}) = \sum_{\alpha \in A} x^\alpha \Pr(F^\alpha|F_{N \setminus K})$. Thus by (2)

$$E(X_K|F_{N \setminus K}) \leq z^{-a} \sum_{\alpha \in A} x^\alpha \Pr(F^\alpha) = z^{-a} E(X_K).$$

Analogously, $E(X_K|F_{K \setminus N}) \geq z^a E(X_K)$. As an arbitrary X_K is the limit of such step functions, (3) follows. \yenmark

7.2 Proofs for Section 3.3

Proof of Lemma 2 Wlog, let $K = \{1, 2, \dots, \kappa\}$. Define the Bernoulli process with κ independent trials with success probability \bar{p}_i in trial i . Let $x_i \in \{0, 1\}$ be the outcome of trial i and let $X^k = \sum_{i=1}^k x_i$. We claim that $X_K \equiv X^\kappa$ FO SD Q_K given $F_{N \setminus K}$. The proof is inductive. Let Q^k be the number of F_1, \dots, F_{k-1} that occur. Trivially, X^0 FO SD Q^0 , since both are identically 0. Suppose X^{k-1}

FOSD Q^{k-1} given $F_{N \setminus K}$. Then, for $r \in \{0, \dots, k\}$,

$$\begin{aligned} \Pr(Q^k \leq r \mid F_{N \setminus K}) &= \Pr(Q^{k-1} < r \mid F_{N \setminus K}) \\ &\quad + \Pr(F_k^c \mid \{Q^{k-1} = r\} \cap F_{N \setminus K}) \Pr(Q^{k-1} = r \mid F_{N \setminus K}) \\ &\geq \Pr(X^{k-1} < r \mid F_{N \setminus K}) + (1 - \bar{p}_k) \Pr(X^{k-1} = r \mid F_{N \setminus K}) \\ &= \Pr(X_K \leq r). \end{aligned}$$

The inequality uses z -independence and the inductive hypothesis.

Similarly, if Y_K is the number of successes in a Bernoulli process with success probabilities \underline{p}_i then given $F_{N \setminus K}$, Q_K FOSD Y_K .

We want a large-deviations inequality for the bounding Bernoulli processes. As X_K is a sum of non-identical independent Bernoulli trials, a slight alteration to the usual proof of Cramér's Theorem (e.g., Shirayev (1996) p.68) is necessary. Let $\pi = \frac{1}{\kappa} \sum_i \bar{p}_i$. Then, for any $\lambda > 0$,

$$\Pr\left(\frac{X_K}{\kappa\pi} > \phi\right) = \Pr\left(e^{\lambda X_K / \kappa\pi} \geq e^{\lambda\phi}\right) \leq \frac{E\left(e^{\lambda X_K / \kappa\pi}\right)}{e^{\lambda\phi}}$$

by Markov's inequality. Note also that $\Pr\left(\frac{X_K}{\kappa\pi} > \phi\right) = 0$ trivially when $\pi\phi \geq 1$.

Now, as X_K is a sum of independent random variables

$$\begin{aligned} Ee^{\lambda X_K / \kappa\pi} &= \prod_{i \in K} \left(1 - \bar{p}_i + \bar{p}_i e^{\lambda / \kappa\pi}\right) \tag{13} \\ &= \exp\left(\log \prod_{i \in K} \left(1 - \bar{p}_i + \bar{p}_i e^{\lambda / \kappa\pi}\right)\right) \\ &= \exp\left(\sum_{i \in K} \log\left(1 - \bar{p}_i + \bar{p}_i e^{\lambda / \kappa\pi}\right)\right) \\ &\leq \exp\left(\kappa \log\left(1 - \pi + \pi e^{\lambda / \kappa\pi}\right)\right) \end{aligned}$$

since $\log(1 - x + xe^{\lambda / \kappa\pi})$ is concave in x .

Thus,

$$\begin{aligned} \Pr\left(\frac{X_K}{\kappa\pi} > \phi\right) &\leq \exp\left[-\lambda\phi + \kappa \log\left(1 - \pi + \pi e^{\lambda / \kappa\pi}\right)\right] \tag{14} \\ &= \exp\left[-\kappa \left\{\frac{\lambda}{\kappa\pi} \phi\pi - \kappa \log\left(1 - \pi + \pi e^{\lambda / \kappa\pi}\right)\right\}\right] \\ &= \exp\left[-\kappa \{s\phi\pi - \log(1 - \pi + \pi e^s)\}\right], \end{aligned}$$

where $s \equiv \frac{\lambda}{\kappa\pi}$. Given that $\lambda > 0$ was arbitrary, this holds for all $s > 0$, and so

in particular for $s = \log\left(\frac{(1-\pi)\phi}{1-\pi\phi}\right)$ (this is positive, because $\pi\phi < 1$), yielding

$$\begin{aligned} \Pr\left(\frac{X_K}{\kappa\pi} > \phi\right) &\leq \exp\left[-\kappa\left\{\log\left(\frac{(1-\pi)\phi}{1-\pi\phi}\right)\phi\pi - \log\left(1-\pi + \pi e^{\log\left(\frac{(1-\pi)\phi}{1-\pi\phi}\right)}\right)\right\}\right] \\ &= \exp\left[-\kappa\left\{\phi\pi\log\phi + (1-\phi\pi)\log\left(\frac{1-\phi\pi}{1-\pi}\right)\right\}\right] \end{aligned}$$

As $\log x \geq (x-1)/x$ the second term in the braces is at least $\pi(1-\phi)$. Thus,

$$\Pr\left(\frac{X_K}{\kappa\pi} > \phi\right) \leq \exp[-\kappa\pi(\phi\log\phi + 1 - \phi)]. \quad (15)$$

Choosing $\phi = 3$,

$$\Pr(X_K > 3\kappa\pi) \leq e^{-\kappa\pi(\log 27 - 2)} \leq e^{-\kappa\pi}. \quad (16)$$

Note that $\frac{1}{z}E(Q_K) \geq \sum_{i \in K} \bar{p}_i = \kappa\pi$, and hence $\Pr(X_K > 3\kappa\pi) \geq \Pr(X_K > \frac{3}{z}E(Q_K))$. And, $\sum_{i \in K} \bar{p}_i \geq E(Q_K)$, and so $e^{-\kappa\pi} \leq e^{-E(Q_K)}$. Finally, X_K stochastically dominates Q_K . Taken together with (16), this implies

$$\Pr\left(Q_K > \frac{3}{z}E(Q_K) \mid F\right) \leq e^{-E(Q_K)}$$

giving (5).

The proof for Y_K is similar: Define $\pi = \sum_{i \in K} p_i$. Then, for any $\lambda < 0$, and $0 < \phi < 1$

$$\Pr\left(\frac{Y_K}{\kappa\pi} < \phi\right) = \Pr\left(e^{\lambda X_K / \kappa\pi} \geq e^{\lambda\phi}\right) \leq \frac{E(e^{\lambda X_K / \kappa\pi})}{e^{\lambda\phi}}.$$

The derivation of (13) and (14) is then as before, replacing \bar{p}_i by p_i and $\Pr\left(\frac{X_K}{\kappa\pi} > \phi\right)$ by $\Pr\left(\frac{Y_K}{\kappa\pi} < \phi\right)$. Note in particular that since $\lambda < 0$, $s \equiv \frac{\lambda}{\kappa\pi}$ can once again take on any positive value. Setting $s = \log\left(\frac{(1-\pi)\phi}{1-\pi\phi}\right)$ is once again valid, as $\phi < 1$, hence we arrive at the analog to (15):

$$\Pr\left(\frac{Y_K}{\kappa\pi} < \phi\right) \leq \exp[-\kappa\pi(\phi\log\phi + 1 - \phi)]. \quad (17)$$

Note that $\sum_{i \in K} p_i \geq zE(Q_K)$, so that $\Pr(Y_K < \phi zE(Q_K)) \leq \Pr(Y_K < \phi\kappa\pi)$ and $\exp[-\kappa\pi(\phi\log\phi + 1 - \phi)] \leq \exp[-zE(Q_K)(\phi\log\phi + 1 - \phi)]$. So,

$$\Pr(Q_K < \phi zE(Q_K)) \leq e^{-zE(Q_K)(\phi\log\phi + 1 - \phi)}. \quad (18)$$

Since $\frac{1}{3}\log\frac{1}{3} + 1 - \frac{1}{3} > 0.3$, (4) follows ¥

Proof of Corollary 1 Note that $\Pr(Q_K = 0 | F_{N \setminus K}) \leq \Pr(Q_K \leq \phi E(Q_K))$ for any $\phi > 0$. Equation 18 then gives

$$\begin{aligned} \Pr(Q_K = 0 | F_{N \setminus K}) &\leq \Pr(Q_K \leq \frac{z}{3}E(Q_K)) \\ &\leq e^{-zE(Q_K)(\phi\log\phi + 1 - \phi)} \\ &\leq e^{-z\Pr(Q_K \geq 1)(\phi\log\phi + 1 - \phi)} \end{aligned}$$

since $E(Q_K) \geq \Pr(Q_K \geq 1)$. As this holds for ϕ arbitrarily close to 0,

$$\begin{aligned} \Pr(Q_K \geq 1 | F_{N \setminus K}) &\geq 1 - e^{-z \Pr(Q_K \geq 1)} \\ &= \frac{1 - e^{-z \Pr(Q_K \geq 1)}}{\Pr(Q_K \geq 1)} \Pr(Q_K \geq 1) \end{aligned}$$

For $x \in (0, 1]$, $(1 - e^{-zx})/x$ is minimized at $x = 1$. ¥

7.3 Proofs for Section 3.4

Proof of Lemma 3 Partition $[0, 1]$ into $k \equiv \lceil n^{1-\alpha/4} \rceil$ intervals $\{I_\kappa\}$ of equal length (between $n^{1-\alpha/4}$ and $2n^{1-\alpha/4}$). Let $Q_B(I_\kappa)$ be the number of buyers with values in I_κ . Note that

$$Wn/k = Wn|I_\kappa| \geq E(Q_B(I_\kappa)) \geq wn|I_\kappa| = wn/k.$$

Let $E_{1\kappa} \equiv \{\frac{3}{z}Wn/k \geq Q_B(I_\kappa) \geq \frac{z}{3}wn/k\}$. By Lemma 2,

$$\Pr(E_{1\kappa}) \geq 1 - 2e^{-.3zn/k} \geq 1 - \frac{1}{n^5}$$

for n SL, since $n/k \rightarrow n^{\alpha/4}$. Similarly, let $Q_S(I_\kappa)$ be the number of sellers with values in I_κ , and define $E_{2\kappa} = \{\frac{3}{z}Wn/k \geq Q_S(I) \geq \frac{z}{3}wn/k\}$. Then, $\Pr(E_{2\kappa}) \geq 1 - \frac{1}{n^5}$ for n SL.

Then, $\mathcal{N} \equiv \cap_\kappa (E_{1\kappa} \cap E_{2\kappa})$. As this involves $2k \leq 2n^{1-\alpha/4}$ events,

$$\begin{aligned} \Pr(\mathcal{N}) &\geq 1 - 2 \frac{n^{1-\alpha/4}}{n^5} \\ &\geq 1 - 1/n^4 \end{aligned} \tag{19}$$

for n SL.

Finally, note that for n SL, any interval I of length at least $\frac{1}{n^{1-\alpha/3}}$ contains at least $k|I|/n - 2 \geq k|I|/2n$ elements of $\{I_\kappa\}$. So, in a normal realization,

$$Q_B(I) \geq \frac{k|I|}{2n} \frac{z}{3} wn/k = \frac{zw}{6} |I|.$$

Similarly, I intersects with at most $2k|I|/n$ elements of $\{I_\kappa\}$ and so $Q_B(I) \leq \frac{6}{z}W|I|$. The argument for $Q_S(I)$ is analogous. ¥

Proofs for Section 4.2

Proof of Lemma 5 If $\frac{E(V|T)}{n} \not\rightarrow 0$, then along a subsequence, $\Pr(V > \gamma n|T) > \gamma$ for some γ . Given $\{V > \gamma n\}$, if one selects $\frac{\gamma n}{2}$ of the buyers at random, the probability that none trades is at most $(1 - \gamma)^{\frac{\gamma n}{2}} \leq 1/8$ for n SL, and so there is a $7/8$ probability of at least one trader. Since this is true in expectation, it must be true for some particular set G_B of $\frac{\gamma n}{2}$ buyers. Similarly, there is a set G_S

of $\gamma n/2$ sellers such that conditional on $\{V > \gamma n\}$ at least one is a trader with probability $7/8$. Let $G \equiv G_S \cup G_B$, and let T_G be the event that at least one buyer and one seller in G trades. Then, $\Pr(T_G | \{V > \gamma n\}) \geq 1 - 2(1/8) = 3/4$. So, $\Pr(T_G \cap \{V > \gamma n\}) \geq 3/4 \Pr(V > \gamma n)$. As $T_G \subseteq T$,

$$\begin{aligned} \Pr(\{V > \gamma n\} | T_G) &= \frac{\Pr(T_G \cap \{V > \gamma n\})}{\Pr(T_G)} \\ &\geq \frac{3/4 \Pr(V > \gamma n)}{\Pr(T)} \\ &\geq 3\gamma/4. \end{aligned}$$

Since G has only $\frac{\gamma n}{2}$ buyers or sellers, $T_G \cap \{V > \gamma n\}$ implies that there are at least $\frac{\gamma n}{2}$ buyers and sellers trading in $N \setminus G$. Let X be this event. So, $\Pr(X | T_G) \geq 3\gamma/4$.

Let p^* be such that $\Pr(p \geq p^* | X \cap T_G) \geq \frac{1}{2}$ and $\Pr(p \leq p^* | X \cap T_G) \geq \frac{1}{2}$. Let Q_S be the number of sellers in $N \setminus G$ with $b_i \leq p^*$ and Q_B the number of buyers in $N \setminus G$ with $b_i \geq p^*$. Then,

$$E(Q_S | T_G) \geq \Pr(X \cap \{p \leq p^*\} | T_G) \frac{\gamma n}{2} \geq \frac{1}{2} \Pr(X | T_G) \frac{\gamma n}{2} \geq 3\gamma^2 n / 16,$$

and so

$$E(Q_S) = \sum_{i \in N_S \setminus G} \Pr(b_i \leq p^*) \geq z \sum_{i \in N_S \setminus G} \Pr(b_i \leq p^* | T_G) = z E(Q_S | T_G) = 3z\gamma^2 n / 16.$$

Thus by Lemma 2 $\Pr(Q_S = 0) \rightarrow 0$. Similarly $\Pr(Q_B = 0) \rightarrow 0$. But then, $\Pr(T) \rightarrow 1$. ¥

Proof of Proposition 1 Fix \mathcal{A}^n and a non-trivial equilibrium. Let $\phi_B \equiv \max_{N_B} b_i$ be the highest buy bid submitted and let $\phi_S \equiv \min_{N_S} b_i$ be the lowest sell bid. Note that $\Pr(\phi_B \geq x)$ is decreasing and continuous from the left. Similarly, $\Pr(\phi_S \leq x)$ is increasing and continuous from the right. Let $v^* \in [0, 1]$ have the property that for all $x \in [0, v^*)$, $\Pr(\phi_B \geq x) \geq \Pr(\phi_S \geq x)$, while for all $x \in (v^*, 1]$, $\Pr(\phi_B \geq x) \leq \Pr(\phi_S \geq x)$. Let

$$\delta \equiv \min \{ \Pr(\phi_B \geq v^*), \Pr(\phi_S \leq v^*) \}.$$

Note that $\Pr(\phi_B > v^*) \leq \delta$. This is trivial if $\Pr(\phi_B \geq v^*) = \delta$. If $\Pr(\phi_B \geq v^*) > \delta$, then $\Pr(\phi_S \geq v^*) = \delta$. But then since $\Pr(\phi_S \leq x)$ is continuous from the right,

$$\begin{aligned} \Pr(\phi_B > v^*) &= \lim_{v \downarrow v^*} \Pr(\phi_B \geq v) \\ &\leq \lim_{v \downarrow v^*} \Pr(\phi_S \geq v) \\ &= \Pr(\phi_S \geq v) = \delta. \end{aligned}$$

Analogously, $\Pr(\phi_S < v^*) \leq \delta$.

Assume that $\Pr(\phi_S \leq v^*) = \delta$. Then, $\Pr(T \cap \{p \leq v^*\}) \leq \Pr(\phi_S \leq v^*) = \delta$, while $\Pr(T \cap \{p > v^*\}) \leq \Pr(\phi_B > v^*) \leq \delta$. Similarly, if $\Pr(\phi_B \geq v^*) = \delta$, then, $\Pr(T \cap \{p < v^*\}) \leq \Pr(\phi_S < v^*) \leq \delta$, while $\Pr(T \cap \{p \geq v^*\}) \leq \Pr(\phi_B \geq v^*) = \delta$. So $\Pr(T) \leq 2\delta$.

Now, $\{\phi_S \leq v^*\} = \cup_{i \in N_S} \{b_i \leq v^*\}$. Hence, by Corrolary 1,

$$\begin{aligned} \Pr(\phi_S \leq v^* | F_{N_B}) &\geq (1 - e^{-z}) \Pr(\phi_S \leq v^*) \\ &\geq (1 - e^{-z}) \delta \end{aligned} \quad (20)$$

for any F_{N_B} . So,

$$\begin{aligned} \Pr(T) &\geq \Pr(\{\phi_B \geq v^*\} \cap \{\phi_S \leq v^*\}) \\ &\geq \Pr(\phi_S \leq v^* | \phi_B \geq v^*) \Pr(\phi_B \geq v^*) \\ &\geq (1 - e^{-z}) \delta^2. \end{aligned} \quad (21)$$

Assume that $v^* \leq 1/2$. (If not, the proof below applies, *mutatis mutandis*, to the sellers). Fix an arbitrary buyer i . Let $\phi_B^i \equiv \max_{N_B \setminus \{i\}} b_i$. Now,

$$\begin{aligned} \Pr(\phi_B^i < 2/3) &= 1 - \Pr(\phi_B^i \geq 2/3) \\ &\geq 1 - \Pr(\phi_B > v^*) \\ &\geq 1 - \delta. \end{aligned}$$

Let $J \equiv [5/6, 1]$. By Lemma 1,

$$\Pr(\phi_B^i < 2/3 | v_i \in J) \geq z(1 - \delta). \quad (22)$$

By (20),

$$\Pr(\phi_S \leq v^* | v_i \in J, \phi_B^i < v^*) \geq (1 - e^{-z}) \delta. \quad (23)$$

Let d_i be the deviation for i that whenever $v_i \in J$ and the original strategy specified a bid below v^* , he bids $2/3$ instead. Under this strategy, he wins an object with probability at least $\Pr(\phi_B^i < 2/3, \phi_S \leq v^*, v_i \in J)$, which by (22) and (23) is at least

$$\Pr(v_i \in J) z (1 - \delta) (1 - e^{-z}) \delta,$$

and earns at least $\frac{1}{6}$ when he does so. So,

$$B_i \geq \Pr(v_i \in J) z (1 - \delta) (1 - e^{-z}) \delta \frac{1}{6} - \pi_i,$$

where π_i is i 's expected equilibrium profit.

Summing across buyers, and applying Lemma 4,

$$z(1 - \delta) (1 - e^{-z}) \delta \frac{1}{6} \sum_{N_B} \Pr(v_i \in J) - \sum_{N_B} \pi_i \leq \Pr(T). \quad (24)$$

By A2 $\sum_{N_B} \Pr(v_i \in J) \geq \frac{1}{6}wn$ for $n \geq 6$. As the gains to a buyer from any given trade are at most 1, and V buyers trade,

$$\sum_{N_B} \pi_i \leq \Pr(T)E(V | T).$$

Substituting into (24) gives

$$z(1 - \delta)(1 - e^{-z})\delta \left(\frac{1}{6}\right)^2 wn - 2\delta \Pr(T)E(V | T) \leq \Pr(T).$$

Using $\Pr(T) \leq 2\delta$, and dividing through by $2\delta > 0$ gives

$$\frac{1}{72}wz(1 - \delta)(1 - e^{-z})n - E(V | T) \leq 1.$$

For this to hold for large n , either $(1 - \delta)$ must be close to 0, in which case, $\Pr(T) \geq (1 - e^{-z})\delta^2 \not\rightarrow 0$ (by (21)) or, $E(V | T)$ must grow like n . But then, by Lemma 5, $\Pr(T) \rightarrow 1$. \nexists

7.4 Proofs for Section 4.3

We will prove stronger results that will be useful when we turn to the multiple unit case. Fix an integer $m \geq 1$. Redefine \overline{ug} as the m^{th} bid above \overline{cg} . As before, let $ug \equiv (\overline{cg}, \overline{ug})$. When $m = 1$, we have the original case.

Proof of Lemma 6 Let $v \equiv E(|ug|)$, and let us show that for n SL, $v \leq \frac{1}{n^{1-\alpha_4}}$. Assume this is false along a subsequence. Then

$$\begin{aligned} v &= \Pr(\mathcal{N})E(|ug| | \mathcal{N}) + (1 - \Pr(\mathcal{N}))E(|ug| | \mathcal{N}^c) \\ &\leq \Pr\left(\mathcal{N} \cap \left\{|ug| > \frac{v}{2}\right\}\right) E\left(|ug| \mid \mathcal{N} \cap \left\{|ug| > \frac{v}{2}\right\}\right) \\ &\quad + \Pr\left(\mathcal{N} \cap \left\{|ug| \leq \frac{v}{2}\right\}\right) E\left(|ug| \mid \mathcal{N} \cap \left\{|ug| \leq \frac{v}{2}\right\}\right) + \frac{1}{n^4} \text{ (for } n \text{ SL)} \\ &= \Pr\left(\mathcal{N} \cap \left\{|ug| > \frac{v}{2}\right\}\right) E\left(|ug| \mid \mathcal{N} \cap \left\{|ug| > \frac{v}{2}\right\}\right) + \frac{v}{2} + \frac{1}{n^4}. \end{aligned}$$

So, for n SL

$$\Pr\left(\mathcal{N} \cap \left\{|ug| > \frac{v}{2}\right\}\right) E\left(|ug| \mid \mathcal{N} \cap \left\{|ug| > \frac{v}{2}\right\}\right) > \frac{v}{3}. \quad (25)$$

Consider $d_i(b_i, v_i) = \max\{b_i, v_i - \frac{v}{2}\}$. Consider $\mathcal{N} \cap \{|ug| > \frac{v}{2}\}$. Since $|ug| > \frac{v}{2}$, any buyer in the top half of $|ug|$ is a winner after d_i , and at most m were winners before (since buyers bid at most v_i , and by definition, there are only m bids in $(\overline{cg}, \overline{ug})$). By Lemma 3 (which applies since $|ug|/2 > v/2 > \frac{1}{2n^{1-\alpha_4}} > \frac{1}{n^{1-\alpha/3}}$

for n SL), the number of new winners is at least $w' \frac{1}{2} n |ug| - m \geq \frac{w'}{4} n |ug|$ for n SL. Each new winner earns at least $\frac{v}{2}$. So, using (25)

$$\begin{aligned} \sum B_i &\geq \frac{v w'}{2} \frac{1}{4} n E \left(|ug| \mid \mathcal{N} \cap \left\{ |ug| > \frac{v}{2} \right\} \right) \Pr \left(\mathcal{N} \cap \left\{ |ug| > \frac{v}{2} \right\} \right) \\ &\geq \frac{v w'}{2} \frac{1}{4} n \frac{v}{3}. \end{aligned}$$

But, by Lemma 4, $\sum B_i \leq v$, and so

$$\frac{v w'}{2} \frac{1}{4} n \frac{v}{3} \leq v$$

or

$$nv \leq \frac{24}{w'}.$$

For n SL, this contradicts $v \geq \frac{1}{n^{1-\alpha_4}}$. The argument for sellers is analogous. \nexists

Proof of Lemma 7 Assume $\Pr(|ug| > x) > \frac{1}{x^2 n^{2-\alpha_3}}$ along a subsequence where $x > \frac{1}{n^{1-\alpha_3/2}}$ (for smaller x , $\frac{1}{x^2 n^{2-\alpha_3}} \geq 1$, and the claim is vacuous). Then, for n SL

$$\Pr(\mathcal{N} \cap \{|ug| > x\}) > \Pr(\{|ug| > x\}) - \frac{1}{n^4} > \Pr(\{|ug| > x\})/2.$$

Consider $d_i(b_i, v_i) = \max\{b_i, v_i - \frac{x}{2}\}$. As before, given $\mathcal{N} \cap \{|ug| > x\}$, Lemma 3 implies that there are $w'nx - m > w'nx/2$ new winners, each earning $x/2$ (note that $x > \frac{1}{n^{1-\alpha_3/2}} > \frac{1}{n^{1-\alpha_3}}$, so Lemma 3 does apply). So,

$$\sum_{N_S} B_i \geq \Pr(|ug| > x) \frac{w'nx^2}{4}.$$

But, using Lemma 4 and Lemma 6, $\sum B_i \leq \frac{1}{n^{1-\alpha_4}}$. So,

$$\Pr(|ug| > x) \frac{w'nx^2}{4} \leq \frac{1}{n^{1-\alpha_4}},$$

Rearranging

$$\Pr(|ug| > x) \leq \frac{4}{w'n^{2-\alpha_4}}.$$

For n SL, this contradicts $\Pr(|ug| > x) > \frac{1}{x^2 n^{2-\alpha_3}}$. \nexists

Proof of Lemma 8 For buyer i , consider $d_i(b_i, v_i) \equiv \max\{b_i, v_i - x\}$. If $i \in L_B(x)$, then i wins an extra object and earns at least x . By Lemma 4 and Lemma 6,

$$xE(l_B(x)) \leq \sum B_i \leq E(|ug|) \leq \frac{1}{n^{1-\alpha_4}}$$

and so

$$E(l_B(x)) \leq \frac{1}{xn^{1-\alpha_4}}. \quad (26)$$

establishing the first claim. Now, note that in each realization,

$$SL_B(1/n) = \frac{1}{n}l_B(1/n) + \int_{1/n}^1 l_B(x)dx.$$

(This is easily seen by noting that $SL_B(1/n)$ is a consumer surplus calculation for demand curve $l_B(\cdot)$ up to demand $Q = l_B(1/n)$. Therefore, by Fubini's theorem

$$\begin{aligned} E(SL_B(1/n)) &= \frac{1}{n}E(l_B(1/n)) + \int_{1/n}^1 E(l_B(x))dx \\ &\leq \frac{1}{n^{1-\alpha_4}} + \int_{1/n}^1 \frac{1}{xn^{1-\alpha_4}}dx \text{ using (26) twice} \\ &= \frac{1}{n^{1-\alpha_4}}(1 + \log n) \\ &\leq \frac{1}{n^{1-\alpha_3}} \text{ (for } n \text{ SL)} \end{aligned}$$

which establishes the second claim. Repeat for sellers. \forall

7.5 Proofs for Section 4.4

Proof of Lemma 9 Suppose the lemma is false, so that there exists a sequence $\{n_t\}, \{x_t\}$ satisfying $n_t \rightarrow \infty$ and $x_t \geq 1/n_t^{1-\alpha_2/2}$ such that $\Pr(|cg| \geq x_t) \geq 1/n_t^{2-\alpha_2}x_t^2$ (the claim is vacuous for smaller x_t).

Step 1. Sparse Intervals.

Recall from the proof of Lemma 3 the partition of $[0, 1]$ into $k \equiv \lceil n^{1-\alpha/4} \rceil$ disjoint intervals $\{I_\kappa\}$ of equal length between $1/n^{1-\alpha/4}$ and $2/n^{1-\alpha/4}$. Let $M(I_\kappa)$ be the number of bids in I_κ . Let $\omega = zw'/24$. Say I_κ is *sparse* if $E(M(I_\kappa)) < \omega n^{\alpha/4}$. Let X be the set of sparse intervals. For $\kappa \in X$, let $E_{3\kappa} \equiv \{M(I_\kappa) \leq \frac{3}{z}\omega n^{\alpha/4}\}$ be the event that there are not "too many" bids in I_κ . For any given $\tau \in [0, 1]$ consider the process in which at step one, values and bids are drawn according to the distributional strategy μ , and at stage two, each bid is randomly and independently replaced by a bid in I_κ with probability τ . Let $M_\tau(I_\kappa)$ be the random variable given the number of bids in I_κ for this process. Clearly, $M_\tau(I_\kappa)$ stochastically dominates $M(I_\kappa)$ for any τ . Choose τ^* such that $E(M_{\tau^*}(I_\kappa)) = \omega n^{\alpha/4}$. Then, Lemma 2 implies that

$$\begin{aligned} \Pr(E_{3\kappa}) &\geq \Pr\left(M_{\tau^*}(I_\kappa) \leq \frac{3}{z}\omega n^{\alpha/4}\right) \\ &\geq 1 - e^{-\omega n^{\alpha/4}} \\ &\geq 1 - \frac{1}{n^5} \end{aligned}$$

for n SL.

For $\kappa \notin X$, let $E_{3\kappa} \equiv \{M(I_\kappa) \geq \frac{z}{3}\omega n^{\alpha/4}\}$ be the event that there are not “too few” bids in I_κ . Lemma 2 implies that for n SL

$$\Pr(E_{3\kappa}) \geq 1 - e^{-z\omega n^{\alpha/4}} \geq 1 - \frac{1}{n^5}.$$

Let $\mathcal{N}' \equiv \mathcal{N} \cap (\cap_\kappa E_{3\kappa})$. Arguing as in the proof of Lemma 3, for n SL

$$\begin{aligned} \Pr(\mathcal{N}') &\geq 1 - 3 \frac{n^{1-\alpha/4}}{n^5} \\ &\geq 1 - 1/n^4. \end{aligned} \tag{27}$$

Step 2. Sparse Regions and the Endpoints of Competitive Gaps. Assemble maximal groups of adjacent sparse intervals into sparse *regions*. Let $\{J^\lambda\}_{\lambda \in \Lambda}$ be the set of sparse regions that are longer than $\frac{x}{2}$. For n SL, $\frac{z}{3}\omega n^{\alpha/4} > 1$. So, given \mathcal{N}' , for all n SL, each non-sparse interval contains at least 1 bid and so cg cannot contain a non-sparse interval; cg can include at most a J^λ and parts of the two non-sparse intervals immediately adjacent. These two intervals, having length at most $2/n^{1-\alpha/4}$ become arbitrarily short compared to $x \geq 1/n^{1-\alpha_2/2}$. Hence, given \mathcal{N}' , and for n SL a competitive gap of length x must (a) have intersection of length at least $x/2$ with some J^λ , and (b) intersect at most one J^λ .

Let J_y^λ , $y \in [0, 1]$ be the point a y^{th} of the way up the interval J^λ . Our first lemma says that it is very unlikely that the competitive gap ends a long way from the end of a J^λ .

Lemma 10 *For all n SL*

$$\begin{aligned} \Pr(\overline{cg} \in \cup_\lambda [J_0^\lambda, J_{4/5}^\lambda]) &\leq \frac{1}{12n^{2-\alpha_2}x^2}, \\ \Pr(\underline{cg} \in \cup_\lambda [J_{1/5}^\lambda, J_1^\lambda]) &\leq \frac{1}{12n^{2-\alpha_2}x^2}. \end{aligned}$$

Proof Consider the event $\{\overline{cg} \in [J_0^\lambda, J_{4/5}^\lambda]\} \cap \mathcal{N}'$ for some $\lambda \in \Lambda$. Let $y \equiv J_1^\lambda - \overline{cg}$. As $\overline{cg} \in [J_0^\lambda, J_{4/5}^\lambda]$, $y/2 \geq x/10 \geq 1/n^{1-\alpha/4}$. So, By Lemma 3, the number of values in $[J_1^\lambda - y/2, J_1^\lambda]$ is at least $w'yn/2$. On the other hand, by Step 1, given \mathcal{N}' , each $I_\kappa \subseteq [J_1^\lambda - y/2, J_1^\lambda]$ includes at most $\frac{z}{3}\omega n^{\alpha/4} = \frac{z}{3}(zw'/24)n^{\alpha/4} = w'n^{\alpha/4}/8$ bids. For n SL, this implies that the number of bids in $[J_1^\lambda - y/2, J_1^\lambda]$ is at most $w'yn/4$ (by the same argument as in the proof of Lemma 3). Thus, given $\{\overline{cg} \in [J_0^\lambda, J_{4/5}^\lambda]\} \cap \mathcal{N}'$, there are at least $w'yn/2 - w'yn/4 = w'yn/4$ players with value in $[J_1^\lambda - y/2, J_1^\lambda]$ but bid below $\overline{cg} = J_1^\lambda - y$. But then,

$$SL_B(y/2) \geq \frac{w'yn}{4} \frac{y}{2}.$$

As $\overline{cg} \in [J_0^\lambda, J_{4/5}^\lambda]$, $y \geq x/5$. So, whenever $\{\overline{cg} \in [J_0^\lambda, J_{4/5}^\lambda]\} \cap \mathcal{N}'$,

$$\begin{aligned}
SL_B(x/10) &\geq SL_B(y/2) \\
&\geq \frac{w'nx^2}{200}.
\end{aligned}$$

And, for n SL,

$$\begin{aligned}
\Pr\left(\left\{\overline{cg} \in [J_0^\lambda, J_{4/5}^\lambda]\right\} \cap \mathcal{N}'\right) &\geq \Pr\left(\overline{cg} \in \cup_\lambda [J_0^\lambda, J_{4/5}^\lambda]\right) - \frac{1}{n^4} \\
&\geq \frac{1}{2} \Pr\left(\overline{cg} \in \cup_\lambda [J_0^\lambda, J_{4/5}^\lambda]\right).
\end{aligned}$$

Thus,

$$E\left(SL_B\left(\frac{x}{10}\right)\right) \geq \frac{w'nx^2}{400} \Pr\left(\overline{cg} \in \cup_\lambda [J_0^\lambda, J_{4/5}^\lambda]\right).$$

Now, for n SL $\frac{x}{10} > \frac{1}{n}$, and hence $SL_B(\frac{x}{10}) < SL_B(\frac{1}{n})$. However, $E(SL_B(1/n)) \leq \frac{1}{n^{1-\alpha_3}}$ by Lemma 8. Thus,

$$\frac{1}{n^{1-\alpha_3}} \geq \frac{w'nx^2}{400} \Pr\left(\overline{cg} \in \cup_\lambda [J_0^\lambda, J_{4/5}^\lambda]\right).$$

Rearranging,

$$\begin{aligned}
\Pr\left(\overline{cg} \in \cup_\lambda [J_0^\lambda, J_{4/5}^\lambda]\right) &\leq \frac{400}{n^{2-\alpha_3}x^2} \\
&\leq \frac{1}{12n^{2-\alpha_2}x^2}
\end{aligned}$$

for n SL. Repeat for sellers in the lower fifth to get the second claim. \yenmark

Step 4. Relative Probabilities of competitive and supporting gaps. Let $cg_\lambda \equiv \left\{cg \supseteq [J_{1/5}^\lambda, J_{4/5}^\lambda]\right\}$, and let $c_\lambda \equiv \Pr(cg_\lambda)$. Similar, let $lg_\lambda \equiv \left\{lg \supseteq [J_{1/5}^\lambda, J_{2/5}^\lambda]\right\}$, and $l_\lambda \equiv \Pr(lg_\lambda)$. Finally, let $ug_\lambda \equiv \left\{ug \supseteq [J_{3/5}^\lambda, J_{4/5}^\lambda]\right\}$, and $u_\lambda \equiv \Pr(ug_\lambda)$. Our next lemma shows that for some λ , c_λ is both non-trivial, and much larger than either l_λ or u_λ .

Lemma 11 *For n SL, there exists λ such that*

$$c_\lambda > \frac{1}{n^4}, \tag{28}$$

and such that

$$\frac{l_\lambda + u_\lambda}{c_\lambda} \leq \frac{4}{n^{\alpha_2 - \alpha_3}}. \tag{29}$$

Proof As $\Pr(|cg|^2 \geq x) > n^{\alpha_2-2}x^{-2}$ and $\Pr(\mathcal{N}') \geq 1 - 1/n^4$, for n SL $\Pr\left(\left\{|cg|^2 \geq x\right\} \cap \mathcal{N}'\right) \geq \frac{5}{6}n^{\alpha_2-2}x^{-2}$. By Lemma 10, the probability of a competitive gap in J^λ not including the middle $3/5$ is also less than $\frac{1}{6}n^{\alpha_2-2}x^{-2}$ for n SL. Therefore, $\sum_{\lambda \in \Lambda} c_\lambda \geq \frac{4}{6}n^{\alpha_2-2}x^{-2}$. Let Λ' denote the subset of regions with $c_\lambda > 1/n^4$. There are at most n regions. Thus,

$$\sum_{\lambda \in \Lambda \setminus \Lambda'} c_\lambda \leq \frac{n}{n^4} \leq \frac{1}{6}n^{\alpha_2-2}x^{-2}$$

for n SL. Thus

$$\sum_{\lambda \in \Lambda'} c_\lambda \geq \frac{1}{2}n^{\alpha_2-2}x^{-2}.$$

From Lemma 7, for n SL $\sum_{\lambda \in \Lambda'} l_\lambda + u_\lambda \leq 2n^{\alpha_3-2}x^{-2}$. Thus

$$\frac{\sum_{\lambda \in \Lambda'} l_\lambda + u_\lambda}{\sum_{\lambda \in \Lambda'} c_\lambda} \leq \frac{2n^{\alpha_3-2}x^{-2}}{\frac{1}{2}n^{\alpha_2-2}x^{-2}} = \frac{4}{n^{\alpha_2-\alpha_3}}.$$

for n SL. Since this is true on average, it must be true for at least one $\lambda \in \Lambda'$. \forall

In what follows, we refer to a λ for which Lemma 11 holds. Let $\tilde{c} \equiv \Pr(cg \supseteq [J_{2/5}^\lambda, J_{3/5}^\lambda])$ be the probability of a competitive gap including the middle fifth of J^λ . We will show that \tilde{c} is close to 1. The idea is that the only way to have c_λ be large relative to $u = u_\lambda$ and $l = l_\lambda$ will be for players to almost always get it almost right.

Let $U_i \equiv \{b_i \geq J_{4/5}^\lambda\}$ be the event that i bids up and $\tilde{U}_i \equiv \{b_i \geq J_{2/5}^\lambda\}$ be the event that i bids weakly up. Symmetrically, let $D_i \equiv \{b_i \leq J_{1/5}^\lambda\}$ and $\tilde{D}_i \equiv \{b_i \leq J_{3/5}^\lambda\}$ be the events that i bids down and weakly down. Let $p_i \equiv \Pr(U_i)$, $\tilde{p}_i \equiv \Pr(\tilde{U}_i)$, $q_i \equiv \Pr(D_i)$ and $\tilde{q}_i \equiv \Pr(\tilde{D}_i)$. Order the players so that $\tilde{q}_1 \leq \tilde{q}_2 \leq \dots \leq \tilde{q}_{2n}$.

Step 5. A preliminary inequality. Define

$$A_{-j}^i \equiv \bigcap_{j' > i, j' \neq j} D_{j'}$$

as the event that all players after i not including j bid down. Then, for any event F involving $1, 2, \dots, i$,

$$\begin{aligned} \Pr(A_{-j}^i | F) &\leq \prod_{j' > i, j' \neq j} \Pr(D_{j'} | D_{j'-1}, \dots, D_{i+1}, F) \\ &= \exp \left[\sum_{j' > i, j' \neq j} \log \Pr(D_{j'} | D_{j'-1}, \dots, D_{i+1}, F) \right] \\ &\leq \exp \left[\sum_{j' > i, j' \neq j} (\Pr(D_{j'} | D_{j'-1}, \dots, D_{i+1}, F) - 1) \right] \quad (\text{since } \log x \leq x - 1) \end{aligned} \tag{30}$$

$$\begin{aligned}
&\leq \exp \left[- \sum_{j' > i, j' \neq j} \Pr(D_{j'}^c \mid D_{j'-1}, \dots, D_{i+1}, F) \right] \\
&\leq e^{-z \sum_{j' > i, j' \neq j} 1 - q_{j'}} \text{ (by } z\text{-independence)} \\
&\leq e^{-z \sum_{j' > i, j' \neq j} p_{j'}} \text{ (since } 1 - q_{j'} \geq p_{j'}) \\
&\leq e^{-z(-1 + \sum_{j' > i} p_{j'})}.
\end{aligned}$$

Step 6. Two Bounds. Recall that $cg_\lambda \equiv \{cg \supseteq [J_{1/5}^\lambda, J_{4/5}^\lambda]\}$. Let cg_λ^{ij} , $i < j$, be the event cg_λ where i and j are the two highest indexed players for whom U_i holds. Let $c_\lambda^{ij} = \Pr(cg_\lambda^{ij})$. Let F^i be the event that $U_{j'}$ holds for $j' = i$ and for $n-2$ other $j' \in \{1, \dots, i\}$, while $D_{j'}$ holds for all other $j' \in \{1, \dots, i\}$. Then,

$$\begin{aligned}
c_\lambda^{ij} &= \Pr(F^i \cap A_{-j}^i \cap U_j) \\
&= \Pr(U_j \mid F^i \cap A_{-j}^i) \Pr(F^i \cap A_{-j}^i) \\
&= \frac{\Pr(\tilde{D}_j \mid F^i \cap A_{-j}^i)}{\Pr(\tilde{D}_j \mid F^i \cap A_{-j}^i)} \Pr(U_j \mid F^i \cap A_{-j}^i) \Pr(F^i \cap A_{-j}^i) \\
&\leq \frac{p_j}{z^2 \tilde{q}_j} \Pr(\tilde{D}_j \mid F^i \cap A_{-j}^i) \Pr(F^i \cap A_{-j}^i) \\
&= \frac{p_j}{z^2 \tilde{q}_j} \Pr(\tilde{D}_j \cap F^i \cap A_{-j}^i).
\end{aligned}$$

Let ug_λ^i be the event ug_λ , where i is the last player to bid up, and let $u_\lambda^i = \Pr(ug_\lambda^i)$. When $\tilde{D}_j \cap F^i \cap A_{-j}^i$ holds, i is the last player to bid up and in total $n-1$ players bid up while the rest bid weakly down. Thus, $\{\tilde{D}_j \cap F^i \cap A_{-j}^i\} \subseteq ug_\lambda$, and so $\Pr(\tilde{D}_j \cap F^i \cap A_{-j}^i) \leq u_\lambda^i$. The previous equation thus implies

$$c_\lambda^{ij} \leq \frac{p_j}{z^2 \tilde{q}_j} u_\lambda^i. \quad (31)$$

Another bound on c_λ^{ij} comes from (30):

$$c_\lambda^{ij} \leq \Pr(A_{-j}^i) \leq e^{-z(-1 + \sum_{j' > i} p_{j'})}. \quad (32)$$

Step 7. Up and Down Players. We next show that for all n SL, $\Pr(\tilde{D}_i) \leq \frac{1}{4}$ for $i \leq n$, and $\Pr(\tilde{U}_i) \geq \frac{1}{4}$ for $i > n$.

Consider the first claim. Suppose that $\tilde{q}_n > 0$ (if $\tilde{q}_n = 0$ the result is immediate). Let $c_\lambda^i \equiv \sum_{j > i} c_\lambda^{ij}$ be the probability that cg_λ occurs, where i is the second last up player. Let $i^* \leq n$ be the last index with the property that $\sum_{j' > i} p_{j'} > n^{(\alpha_2 - \alpha_3)/2}$. Then,

$$\begin{aligned}
\sum_{i \leq i^*} c_\lambda^i &= \sum_{i \leq i^*} \sum_{j > i} c_\lambda^{ij} \\
&\leq n^2 e^{-z(-1 + \sum_{j' > i} p_{j'})} \text{ (using (32))} \\
&\leq n^2 e^{-z(-1 + n^{(\alpha_2 - \alpha_3)/2})} \\
&\leq \frac{1}{2n^4} \text{ (for } n \text{ SL)} \\
&\leq \frac{1}{2} c_\lambda \text{ (using (28))}
\end{aligned}$$

So, as $c_\lambda = \sum_{i \geq n-1} c_\lambda^i$, for n SL,

$$\begin{aligned}
\frac{1}{2} c_\lambda &\leq \sum_{i > i^*, i \geq n-1} c_\lambda^i \\
&\leq \sum_{i > i^*, i \geq n-1} \sum_{j > i} \frac{p_j}{z^2 \tilde{q}_j} u_\lambda^i \text{ (using (31))} \\
&\leq \frac{1}{z^2 \tilde{q}_n} \sum_{i > i^*, i \geq n-1} \sum_{j > i} p_j u_\lambda^i \text{ (since } \tilde{q}_i \text{ is increasing)} \\
&\leq \frac{n^{(\alpha_2 - \alpha_3)/2}}{z^2 \tilde{q}_n} \sum_{i > i^*, i \geq n-1} u_\lambda^i \text{ (by choice of } i^*) \\
&\leq \frac{n^{(\alpha_2 - \alpha_3)/2}}{z^2 \tilde{q}_n} u_\lambda \\
&\leq \frac{n^{(\alpha_2 - \alpha_3)/2}}{z^2 \tilde{q}_n} \frac{4}{n^{\alpha_2 - \alpha_3}} c_\lambda \text{ (by (29))} \\
&= \frac{4}{z^2 \tilde{q}_n} \frac{1}{n^{(\alpha_2 - \alpha_3)/2}} c_\lambda.
\end{aligned}$$

Comparing the first and last expressions, $\tilde{q}_n \rightarrow 0$, and so in particular, $q_i \leq 1/4$ all $i \leq n$ for all n SL.

If the players are ordered so that \tilde{p}_i increases, this argument can be repeated considering events in which $n-1$ of the first i players bid down and the others bid up. Thus there are n players for which $\tilde{p}_i \rightarrow 0$. As $\tilde{p}_i + \tilde{q}_i \geq 1$ these players are disjoint from players $1, \dots, n$, and so must be the players $\{n+1, \dots, 2n\}$.

Let $R \equiv \cap_{i \leq n} U_i \cap_{i > n} D_i = c g_\lambda^{n-1, n}$ be the event that all the players bid according to their type (and a competitive gap occurs).

Step 8. A lower bound for $\Pr(R)$: We already know that $\sum_{i \leq i^*, j > i} c_\lambda^{ij} \leq \frac{1}{2} c_\lambda$ for n SL. We will show that for n SL $\sum_{i > i^*, j > n+1} c_\lambda^{ij} \leq \frac{1}{4}$. Since R is the only

event left, it would then follow that $\Pr(R) \geq \frac{c_\lambda}{4}$. So, as in Step 7 note that

$$\begin{aligned}
\sum_{i>i^*, j \geq n+1} c_\lambda^{ij} &\leq \sum_{i>i^*, j \geq n+1} \frac{p_j}{z^2 \tilde{q}_j} u_\lambda^i \text{ (using (31))} \\
&\leq \frac{1}{z^2 \tilde{q}_{n+1}} \sum_{i>i^*} u_\lambda^i \sum_{j>i} p_j \text{ (note the } n+1 \text{)} \\
&\leq \frac{n^{(\alpha_2 - \alpha_3)/2}}{2z^2} \sum_{i>i^*} u_\lambda^i \text{ (by choice of } i^*, \text{ and since } \tilde{q}_{n+1} \cong 1) \\
&\leq \frac{n^{(\alpha_2 - \alpha_3)/2}}{2z^2} u_\lambda \\
&\leq \frac{n^{(\alpha_2 - \alpha_3)/2}}{2z^2} \frac{1}{n^{\alpha_2 - \alpha_3}} c_\lambda \text{ (by (29))} \\
&= \frac{1}{2z^2} \frac{1}{n^{(\alpha_2 - \alpha_3)/2}} c_\lambda \\
&\leq \frac{1}{4} c_\lambda \text{ (for } n \text{ SL).}
\end{aligned}$$

Step 9. A Persistent Competitive Gap. Let $\tilde{R} \supset R$ be the event that all the players get it nearly right — the first n players are not bidding below $J_{3/5}^\lambda$ and the others are not bidding above $J_{2/5}^\lambda$. For $i > n$, define R_{-i} to be the event that all players except i play according to type. If R_{-i} occurs and player i bids weakly up, then there will be a $lg \supset [J_{1/5}^\lambda, J_{2/5}^\lambda]$. Thus,

$$\begin{aligned}
u_\lambda &\geq \sum_{i>n} \Pr(R_{-i}) \Pr(\tilde{U}_i | R_{-i}) \\
&\geq z \Pr(R) \sum_{i>n} \tilde{p}_i \text{ (by } z\text{-independence and since } R \subseteq R_{-i}) \\
&\geq \frac{zc_\lambda}{4} \sum_{i>n} \tilde{p}_i
\end{aligned}$$

Since $\frac{u_\lambda}{c_\lambda} \rightarrow 0$,

$$\sum_{i>n} \tilde{p}_i \rightarrow 0.$$

Arguing symmetrically,

$$\sum_{i \leq n} \tilde{q}_i \rightarrow 0.$$

With the ordering described in Step 2, \tilde{R} occurs when the first n players do not bid weakly down and the last n players do not bid weakly up, that is, $\tilde{R} = (\cap_{i \leq n} \tilde{D}_i^c) \cap (\cap_{i > n} \tilde{U}_i^c)$. Thus,

$$\Pr(\tilde{R}) \geq 1 - \sum_{i \leq n} \tilde{q}_i - \sum_{i > n} \tilde{p}_i \rightarrow 1.$$

Step 10. **A Contradiction.** When $\tilde{R} \cap \mathcal{N}'$ occurs, $[J_{2/5}^\lambda, J_{3/5}^\lambda]$ is in the interior of cg (players $1, \dots, n$ bid strictly above $J_{3/5}$, and other players strictly below $J_{2/5}$). And since the probability of trade is bounded away from 0, and since $\Pr(\tilde{R}) \rightarrow 1$, there is at least one buyer in $\{1, \dots, n\}$ and at least one seller in $\{n+1, \dots, 2n\}$.

Let p^* be the expected price conditional on \tilde{R} . Either $p^* \leq J_{1/2}^\lambda$, or $p^* \geq J_{1/2}^\lambda$. Wlog, assume $p^* \geq J_{1/2}^\lambda$. Let x^λ be the length of J^λ . By construction, $x^\lambda \geq \frac{1}{2}x \geq \frac{1}{2n^{1-\alpha_2/2}}$

Assume first that $J_1^\lambda \geq 1 - 3x^\lambda$. Consider any buyer in $\{1, \dots, n\}$. A bid of $J_{2/5}^\lambda$ wins whenever \tilde{R} occurs, and forces the price to at most $J_{2/5}^\lambda$. So, conditional on \tilde{R} , the buyer's expected gain from lowering the price is at least $J_{1/2}^\lambda - J_{2/5}^\lambda \geq \frac{x^\lambda}{6}$. On the other hand, when \tilde{R} does not occur, he may go from being a winner to a loser. But, for this to happen, it must be that $\underline{cg} \geq J_{2/5}^\lambda$. But then i 's lost profit is at most $1 - \underline{cg} \leq 4x^\lambda$. Since $\Pr(\tilde{R}) \rightarrow 1$,

$$\Pr(\tilde{R}) \frac{x^\lambda}{6} - (1 - \Pr(\tilde{R}))4x^\lambda$$

is eventually positive, and we have a contradiction.

Assume $J_1^\lambda < 1 - 3x^\lambda$. Given \mathcal{N}' , the number of buyers with value in $(J_1^\lambda + 2x^\lambda, J_1^\lambda + 3x^\lambda)$ is at least $w'nx^\lambda$. But, by Lemma 8 for n SL

$$E(\#U(x^\lambda)) \leq \frac{1}{n^{1-\alpha_4}x^\lambda}.$$

It follows that for n SL, at least half the buyers with value in $(J_1^\lambda + 2x^\lambda, J_1^\lambda + 3x^\lambda)$ trade conditional on $\tilde{R} \cap \mathcal{N}'$ (and so bid above $J_{3/5}^\lambda$) Consider the deviation that any buyer with value in $(J_1^\lambda + 2x^\lambda, J_1^\lambda + 3x^\lambda)$ and bid above J^λ bids $J_{2/5}^\lambda$ instead. Given $\tilde{R} \cap \mathcal{N}'$, this gains the buyer at least $x^\lambda/6$. Given \mathcal{N}' , the number of players for in $(J_1^\lambda + 2x^\lambda, J_1^\lambda + 3x^\lambda)$ is at least $w'nx^\lambda$ and at most $W'nx^\lambda$. So, given $R \cap \mathcal{N}'$, the expected sum of gains is at least $\frac{w'nx^\lambda}{2} \frac{x^\lambda}{6}$. The loss from such a buyer going from being a winner to a loser is again at most $4x^\lambda$. Given $\mathcal{N}' \setminus \tilde{R}$, there are at most $W'nx^\lambda$ such buyers. In \mathcal{N}'^c , the worst case is that all n buyers are in $(J_1^\lambda + 2x^\lambda, J_1^\lambda + 3x^\lambda)$. So, the expected sum of losses is at most

$$\begin{aligned} & \Pr(\mathcal{N}' \setminus \tilde{R})W'nx^\lambda 4x^\lambda + \Pr(\mathcal{N}'^c)n4x^\lambda \\ & \leq (1 - \Pr(\tilde{R}))W'nx^\lambda 4x^\lambda + \frac{1}{n^4}n4x^\lambda \end{aligned}$$

and thus, since the deviation cannot be profitable,

$$\Pr(\tilde{R}) \frac{w'nx^\lambda}{2} \frac{x^\lambda}{6} \leq (1 - \Pr(\tilde{R}))W'nx^\lambda 4x^\lambda + \frac{1}{n^4}n4x^\lambda.$$

Dividing both sides by $n(x^\lambda)^2$,

$$\begin{aligned}
\Pr(\tilde{R}) \frac{w'}{12} &\leq 4W' \left(1 - \Pr(\tilde{R})\right) + \frac{4}{n^2 x^\lambda} \\
&\leq 4W' \left(1 - \Pr(\tilde{R})\right) + \frac{8}{n^2 x} \quad (\text{since } x^\lambda \geq \frac{1}{2}x) \\
&\leq 4W' \left(1 - \Pr(\tilde{R})\right) + \frac{8}{n^2 \frac{1}{n^{1-\alpha_2/2}}} \\
&= 4W' \left(1 - \Pr(\tilde{R})\right) + \frac{8}{n^{1+\alpha_2/2}}.
\end{aligned}$$

The LHS goes to $w'/12$, while the RHS goes to 0, a contradiction. \nexists

7.6 Proofs for Section 5

Let x be the random variable $\overline{sg}_m - \underline{cg}$. In an m unit demand/supply setting, this is the maximum impact of raising a buyer's bid vector on price. Let p be the price. We will show that in expectation buyers achieve within $\frac{1}{2n^{1-\alpha}}$ of the consumer surplus if they can price take at p . A symmetric argument applies to sellers. But, the sum of consumer and producer surplus at an arbitrary p is at least as large as the surplus at the Walrasian price. So, this both establishes that the market achieves within $1/n^{1-\alpha}$ of the efficient surplus and that price must be asymptotically Walrasian (else the market achieves more than the feasible surplus, a contradiction). Finally, from NAG and NAA, expected feasible surplus grows like n , and the result follows.

Consider the truth-telling deviation $d_i(b_i, v_i) = v_i$, remembering that v_i and b_i are now vectors in $[0, 1]^m$. Let W be the set of ih , $i \in N_B$ that are allocated an object. Let $SL_{ih} = 0$ if i wins an object h , and let $SL_{ih} = \max[v_{ih} - p, 0]$ otherwise. So, SL_{ih} gives the loss in consumer surplus compared to taking at p from i not winning object h .

In any given realization, think about moving from b_i to v_i one bid at a time, starting from b_{i1} . Let \hat{C}_{ih} be the cost to i from raising bid h in terms of raising the price paid on units already won, and \hat{B}_{ih} the profit to i of winning an extra unit.

If $v_{ih} < \underline{cg}$, then raising b_{ih} to v_{ih} is irrelevant to both p and the allocation. Hence, $\hat{B}_{ih} - \hat{C}_{ih} = 0$. And, since $v_{ih} < p$, $SL_{ih} = 0$.

If $v_{ih} \in [\underline{cg}, \overline{sg} + 2mx]$, then raising b_{ih} to v_{ih} may raise the price on units already won by as much as x . So, $\hat{B}_{ih} - \hat{C}_{ih} \geq -(m-1)x$. And, since $v_{ih} \in [\underline{cg}, \overline{sg} + 2mx]$ and $p \geq \underline{cg}$, $SL_{ih} \leq (2m+1)x$. In any normal realization, the number of such ih is at most $Kn x$ for some $K < \infty$. In a non-normal realization, there are at most nm values in this range. Hence, the expected number of such values is at most

$$\begin{aligned}
&\left(1 - \frac{1}{n^4}\right) Kn x + \frac{1}{n^4} nm \\
&\leq Kn x + \frac{m}{n^3}
\end{aligned}$$

Consider $ih \in W$ such that $v_{ih} > \overline{sg} + 2mx$. In any realization, at most 1 player who is winning an object is also in a position to affect the price by changing the associated bid. And, the impact of that bid on price is at most x . Hence,

$$\sum_{\{ih \in W | v_{ih} > \overline{sg} + 2mx\}} \hat{C}_{ih} \leq x$$

and

$$\sum_{\{ih \in W | v_{ih} > \overline{sg} + 2mx\}} SL_{ih} = 0.$$

Finally, consider $ih \notin W$ such that $v_{ih} > \overline{sg} + 2mx$. Then, by deviating to $b_{ih} = v_{ih}$, i raises the price on at most $m - 1$ previous units by at most x . But, i also wins an extra object at price at most \overline{sg} . So,

$$\hat{B}_{ih} - \hat{C}_{ih} \geq v_{ih} - \overline{sg} - (m - 1)x \geq \frac{v_{ih} - p}{2} = \frac{SL_{ih}}{2}.$$

Since we are in equilibrium

$$\begin{aligned} 0 &\geq E \left(\sum_{ih} \hat{B}_{ih} - \hat{C}_{ih} \right) & (33) \\ &= E \left(E \left(\sum_{\{ih \notin W | v_{ih} > \overline{sg} + 2mx\}} \hat{B}_{ih} - \hat{C}_{ih} \middle| x \right) \right) + E \left(E \left(\sum_{\{ih \in W | v_{ih} > \overline{sg} + 2mx\}} \hat{B}_{ih} - \hat{C}_{ih} \middle| x \right) \right) \\ &\quad + E \left(E \left(\sum_{\{ih | v_{ih} \in [\underline{cg}, \overline{sg} + 2mx]\}} \hat{B}_{ih} - \hat{C}_{ih} \middle| x \right) \right) \\ &\geq E \left(E \left(\sum_{\{ih | v_{ih} > \overline{sg} + 2mx\}} \frac{SL_{ih}}{2} \middle| x \right) \right) - E(x) - E \left(E \left(\sum_{\{ih | v_{ih} \in [\underline{cg}, \overline{sg} + 2mx]\}} (m - 1)x \middle| x \right) \right) \\ &\geq E \left(E \left(\sum_{\{ih | v_{ih} > \overline{sg} + 2mx\}} \frac{SL_{ih}}{2} \middle| x \right) \right) - E(x) - E \left(\left(Knx + \frac{m}{n^3} \right) (m - 1)x \right) \\ &\geq E \left(E \left(\sum_{\{ih | v_{ih} > \overline{sg} + 2mx\}} \frac{SL_{ih}}{2} \middle| x \right) \right) - 2E(x) - K''nE(x^2) \text{ (for } n \text{ SL)} \end{aligned}$$

Let H be the cumulative for x . By Lemmas 7 and 9, $H(x) \leq \frac{1}{n^2 - \alpha_2 x^2}$ for all

$x > \frac{1}{n}$. Hence,

$$\begin{aligned}
nE(x^2) &= \int_0^1 nx^2 dH(x) \\
&= \int_0^1 n2x[1 - H(x)]dx \\
&\leq 0 + \int_0^{1/n} 2nxdx + \int_{1/n}^1 2nx \frac{1}{n^{2-\alpha_2}x^2} dx \\
&= nx^2 \Big|_0^{1/n} + \frac{2}{n^{1-\alpha_2}} \int_{1/n}^1 \frac{1}{x} dx \\
&\leq \frac{1 + 2 \log n}{n^{1-\alpha_2}} \\
&\leq \frac{1}{n^{1-\alpha_1}} \text{ (for } n \text{ SL)}.
\end{aligned}$$

And, $E(x) \leq \frac{1}{n^{1-\alpha_1}}$ as well (a simple integration by parts). So, (33) yields

$$\begin{aligned}
E \left(E \left(\sum_{\{ih|v_{ih} > \overline{sg} + 2mx\}} SL_{ih} \Big| x \right) \right) &\leq 2 \left(\frac{1}{n^{1-\alpha}} + K'' \frac{1}{n^{1-\alpha}} \right) \\
&= K''' \frac{1}{n^{1-\alpha}}
\end{aligned}$$

But then,

$$\begin{aligned}
E \left(\sum_{ih} SL_{ih} \right) &= E \left(E \left(\sum_{\{ih|v_{ih} > \overline{sg} + 2mx\}} SL_{ih} \Big| x \right) \right) + E \left(E \left(\sum_{v_{ih} \in [\underline{cg}, \overline{sg} + 2mx]} SL_{ih} \Big| x \right) \right) \\
&\leq K''' \frac{1}{n^{1-\alpha_1}} + E \left(Knx + \frac{m}{n^3} \right) (2m + 1)x \\
&\leq K''' \frac{1}{n^{1-\alpha_1}} + \frac{2m + 1}{n^3} xmE(x) + K'''' E(nx^2) \\
&\leq \frac{1}{2n^{1-\alpha}} \text{ for } n \text{ SL}
\end{aligned}$$

Arguing analogously for sellers, $E(\sum_{ih} SL_{ih}) \leq \frac{1}{2n^{1-\alpha}}$. Hence, the expected sum of consumer and producer surplus is within $1/n^{1-\alpha}$ of that achieved by the Walrasian outcome, and we are done. ¥