Effect of Optical Defocus on Detection and Recognition of Vanishing Optotype Letters in the Fovea and Periphery.

Nilpa Shah<sup>1</sup>, Steven C. Dakin<sup>1</sup>, Roger S. Anderson<sup>1,2</sup>

- NIHR Biomedical Research Centre for Ophthalmology, Moorfields Eye Hospital NHS Foundation Trust & UCL Institute of Ophthalmology, London, UK.
- Vision Science Research Group, University of Ulster, Coleraine, Northern Ireland, UK.

**Correspondence:** N. Shah, UCL Institute of Ophthalmology, 11-43 Bath Street, London EC1V 9EL. Tel: +447949 705 384, E-mail: nilpa.shah@ucl.ac.uk

## Word Count: 4561

**Grant Information:** Supported by a Fight for Sight studentship, by Moorfields Special Trustees and by an award from the NIHR Biomedical Research Centre for Ophthalmology, Moorfields Eye Hospital & UCL Institute of Ophthalmology, London.

**PURPOSE.** Vanishing Optotypes (VO's) are pseudo high-pass letters whose mean luminance matches the background so that they 'vanish' when recognition acuity threshold is reached in the fovea. We determined the effect of increasing blur on acuity for these optotypes and conventional letters, in both foveal and extrafoveal viewing.

**METHODS.** Detection and recognition thresholds were determined separately for each of the 26 letters of both a conventional and VO alphabet, both in the fovea and at 10 degrees in the horizontal temporal retina, under varying degrees of positive dioptric blur.

**RESULTS.** In the fovea, detection and recognition thresholds were similar for individual VO's, increased steadily with blur, and separated somewhat at higher levels of defocus (3D). While the recognition thresholds for VO's changed on average by 0.28 logMAR/dioptre, those for conventional letters changed more rapidly by 0.35 logMAR/dioptre. In the periphery, recognition thresholds were significantly higher than detection thresholds for the 'Vanishing' Optotypes at 0D blur; both thresholds increased steadily thereafter, converging as blur increased. Peripheral recognition acuity displayed a loss of only 0.09 logMAR/dioptre. In both the fovea and periphery, the inter-letter variation in recognition acuity was much lower for VO's than conventional letters (0.04 vs. 0.09 logMAR).

**CONCLUSIONS.** Outside the fovea, high-pass 'Vanishing' Optotypes display significant differences in their detection and resolution thresholds up to +7D blur, with a logMAR/diopter loss of a quarter that of the fovea. The lower inter-letter legibility differences indicate that VO letters may be better stimuli from which to design clinical letter charts.

### INTRODUCTION.

Any test of visual acuity should deliver accurate, precise and repeatable measurements in order to reliably identify significant change in performance resulting from either abnormality or therapy. There is increasing evidence that conventional letters charts, including logMAR charts, are failing to do so, displaying test-retest variability of between 0.06 and 0.19 log units even for normal, focused eyes,<sup>1-11</sup> with this variability increasing significantly with the presence of optical defocus<sup>12</sup> or retinal disease.<sup>13</sup>

"Vanishing Optotype" (VO) targets, first described by Howland et al.,<sup>14</sup> have a pseudo-high-pass design and are typically constructed of a dark core surrounded by light edges, or vice versa, which results in the mean luminance of the letter matching that of the background. Their construction means that, like gratings, size thresholds for detection and recognition of VO's are closely matched in the fovea when the letters are well focused and, unlike conventional letters, the characters 'vanish' almost as soon as the recognition limit is exceeded. However, the relative legibility of Vanishing Optotype letters across the whole alphabet has not yet been examined.

Conventional letters make good stimuli to detect defocus in the fovea because their rich spatial frequency spectra make them especially vulnerable to the effects of phase reversals associated with defocus, something which has much less effect on grating acuity.<sup>15</sup> However, the relative vulnerability of Vanishing Optotypes to the effects of optical defocus remains unknown.

Many subjects for whom we require measures of letter acuity, such as patients with age-related macular degeneration (AMD), have lost foveal function and view extrafoveally. Although detection and recognition thresholds for VO's are similar in the fovea, numerous studies of peripheral acuity, employing gratings with the same mean luminance as their surround (as with VO letters), have found a significant difference between detection and resolution thresholds outside the fovea, indicating that peripheral grating resolution is limited, not by optics, but by retinal ganglion cell sampling density.<sup>16</sup> Further evidence for the sampling limited nature of peripheral grating acuity comes from the observation that resolution acuity remains robust to optical blur up to 3-4 dioptres.<sup>17,18</sup> In this study, we also wished to determine the differences in detection and recognition performance for individual letters of VO design in peripheral vision under differing levels of defocus.

Our goals were thus to (i) determine VO legibility on an individual letter basis under foveal and extrafoveal presentation, (ii) examine the effect of optical defocus on acuity for VO compared to conventional letters (both foveally and extrafoveally) and (iii) determine if differences in detection and recognition performance exist peripherally for individual VO letters under these differing levels of defocus.

#### **METHODS**

Ethical approval for this study was obtained from the UCL Research Ethics Committee and all procedures adhered to the tenets of the declaration of Helsinki. Testing using both conventional and Vanishing Optotype letters was predominantly conducted on two experienced psychophysical observers (NS and RSA), with no ocular abnormalities and corrected visual acuities of 6/5 or better. As a control, we also employed a naïve observer when testing under zero-blur conditions.

The Vanishing Optotypes were constructed with an inner black 'core' flanked by a white border of half the width of the central section. This resulted in a target with the same mean luminance as the background (53cd/m<sup>2</sup>) and an on-screen contrast of 98%. The appearance of several different Vanishing Optotypes can be viewed down the left side of Figure 1.

Insert Figure 1 about here. Full size.

For the conventional letter measurements, black letters were presented on a white background (B/W) of luminance 113 cd/m<sup>2</sup>, thus yielding the same on-screen contrast as the Vanishing Optotypes (98%). Since this higher background luminance could potentially introduce an additional influence on acuity estimates relative to the VO test conditions, control measurements were also made with white letters on the same grey background (W/G) as the Vanishing Optotypes (53cd/m<sup>2</sup>) in the zero defocus condition. While this afforded the same individual retinal illuminance as for VO letters, it resulted in a lower on-screen contrast of 35%.

All letters had the same outline form. For both stimulus types, the letter height and width were 5 times the 'stroke width', which in the case of the Vanishing Optotypes incorporated both the dark middle bar and its two white flanks. All optotype stimuli were generated using MATLAB v7.6 (Mathworks Inc) and were presented on a high-resolution (1280 x 1024 pixels) Dell Trinitron P992 CRT monitor (Dell Corp. Ltd,

Bracknell, Berkshire, UK) driven by an Apple Macintosh computer. True 14-bit contrast resolution was achieved using a Bits++ video processor (Cambridge Research Systems, Ltd., Rochester, UK).

Scaling of stimuli was achieved using the OpenGL capabilities of the computer's built-in graphics card (ATI Radeon X1600). This (bilinear interpolation) procedure allowed us to display stimuli of arbitrary size with sub-pixel resolution while retaining accurate representation of their (balanced) luminance structure.

Foveal and peripheral visual acuity measurements were made monocularly in the right eyes of both subjects, using both conventional and Vanishing Optotypes. All peripheral measurements were made at 10 degrees eccentricity in the nasal field. This eccentricity was chosen, on the one hand, because the peripheral refractive error of both subjects differed little from that of the fovea at that location and, on the other hand, to prevent larger, defocused letters from encroaching on the fovea. In addition, it is not uncommon to find patients with age-related macular degeneration (AMD) displaying a preferred retinal locus around this eccentricity.<sup>19</sup>

The refractive error was determined and corrected prior to the start of each testing session using trial lenses which were employed for both on- and off-axis testing. Little difference was found between the two locations (subject NS -0.25D fovea, 0D periphery; subject RSA +0.50DS for both fovea and periphery). All foveal testing was conducted under low room illumination at a viewing distance of 8 m, at which the letters could vanish without pixilation effects. All peripheral testing was conducted under the same conditions at 1.6 m; at this near distance the screen subtended 11.6 x 9.8 degrees and one pixel subtended 0.55 minutes of arc.

Each subject underwent the following tests: detection and recognition of conventional optotypes (both B/W and W/G) and detection and recognition of Vanishing Optotypes, under foveal and peripheral viewing conditions i.e. twelve test conditions in total. Detection and recognition tasks were conducted in separate runs.

All 26 different alphabet letters were used for each test condition. In the detection tasks using the method of limits, a letter was initially displayed at sub-threshold size and the observer moved the computer mouse to progressively increase the letter-size until the optotype was just visible, which they indicated by clicking the mouse button. In the recognition task, the observer was again requested to increase the size of the optotype, but this time was required to indicate the letter-size when the letter identity was just discernable. The observer then verbally reported the letter identity to the experimenter; if the letter identity was incorrectly reported, the observer was required to continue increasing the size until able to correctly identify it. Viewing time was not restricted and five presentations were made, in a randomly interleaved order, for each letter of the alphabet under each condition.

We chose to employ an ascending method of limits (MOL) rather than forced choice procedures for nearly all of the testing. The reasons for this were several. Firstly, employing forced-choice procedures in an individual-letter comparison experiment requires twenty-six different interleaved staircases resulting in more than eight hundred presentations for each run. When this is undertaken for both detection (2 intervals per presentation) and recognition, and for each of eight blur conditions in both the fovea and periphery, this procedure inevitably introduces enormous variability as a consequence of observer fatigue. Secondly, it becomes more difficult to quantitatively compare detection and recognition thresholds because the thresholds would be measured under different forced choice conditions (2 interval forced choice for detection, 26 alternative forced choice for resolution) yielding a different guess rate and potentially boosting performance for the smaller number of alternatives (detection). Thirdly, in previous experiments<sup>20</sup> we observed that circular letters, such as O and Q, are most commonly confused with each other and can behave as something of a subset within a 26 alternative test. If, under forced choice conditions, an observer is able to determine that a letter is round, but unable to precisely identify which one, s/he will often bias towards a particular letter (e.g. O) thus artificially boosting 'performance' for that letter. Under MOL procedures, rather than forcing a decision, the letter must be increased in size until the observer can more confidently report its identity, yielding a truer threshold for that individual letter. Finally, lacking previous similar high-pass letter studies, we had no prior expectations of which of the 26 letters, if any, might display differences between detection and recognition acuity, thereby affecting our criterion. However, in order to assess the possibility that criterion-based methods may permit bias of the results in non-naive observers, for comparison and control purposes, we employed forced-choice reversal staircase procedures (QUEST) in the zero-blur condition for one of the trained observers.

In the second stage of the study, both observers repeated the task with levels of defocus increasing, from their initial refraction, in +1D steps to a maximum of +3D under foveal viewing conditions and +7D under peripheral viewing conditions (beyond +3D of foveal blur the letters became too large to be generated on the screen at the test distance employed). Detection and recognition thresholds were measured in the same way as described above for the Vanishing Optotypes, but only recognition thresholds were measured for B/W conventional letters.

The average detection and recognition size was recorded and converted to 'logMAR' where, for the Vanishing Optotypes, the 'stroke width' was taken to include both the central dark bar and its white flanks.

## RESULTS

Figure 2 displays both the detection and recognition values for the Vanishing Optotypes, and the recognition values for conventional letters, presented under different levels of blur in both the fovea (2a) and at 10 degrees (2b). Each point is the average across the 26 letters and both subjects. Error bars represent the SD of the 26 letter thresholds under each blur condition, averaged for both subjects.

Performance for recognition of conventional letters was significantly 'better' than either detection or recognition performance for the Vanishing Optotypes at zero blur (-0.01 logMAR vs. 0.11 & 0.14 logMAR; p<0.0001, one-way ANOVA). Under foveal viewing conditions, detection and recognition thresholds for Vanishing Optotypes were similar, increasing steadily with blur and separating somewhat at higher levels of defocus. However, while the recognition thresholds for Vanishing Optotypes changed by 0.28 logMAR/dioptre on average, those for conventional letters changed more rapidly by 0.35 logMAR/dioptre so that, after +1D blur, performance was 'worse' for conventional letters than Vanishing Optotypes (p<0.01, paired t-test). The same pattern was observed for both subjects.

At 10 degrees in the periphery (Figure 2b), recognition thresholds for the conventional letters is again lower than for the Vanishing Optotypes at 0D blur and increases roughly in parallel with it as blur increases, converging somewhat at +7D.

Recognition thresholds were significantly higher than detection thresholds for the Vanishing Optotypes at 0D blur (0.85 vs. 0.56 logMAR; p<0.0001, one-way ANOVA test). Thresholds for recognition appear to be little affected until around +1D blur but increases steadily thereafter, resulting in a change of approximately 0.09 logMAR/dioptre over the full 7D blur range.

The lower part of Figure 3a displays the detection and recognition values for the individual black-on-white (B/W) conventional–form letters in the fovea for 0D blur (average of both subjects). Unsurprisingly, and owing to the large difference between the mean luminance of the letters and the background, both subjects displayed significantly lower detection thresholds than recognition thresholds (mean -0.59 logMAR for detection and -0.01 logMAR for recognition; error bars represent the SE of the five threshold measurements for each letter). It can also be seen that there are significant between-letter threshold differences for both detection and recognition, with detection displaying significantly less variation than recognition (SD 0.04 vs. 0.09 logMAR, mean of both subjects). For recognition, the highest thresholds were observed for the circular letters (CGOQ), which seemed to behave as a separate subset. For the W/G conventional letters (not plotted), performance was qualitatively and quantitatively very similar to that for the higher contrast B/W letters, the only observable difference being slightly increased thresholds for letter detection than in the B/W case (-0.47 rather than -0.59 logMAR).

We already noted that the Vanishing Optotypes display very similar average detection and recognition performance in the fovea, but the lower part of Figure 3b indicates that this also applies on an individual letter basis. As a consequence, while the SD of the between-letter differences for detection is similar to conventional letters (0.05 vs. 0.04 logMAR), the variation for recognition is much lower with VO's (0.04 vs. 0.09

logMAR, mean of all letters and both subjects). Results under forced-choice staircase conditions, and for the naïve observer, were qualitatively similar but are not shown. Looking at the results under peripheral viewing, the conventional B/W letters (Figure 4a) display even larger differences between detection (mean -0.08 logMAR) and recognition (mean 0.63 logMAR) than in the fovea. Between-letter threshold differences display the same degree of variation as in the fovea for detection (SD 0.03 logMAR) and slightly greater variation for recognition (SD 0.12 logMAR), with the circular letters again displaying the highest recognition thresholds. The W/G letters again displayed very similar qualitative and quantitative performance to B/W and are not plotted.

Unlike in the fovea, the Vanishing Optotypes exhibit quite large differences between detection and recognition thresholds in the periphery, for all letters and for both subjects (Figure 4b, lower part). The magnitude of this difference again varied with the letter in question. While the performance variation between letters was higher than in the fovea for these optotypes (SD 0.07 logMAR for both detection and recognition), performance variation for recognition was notably lower than for conventional letters at the same location. Acuities and SD's for all tasks at 0D blur are summarized in Table 1. Again, results under forced-choice staircase conditions, and for the naïve observer, were qualitatively similar but are not shown.

The upper half of Figure 3a displays the letter recognition thresholds for individual B/W conventional letters under the maximum +3D blur conditions in the fovea. While performance fell by an average of 1.05 logMAR under +3D blur, the between-letter variability displays a very similar pattern to zero blur.

The upper half of 3b displays both detection and recognition thresholds for the individual Vanishing Optotypes in the fovea. Interestingly, it can be seen that the

detection and recognition thresholds increasingly separate with increased blur, most noticeably for the circular letters (CGOQ), which again begin to behave as a separate subset.

The upper parts of Figure 4 (a & b) display the corresponding results for the periphery with the maximum +7D of blur. For conventional letters (4a), recognition performance on average decreased by 0.83 logMAR over this range, with a between-letter standard deviation very similar to 0D blur (0.11 logMAR). The circular letters (CGOQ) again appear to behave as a separate subset.

For the Vanishing Optotypes (4b), both detection and resolution performance declined with blur, but detection more so. The 0.29 logMAR average difference observed between the two thresholds at 0D blur narrowed to 0.13 logMAR at +7D blur.

### DISCUSSION

In a previous study,<sup>20</sup> we found that acuity thresholds measured with Vanishing Optotype letters are less variable than corresponding measures made with conventional letters. Furthermore, we found that acuity estimates with VO's vary less with the number of available alternatives. In explaining this finding we proposed that attenuating the low spatial frequency components renders the letters more equally resolvable, and this in turn results in lower variability in acuity thresholds based on letter discrimination.

In the current study we sought to separately measure the detection and recognition acuity thresholds for these optotypes on an individual basis, under both foveal and extrafoveal viewing. This allows us to, firstly, determine if between-letter performance is in fact less variable than for conventional letters and, secondly, to determine if detection and recognition thresholds continue to be closely similar outside the fovea for these letters, and at different levels of blur.

It is apparent from Figure 2 that, in agreement with our previous study, foveal acuity thresholds for VO's are significantly larger than for conventional letters in the focused condition. This was not surprising given that most of the low frequency information has been removed from the stimuli, requiring the letter to increase in size in order for the visual system to utilise the higher frequency content. Surprising however, is the observation that, while threshold letter size for VO recognition is larger in the focused condition, it becomes slightly, but significantly, smaller than for conventional letters as defocus increases above 1 diopter. Why might this happen? Since the high-pass VO letters contain less information at low frequencies, in the focused state they must necessarily become initially larger so that these higher *object* frequencies become lower in terms of *retinal* frequency in order for the visual system to resolve them and recognise the letter. When letters become progressively defocused, it is the higher retinal frequencies that first start to phase-reverse, which some previous studies have claimed results in masking of the lower frequencies, making the letters increasingly difficult to resolve.<sup>21</sup> However, other computational studies have found that this is not necessarily the case. Akutsu et al.<sup>22</sup> reported that removing the spectrum above the first cut-off of the optical transfer function (OTF) had little effect on defocused letter VA. Ravikumar et al.<sup>23</sup> found that, in the presence of positive spherical aberration, the impact of phase correction on letter acuity depended on the sign of the defocus. For positive defocus (as in the present study), the impact on VA was not significantly different for standard, phase-rectified or low-pass filtered defocus, leading them to conclude that the primary cause of acuity loss for positive blur was contrast reduction;

but that was for conventional letters. For VO letters it may be that phase-reversal of the higher frequencies, which make up the lighter edges of the stimuli, causes the edges to become darker and results in the letter strokes effectively becoming thicker and the letter more discriminable. However, it is beyond the scope of the present study to fully determine why performance for heavily defocused VO letters is actually 'better' than for conventional letters.

The results found in the periphery provide interesting additions to our knowledge from previous studies. Unlike in the fovea, there is a significant difference between detection and recognition performance for VO letters, indicating that these letters do not, in fact, vanish extrafoveally. This is in agreement with our previous studies using high-pass targets with lower numbers of alternatives.<sup>24,25</sup> The 'vanishing' adjective may, therefore, be somewhat of a misnomer under these conditions. These letters behave in a partially similar manner to that found in previous studies that employed peripheral gratings with the same mean luminance as their surround. A superiority of detection acuity over recognition/resolution acuity for targets with the same mean luminance as their background, is strong evidence that the *resolution* task is, for the majority of letters, limited by retinal sampling rather than the eye's optics.

However, we found the effect of optical defocus on these letters to display both similarities and differences to the effects found with gratings.<sup>17,18</sup> While the effect of optical defocus is substantially less outside the fovea for both the conventional and VO letters, displaying a logMAR/diopter loss of only about a quarter of that observed in the fovea, the recognition acuity is not quite so robust to blur with VO's as observed in previous studies using gratings. This may be because, unlike for gratings, retinal sampling is not the only significant limiting factor involved in peripheral viewing. While grating appearance remains largely veridical under optical phase

reversal, the increasingly spurious appearance of the letters with increasing defocus renders the letter more ambiguous to the underlying retinal ganglion cell mosaic. Also, while the previous studies found that grating detection and resolution performance are identical by around 3-4 diopters defocus, there remains a difference between the two thresholds for high-pass letters all the way up to 7D (Figure 2b); again this is likely because the spuriously defocused letter permits the eye to detect the presence of contrast that may not be sufficiently veridical to resolve. This would seem to also occur somewhat in the fovea where, as defocus increases, the detection and recognition thresholds of the 'vanishing' optotype begin to separate somewhat (Figure 2a).

Looking at the individual letters more closely, we found that, under zero defocus (lower half of Figure 3a), there are considerable differences in the recognition thresholds (grey squares) for different conventional letters, with an inter-letter range of 0.40 logMAR. The result was the same for W/G as for B/W letters. This variation was even greater under forced choice conditions (0.99 logMAR). When considering only the Sloan letters (vertical arrows) the legibility range reduces to 0.22 logMAR (SD 0.08 logMAR, mean of subjects), but this is still much greater than the desired 0.1 logMAR difference between lines on a conventional logMAR chart. These legibility differences are similar to those found in previous studies,<sup>26</sup> and further calls into question the widespread acceptance that the Sloan letters are closely equal in legibility. As expected, the circular letters (CGOQ) display the lowest acuities and appear to behave as a separate subset, or even as a pair of subsets (e.g. OQ and CG), likely owing to their close similarity to each other and strong dissimilarity to the other letters. The current findings support the notion that, when the within-line recognition differences become greater than the between-line differences, acuity measurement

variability results. This was discussed by McMonnies & Ho<sup>27</sup> who found that chance combinations of easy or difficult letters can lead to significant line-difficulty variation. By reducing the within-line differences, we could reasonably expect testretest variability to improve. For this reason, it may be possible to choose a different set of letters to the Sloan ones, with more closely similar legibility. Grimm et al.<sup>28</sup> have summarised some previous studies that have attempted to do this.

The Vanishing Optotypes (Figure 3b) however, in addition to displaying closely similar detection and recognition thresholds for all letters, display much lower between-letter variation in recognition under zero defocus (lower section); range 0.15 logMAR across all 26 letters which reduces slightly further to 0.13 logMAR for the Sloan set. Our previous study<sup>20</sup> also found lower letter acuity measurement variability for VO letters, suggesting the reason to be the smaller inter-letter legibility differences. If a test-chart's within-line legibility difference is greater than its between-line legibility difference this will limit the reliability of acuity measurement and consequently its ability to register subtle alterations in visual acuity that can signify changes in the disease state. While the purpose of this study was not to assess test-retest variability, which is a property of a specifically constructed instrument, it is possible from the error bars on Figure 2a to observe the lower variability of acuity measurements with Vanishing Optotypes compared to conventional letters. VO letters may thus be more appropriate targets from which to construct acuity charts. Interestingly, the circular VO letters (CGOQ) do not so much behave as a separate subset under zero defocus, but increasingly do so as defocus increases (upper part of Figure 3b). Why should this happen? These four letters are most commonly confused with each other and the information required to distinguish them most likely lies within the higher frequencies; as these higher frequencies are progressively attenuated by the low-pass filtering effects of the blur lenses they will become increasingly difficult to tell apart. The inclusion of these VO letters on a clinical test-chart could thus increase acuity measurement variability under higher levels of defocus.

In peripheral viewing, the inter-letter recognition differences were again substantial for the conventional letters, the circular letters again behaving as a separate subset (Figure 4a) but these differences were again much smaller for VO's (Figure 4b). However, unlike in the fovea, we found significant, and often substantial, differences between detection and recognition thresholds for all of the VO characters (Figure 4b). This difference between detection and recognition thresholds, and the relative robustness to the effects of optical defocus, points towards a sampling limit for recognition, but not detection, of VO letters in peripheral vision.

This has implications for tests such as High-Pass Resolution Perimetry (HRP)<sup>29</sup>, which assumes that the detection and resolution limits remain the same for 'Vanishing' Optotypes in the periphery. While only one optotype is employed by HRP (a ring) it can be seen by observing our results for the 'O' in Figure 4b that its detection and recognition thresholds are significantly different. However, our finding of blur-resistant, potentially sampling-limited recognition performance for 'vanishing' optotypes outside the fovea points towards a patient friendly *in vivo* measure of localized ganglion cell density; in effect a true letter resolution perimetry test.

Finally, in conditions like age-related macular degeneration (AMD), where there is loss of foveal photoreceptors or even a central scotoma necessitating extrafoveal viewing, these optotypes may no longer 'vanish' when recognition fails and the targets remain visible for some time after. This situation, resulting in a discrepancy between detection and recognition, may even be a useful sign of early AMD. Further

work on clinical patients is required to better determine the clinical usefulness of high-pass letters in the detection and diagnosis of diseases such as AMD and glaucoma. In addition, other possibly confounding age-related optical factors such as straylight and lens yellowing may also affect the detection and/or recognition of VO's and these should be investigated in future studies.

## ACKNOWLEDGEMENTS

Supported by a Fight for Sight studentship, by Moorfields Special Trustees and by an award from the NIHR Biomedical Research Centre for Ophthalmology, Moorfields Eye Hospital & UCL Institute of Ophthalmology, London.

### REFERENCES

- 1) Elliott DB, Sheridan M. The use of accurate visual acuity measurements in clinical anti-cataract formulation trials. *Ophthalmic Physiol Opt.* 1988;8:397-401.
- Lovie-Kitchin JE. Validity and reliability of visual acuity measurements. *Ophthalmic Physiol Opt.* 1988;8:363-370.
- Reeves BC, Wood JM, Hill AR. Vistech VCTS 6500 charts within and between – session reliability. *Optom Vis Sci.* 1991;68:728-737.
- Brown B, Lovie-Kitchin J. Repeated visual acuity measurement: establishing the patient's own criterion for change. *Optom Vis Sci.* 1993;70:45-53.
- Vanden Bosch ME, Wall M. Visual acuity scored by the letter-by-letter or probit methods has lower retest variability than the line assignment method. *Eye* 1997;11:411-417.
- Arditi A, Cagenello R. On the statistical reliability of letter-chart visual acuity measurements. *Invest Ophthalmol Vis Sci.* 1993;34:120-129.
- Bailey IL, Bullimore MA, Raasch TW, Taylor HR. Clinical grading and the effects of scaling. *Invest Ophthalmol Vis Sci.* 1991;32:422-432.

- Hazel CA, Elliott DB. The dependency of logMAR visual acuity measurements on chart design and scoring rule. *Optom Vis Sci.* 2002;79:788-792.
- Rosser DA, Laidlaw DA, Murdoch IE. The development of a "reduced logMAR" visual acuity chart for use in routine clinical practice. *Br J Ophthalmol.* 2001;85(4):432-436.
- 10) Rosser DA, Cousens SN, Murdoch IE, Fitzke FW, Laidlaw DA. How sensitive to clinical change are ETDRS logMAR visual acuity measurements? *Invest Ophthalmol Vis Sci.* 2003;44:3278-3281.
- 11) Laidlaw DA, Tailor V, Shah N, Atamian S, Harcourt C. Validation of a computerised logMAR visual acuity measurement system (COMPlog): comparison with ETDRS and the electronic ETDRS testing algorithm in adults and amblyopic children. *Br J Ophthalmol.* 2008;92(2):241-244
- 12) Rosser DA, Murdoch IE, Cousens SN. The effect of optical defocus on the testretest variability of visual acuity measurements. *Invest Ophthalmol Vis Sci.* 2004;45:1076-1079.
- Patel PJ, Chen FK, Rubin GS, Tufail A. Intersession repeatability of visual acuity scores in age-related macular degeneration. *Invest Ophthalmol Vis Sci.* 2008;49:4347-4352.

- 14) Howland B, Ginsburg A, Campbell F. High-pass spatial frequency letters as clinical optotypes. *Vision Res.* 1978;18:1063-1066.
- 15) Thorn F, Schwartz F. Effects of dioptric blur on Snellen and grating acuity. *Optom Vis Sci.* 1990;67(1):3-7.
- 16) Anderson R.S. The psychophysics of glaucoma: improving the structure/function relationship. *Prog Ret Eye Res.* 2006;25:79-97.
- 17) Anderson RS. The selective effect of optical defocus on detection and resolution acuity in peripheral vision. *Curr Eye Res.* 1996;15(3):351-353.
- 18) Wang YZ, Thibos LN, Bradley A. Effects of refractive error on detection acuity and resolution acuity in peripheral vision. *Invest. Ophthalmol. Vis. Sci.* 1997;38:2134–2143.
- 19) Rubin GS, Feely M. The Role of Eye Movements During Reading in Patients with Age-Related Macular Degeneration (AMD). *Neuro-ophthalmology*. 2009;33(3):120-126.
- Shah N, Dakin SC, Redmond T, Anderson RS. Vanishing Optotype acuity: repeatability and effect of the number of alternatives. *Ophthalmic Physiol Opt.* 2001;31:17-22.

- 21) Thorn F, Schwartz F. Effects of dioptric blur on Snellen and grating acuity. *Optom Vis Sci.* 1990;67(1):3-7.Anderson RS. The selective effect of optical defocus on detection and resolution acuity in peripheral vision. *Curr Eye Res.* 1996;15(3):351-353.
- 22) Akutsu H, Bedell HE, Patel SS. Recognition thresholds for letters with simulated dioptric blur. *Optom Vis Sci.* 2000;77:524–530.
- 23) Ravikumar S, Bradley A, Thibos L. Phase changes induced by optical aberrations degrade letter and face acuity. *J Vis.* 2010;10(14):1-12.
- 24) Anderson RS, Ennis FA. Foveal and peripheral thresholds for detection and resolution of vanishing optotype tumbling E's. *Vision Res.* 1999;39:4141-4144.
- 25) Demirel S, Anderson RS, Haggerty K, Thibos LN. Detection and resolution of high-pass (vanishing-optotype) letters in central and peripheral vision. *American Acad Optom Annual Meeting* (abstract).1998
- 26) Alexander KR, Xie W, Derlacki DJ. Visual acuity and contrast sensitivity for individual Sloan letters. *Vision Res.* 1997;37(6):813-819.
- 27) McMonnies CW, Ho A. Letter legibility and chart equivalence. Ophthalmic Physiol Opt. 2000;20(2):142-152.

- 28) Grimm W, Rassow B, Wesemann W, Saur K, Hilz R. Correlation of optotypes with the Landolt ring – a fresh look at the comparability of optotypes. *Optom Vis Sci.* 1994;71(1):6-13.
- 29) Frisen L. High-pass resolution targets in peripheral vision. *Ophthalmol.* 1987;94:1104-1108.

### **FIGURE LEGENDS**

Figure 1. The left side displays different 'vanishing' optotype letters under different gammas. While covering one eye, the reader should observe the figure at a distance of around 4m (or less if not fully corrected optically) and choose the letter with the appropriate gamma so that it disappears into the background. Next, at a distance of 50cm, the reader should observe this same letter extrafoveally while fixating the cross to its right (to achieve a viewing angle of 10 degrees). It should be observed that, while difficult to resolve, the letter is clearly detectable.

Figure 2. The detection and recognition thresholds for the Vanishing Optotypes, and recognition values for the B/W conventional letters under different levels of blur in a) the fovea and b) 10 degrees (average across the 26 letters and both subjects). Error bars represent the SD of the 26 letter thresholds (mean of both subjects).

Figure 3. The lower parts of Figure 3a&b displays the detection and recognition values under 0D blur in the fovea for the individual a) B/W conventional letters and b) Vanishing Optotypes (average of both subjects). The upper part displays the a) recognition values for the B/W conventional letters and b) detection and recognition values for Vanishing Optotypes under +3D blur. Error bars represent the standard error of the five threshold measurements of each letter (average of both subjects). The arrows indicate thresholds for the set of 10 Sloan letters.

Figure 4. The lower part of Figure 4 displays the detection and recognition values under 0D blur in the periphery for the individual a) B/W conventional letters and b)

Vanishing Optotypes (average of both subjects). The upper part displays the a) recognition values for the B/W conventional letters and b) detection and recognition values for Vanishing Optotypes under +7D blur. Error bars represent the standard error of the five threshold measurements for each letter (average of both subjects).

Table 1. Summary of the detection and recognition thresholds and standard deviations (average of both subjects) for all tasks under 0D blur conditions.

Acuity	B/W	B/W	W/G	W/G	Hi-Pass	Hi-Pass
'logMAR'	Detect	Recog	Detect	Recog	Detect	Recog
Fovea	-0.59	-0.01	-0.47	0.01	0.11	0.14
	±0.04	±0.09	±0.05	±0.09	±0.05	±0.04
Periph	-0.08	0.63	0.02	0.73	0.56	0.85
	±0.03	±0.12	±0.06	±0.11	±0.07	±0.07

Table 1. Detection and recognition thresholds under 0D blur.

# PRECIS

High-pass (Vanishing Optotype) letters display smaller inter-letter legibility differences than conventional letters, both foveally and extrafoveally. Outside the fovea, however, they display different detection and recognition thresholds, even for high levels of defocus, indicating that extrafoveal recognition is limited by neural sampling.

Figure 1.

Z	+
Н	+
R	+
S	+
N	
B	
E	
K	

















Figure 4

