

# Sheet 8: More hyperbolic geometry

J. Evans

**The final mark out of 10 (which counts towards your grade) will be calculated as your mark on Q1 plus your mark on Q2 plus your best mark from Q3–5.** I will also award stars: silver for a total mark of 12 or more on any four questions, gold for a total mark of 15 or more on all questions.

## Question 1. (2 marks)

An ideal hyperbolic triangle is a triple of hyperbolic lines  $A, B, C$  such that each pair intersect precisely once on the boundary of hyperbolic space. For example, the semicircle  $\{z : |z| = 1, \operatorname{Im}(z) > 0\}$  and the two vertical half-lines  $\{z = -1 + ib : 0 < b \in \mathbf{R}\}$  and  $\{z = 1 + ib : 0 < b \in \mathbf{R}\}$  at  $x = -1$  and  $x = 1$  form an ideal triangle (the three “vertices” are at  $-1, 1$  and  $\infty$ ). What is the area of an ideal triangle? Why are any two ideal triangles related by an isometry of the hyperbolic plane? [Hint: Try to use 3-transitivity.]

## Question 2. (5 marks)

Working in the disc model of hyperbolic 2-space, let  $\gamma(t) = rt$  be the straight-line path starting at the origin when  $t = 0$  and finishing at radius  $r$  at time 1 and let  $\delta(t) = re^{2\pi it}$  be the circular path at radius  $r$ . A hyperbolic circle of hyperbolic radius  $R$  is defined to be the set of points a fixed hyperbolic distance  $R$  away from a fixed point.

- Find the hyperbolic length of  $\gamma$  and the hyperbolic length of  $\delta$  as functions of  $r$ .
- Deduce that a hyperbolic circle of hyperbolic radius  $R$  centred at the origin is an ordinary Euclidean circle and that the hyperbolic circumference is  $2\pi \sinh(R)$ .
- Show that the area circumscribed by a hyperbolic circle of hyperbolic radius  $R$  is  $2\pi(\cosh(R) - 1)$ .
- We have now seen that a hyperbolic circle centred at the origin looks (in the disc model) like an ordinary Euclidean circle. What if the centre is taken to be at a different point? What if we look at the circle in the upper half-plane?

**Question 3.** (3 marks)

- (a) Working in the upper half-plane model of hyperbolic space, which elements of  $PSL(2, \mathbf{R})$  send the positive imaginary half-axis to itself?
- (b) Let  $\ell$  be a straight ray in the upper half-plane starting at 0. For any  $z \in \ell$ , let  $C_z$  be the unique semicircle centred at 0 passing through  $z$ . Let  $z'$  denote the point where  $C_z$  intersects the positive imaginary axis. Prove that the hyperbolic length of the segment of  $C_z$  between  $z$  and  $z'$  depends only on the ray  $\ell$  and not on the specific choice of a point  $z \in \ell$ . [Hint: Use part (a).]

**Question 4.** (3 marks)

Let  $ABC$  be a hyperbolic triangle with edge lengths  $a, b, c$  opposite angles  $\alpha, \beta, \gamma$ . Starting from the hyperbolic cosine rule  $\cosh(a) = \cosh(b) \cosh(c) - \cos(\alpha) \sinh(b) \sinh(c)$ , prove the hyperbolic sine rule:

$$\frac{\sinh(a)}{\sin(\alpha)} = \frac{\sinh(b)}{\sin(\beta)} = \frac{\sinh(c)}{\sin(\gamma)}.$$

[Hint: You want to show that  $\sinh^2(b) \sinh^2(c) \sin^2(\alpha)$  can be written in terms of  $a, b, c$  in a completely symmetric way.]

**Question 5.** (3 marks)

Consider the semicircle  $C$  centred at  $r \in \mathbf{R}$  with radius  $r$ . What is its image under the Möbius transformation  $g(z) = -1/z$ ? What are the images under  $g$  of the points  $A = r + ri \in C$  and  $B = r(1 + e^{i\pi/4}) \in C$ ? Hence or otherwise, find the length of the segment of  $C$  connecting  $A$  to  $B$ .