

Sheet 6: More on representations

J. Evans

Question 1.

Write out the action of $X, Y \in \mathfrak{sl}(2, \mathbb{C})$ on $\text{Sym}^3(\mathbb{C}^2)$ explicitly.

Question 2.

Decompose the following representations of $\mathfrak{sl}(2, \mathbb{C})$ into irreducible summands:

- | | |
|---|---|
| (a) $\Lambda^2 \text{Sym}^3 \mathbb{C}^2,$ | (e) $\text{Sym}^2 \text{Sym}^3 \mathbb{C}^2,$ |
| (b) $\text{Sym}^2 \text{Sym}^2 \mathbb{C}^2,$ | (f) $\text{Sym}^2 \Lambda^2 \text{Sym}^3 \mathbb{C}^2,$ |
| (c) $\Lambda^3 \text{Sym}^4 \mathbb{C}^2,$ | (g) $\text{Sym}^2 \text{Sym}^4 \mathbb{C}^2,$ |
| (d) $\text{Sym}^3 \text{Sym}^2 \mathbb{C}^2,$ | (h) $\text{Sym}^3 \text{Sym}^4 \mathbb{C}^2,$ |

From your computations in (g) and (h) deduce that there are quadratic and cubic invariants $g_2(a, b, c, d, e)$ and $g_3(a, b, c, d, e)$ for binary quartic polynomials $ax^4 + bx^3y + cx^2y^2 + dxy^3 + ey^4$ under the action of $SL(2, \mathbb{C})$.

Question 3. (Clebsch-Gordan theorem)

Let V denote the standard 2-dimensional representation of $\mathfrak{sl}(2, \mathbb{C})$. Prove that the tensor product $\text{Sym}^m(V) \otimes \text{Sym}^n(V)$ decomposes into irreducible representations

$$\bigoplus_{\substack{k=|m-n| \\ k \equiv m+n \pmod{2}}}^{m+n} \text{Sym}^k(V).$$

Question 4.

Prove that if $\rho: \mathfrak{sl}(2, \mathbb{C}) \rightarrow \mathfrak{gl}(V)$ is a representation and X, Y, H denote the usual basis of $\mathfrak{sl}(2, \mathbb{C})$ satisfying the commutation relations

$$[H, X] = 2X, \quad [H, Y] = -2Y, \quad [X, Y] = H$$

then

$$C := \rho(X)\rho(Y) + \rho(Y)\rho(X) + \frac{1}{2}\rho(H)^2$$

commutes with $\rho(X)$, $\rho(Y)$ and $\rho(H)$. Deduce that if $V = \bigoplus_{\lambda} V_{\lambda}$ is the decomposition of V into eigenspaces of C then each V_{λ} is a subrepresentation. If V is irreducible with highest weight m , deduce that C is the diagonal matrix $(m + \frac{1}{2}m^2) \text{Id}$.

If $R: G \rightarrow GL(V)$ is a representation we say that:

- $M \in V$ is R -invariant if $R(g)M = M$ for all $g \in G$.
- a symmetric bilinear form $B: V \otimes V \rightarrow \mathbf{K}$ is R -invariant if

$$B(R(g)v, R(g)w) = B(v, w)$$

for all $v, w \in V$ and $g \in G$.

If $\rho: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ is a representation, we say that

- $M \in V$ is ρ -invariant if $\rho(X)M = 0$ for all $X \in \mathfrak{g}$.
- a symmetric bilinear form $B: V \times V \rightarrow \mathbf{K}$ is ρ -invariant if

$$B(\rho(X)v, w) + B(v, \rho(X)w) = 0$$

for all $v, w \in V$ and $X \in \mathfrak{g}$.

Question 5.

Let G be a connected Lie group and $R: G \rightarrow GL(V)$ be a representation. Let $\rho = R_*$ be the linearisation of R .

- Prove that $M \in V$ is G -invariant if and only if it is R_* -invariant.
- Prove that a symmetric bilinear form $B: V \times V \rightarrow \mathbf{K}$ is R -invariant if and only if it is R_* -invariant.

Hint: To show M or B is R -invariant it suffices to check it on an exponential chart because the group is connected and connected groups are generated by the image of an exponential chart.

- Let $\text{ad}: \mathfrak{g} \rightarrow \mathfrak{gl}(\mathfrak{g})$ be the adjoint representation $X \mapsto \text{ad}_X$, $\text{ad}_X Y = [X, Y]$. Define the symmetric bilinear form

$$B(X, Y) = \text{Tr}(\text{ad}_X \text{ad}_Y)$$

where Tr denotes the trace. Using some form of the Jacobi identity, prove that B is ad-invariant. This is called the *Killing form*.

Hint: The trace of a commutator of matrices vanishes.

- Let X, H, Y be the usual basis for $\mathfrak{sl}(2, \mathbf{C})$. Check that with respect to this basis

$$\text{ad}_X = \begin{pmatrix} 0 & -2 & \cdots & 0 \\ 0 & 0 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix}, \text{ad}_H = \begin{pmatrix} 2 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & -2 \end{pmatrix}, \text{ad}_Y = \begin{pmatrix} 0 & 0 & \cdots & 0 \\ -1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 2 & \cdots & 0 \end{pmatrix}$$

and hence compute the Killing form on all pairs $B(a, b)$, $a, b \in \{X, H, Y\}$.