

Sloshing, Steklov and corners I: Asymptotics of sloshing eigenvalues

Michael Levitin

Leonid Parnovski

Iosif Polterovich

David A. Sher

2 September 2017; typeset September 2, 2017

Abstract

This is the first in a series of two papers aiming to establish sharp spectral asymptotics for Steklov type problems on planar domains with corners. In the present paper we focus on the two-dimensional sloshing problem, which is a mixed Steklov-Neumann boundary value problem describing small vertical oscillations of an ideal fluid in a container or in a canal with a uniform cross-section. We prove a two-term asymptotic formula for sloshing eigenvalues. In particular, this confirms a conjecture posed by Fox and Kuttler in 1983. We also obtain similar eigenvalue asymptotics for other related mixed Steklov type problems. These results will be used in the second paper of the series, which is concerned with the Steklov spectral asymptotics on polygonal domains.

Contents

1	Introduction and main results	1
1.1	The sloshing problem	1
1.2	Asymptotics of the sloshing eigenvalues	3
1.3	Eigenvalue asymptotics for a mixed Steklov-Dirichlet problem	5
1.4	Outline of the approach	6
1.5	An application to higher order Sturm-Liouville problems	6
1.6	Towards sharp asymptotics for Steklov eigenvalues on polygons	8
1.7	Plan of the paper	9

ML: Department of Mathematics and Statistics University of Reading, Whiteknights, PO Box 220, Reading RG6 6AX, UK; M.Levitin@reading.ac.uk.

LP: Department of Mathematics, University College London, Gower Street, London WC1E 6BT, UK; leonid@math.ucl.ac.uk.

IP: Département de mathématiques et de statistique, Université de Montréal CP 6128 succ Centre-Ville, Montréal QC H3C 3J7, Canada; iossif@dms.umontreal.ca.

DAS: Department of Mathematical Sciences, DePaul University, 2320 N Kenmore Ave, Chicago, IL 60614, USA; dsher@depaul.edu.

2	Construction of quasimodes for triangular domains	10
2.1	The sloping beach problem	10
2.2	Quasimode analysis	11
3	Exponentially accurate quasimodes for angles of the form $\frac{\pi}{2q}$	15
3.1	Hanson-Lewy solutions for angles $\pi/2q$	15
3.2	Exponentially accurate quasimodes for angles of the form $\pi/2q$	19
4	Completeness of quasimodes via higher order Sturm-Liouville equations	20
4.1	An abstract linear algebra result	20
4.2	Eigenvalue asymptotics for higher order Sturm-Liouville problems	21
4.3	Connection to sloshing eigenvalues	23
4.4	Proof of Theorem 1.5.4 for triangular domains	24
4.5	Completeness of quasimodes for arbitrary triangular domains	25
4.6	Completeness of quasi-frequencies: abstract setting	26
4.7	Proof of Proposition 1.6.4	27
5	Domains with curvilinear boundary	29
5.1	Domains which are straight near the surface	29
5.2	Completeness	32
5.3	General curvilinear domains	33
A	Proof of Theorem 2.1.4	34
A.1	Robin-Neumann problem	34
A.2	Peters solution	34
A.3	Verification of Peters solution	35
A.4	Asymptotics of Peters solution as $z \rightarrow \infty$	36
	A.4.1 Error estimates for $R_\alpha(z)$	37
	A.4.2 Evaluation of $g_\alpha(-i)$	40
A.5	Asymptotics of Peters solution as $z \rightarrow 0$	42
A.6	Robin-Dirichlet problem	42
B	Proof of Proposition 4.2.4.	44
B.1	Plan of the argument	44
B.2	Proof of Lemma B.1.2	45
C	Numerical examples	47
C.1	Example illustrating Theorems 1.2.2 and 1.3.2	48
C.2	Example illustrating Remark 1.6.7	48
D	List of recurring notation	50

1 Introduction and main results

1.1 The sloshing problem

Let Ω be a simply connected bounded planar domain with Lipschitz boundary such that $\partial\Omega = S \sqcup \mathcal{W}$, where $S := (A, B)$ is a line segment. Without loss of generality we can assume that in Cartesian coordinates $A = (0, 0)$ and $B = (0, L)$, where $L > 0$ is the length of S .




Figure 1: Geometry of the sloshing problem

Consider the following mixed Steklov-Neumann boundary value problem,

$$\begin{cases} \Delta u = 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = 0 & \text{on } \mathcal{W}, \\ \frac{\partial u}{\partial n} = \lambda u & \text{on } S, \end{cases} \quad (1.1.1)$$

where $\frac{\partial}{\partial n}$ denotes the external normal derivative on $\partial\Omega$.

The eigenvalue problem (1.1.1) is called the *sloshing problem*. It describes small vertical oscillations of an ideal fluid in a container or in a canal with a uniform cross-section which has the shape of the domain Ω . The part $S = (A, B)$ of the boundary is called the *sloshing surface*; it represents the free surface of the fluid. The part \mathcal{W} is called the *walls* and corresponds to the walls and the bottom of a container or a canal. The points A and B at the interface of the sloshing surface and the walls are called the *corner points*. We also say that the walls \mathcal{W} are *straight near the corners* if there exist points $A_1, B_1 \in \mathcal{W}$ such that the line segments AA_1 and BB_1 are subsets of \mathcal{W} .

It follows from general results on the Steklov type problems that the spectrum of the sloshing problem is discrete (see [Agr06, GP17]). We denote the sloshing eigenvalues by

$$0 = \lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots \nearrow \infty,$$




Figure 2: Geometry of the sloshing problem with walls straight near the corners

where the eigenvalues are a priori counted with multiplicities. Let us note that in two dimensions all sloshing eigenvalues are conjectured to be simple, see [KKM04, GP17a]. While in full generality this conjecture remains open, in Corollary 1.2.10 we prove that it holds for all but possibly a finite number of eigenvalues.

Denote by $\mathcal{D} : L^2(S) \rightarrow L^2(S)$ the *sloshing operator*, which is essentially the Dirichlet-to-Neumann operator on S corresponding to Neumann boundary conditions on \mathcal{W} : given a function $f \in L^2(S)$, we have

$$\mathcal{D}_\Omega f = \mathcal{D}f := \left. \frac{\partial}{\partial n}(\mathcal{H}f) \right|_S, \quad (1.1.2)$$

where $\mathcal{H}f$ is the harmonic extension of f to Ω with the homogeneous Neumann conditions on \mathcal{W} . Then the sloshing eigenvalues are exactly the eigenvalues of \mathcal{D} , and the sloshing eigenfunctions are harmonic extensions of the eigenfunctions of \mathcal{D} to Ω .

The physical meaning of the sloshing eigenvalues and eigenfunctions is as follows: an eigenfunction u_k is the fluid velocity potential and $\sqrt{\lambda_k}$ is proportional to the frequency of the corresponding fluid oscillations. The research on the sloshing problem has a long history in hydrodynamics (see [Lam79, Chapter 9] and [Gre87]); we refer to [FK83] for a historical discussion.

1.2 Asymptotics of the sloshing eigenvalues

In the present paper we investigate the asymptotic distribution of the sloshing eigenvalues. It was shown in [San55] that if the boundary of Ω is C^2 -regular in a neighbourhood of the corner points A and B , then as $k \rightarrow +\infty$,

$$\lambda_k L = \pi k + o(k).$$

It follows from the results of [Agr06] on Weyl's law for mixed Steklov type problems that the C^2 assumption can be relaxed to C^1 . In 1983, Fox and Kuttler used numerical evidence to conjecture that the sloshing eigenvalues of domains having both interior angles at the corner points A and

B equal to α , satisfy the two-term asymptotics [FK83, Conjecture 3]:

$$\lambda_k L = \pi \left(k - \frac{1}{2} \right) - \frac{\pi^2}{4\alpha} + o(1), \quad (1.2.1)$$

(note that the numeration of eigenvalues in [FK83] is shifted by one compared to ours). The first main result of the present paper confirms this prediction.

Theorem 1.2.2. *Let Ω be a simply connected bounded Lipschitz planar domain with the sloshing surface $S = (A, B)$ of length L and walls \mathcal{W} which are C^1 -regular near the corner points A and B . Let α and β be the interior angles between \mathcal{W} and S at the points A and B , resp., and assume $0 < \beta \leq \alpha < \pi/2$. Then the following asymptotic expansion holds as $k \rightarrow \infty$:*

$$\lambda_k L = \pi \left(k - \frac{1}{2} \right) - \frac{\pi^2}{8} \left(\frac{1}{\alpha} + \frac{1}{\beta} \right) + r(k), \quad \text{where } r(k) = o(1). \quad (1.2.3)$$

If, moreover, the walls \mathcal{W} are straight near the corners, then

$$r(k) = O \left(k^{1 - \frac{\pi}{2\alpha}} \right). \quad (1.2.4)$$

In particular, for $\alpha = \beta$ we obtain (1.2.1) which proves the Fox-Kuttler conjecture for all angles $0 < \alpha < \pi/2$. Let us note that for $\alpha = \beta = \pi/2$ the asymptotics (1.2.1) has been earlier established in [Dav65, Ur74, Dav74]. Moreover, for $\alpha = \beta = \pi/2$ it was shown that there exist further terms in the asymptotics (1.2.1) involving the curvature of \mathcal{W} near the corner points.

Before stating our next result we require the following definition.

Definition 1.2.5. A corner point $V \in \{A, B\}$ is said to *satisfy a local John's condition* if there exist a neighbourhood \mathcal{O}_V of the point V such that the orthogonal projection of $\mathcal{W} \cap \mathcal{O}_V$ onto the x -axis is a subset of $[A, B]$.

For $\alpha = \pi/2 \geq \beta$ we obtain the following

Proposition 1.2.6. *In the notation of Theorem 1.2.2, let $\alpha = \pi/2 > \beta$, and assume that A satisfies a local John's condition. Then*

$$\lambda_k L = \pi \left(k - \frac{3}{4} \right) - \frac{\pi^2}{8\beta} + r(k), \quad r(k) = o(1). \quad (1.2.7)$$

The same result holds if $\alpha = \beta = \pi/2$, and additionally B satisfies a local John's condition.

If, moreover, the walls \mathcal{W} are straight near both corners, then

$$r(k) = O \left(k^{1 - \frac{\pi}{2\beta}} \right),$$

provided $\beta < \pi/2$ and

$$r(k) = o \left(e^{-k/C} \right),$$

if $\alpha = \beta = \pi/2$.

Proposition 1.2.6 provides a solution to [FK83, Conjecture 4] under the assumption that the corner point corresponding to the angle $\pi/2$ satisfies local John's condition.

Remark 1.2.8. In Theorem 1.2.2, and everywhere further on, C will denote various positive constants which depend only upon the domain Ω .

Remark 1.2.9. Definition 1.2.5 is a local version of John's condition which often appears in sloshing problems, see [KKM04, BKPS10].

Theorem 1.2.2 and Proposition 1.2.6 yield the following

Corollary 1.2.10. *Given a sloshing problem on a domain Ω satisfying the assumptions of Theorem 1.2.2 or Proposition 1.2.6, there exists $N > 0$, such that the eigenvalues λ_k are simple for all $k \geq N$.*

This result partially confirms the conjecture about the simplicity of sloshing eigenvalues mentioned in subsection 1.1.

1.3 Eigenvalue asymptotics for a mixed Steklov-Dirichlet problem

Boundary value problems of Steklov type with mixed boundary conditions admit several physical and probabilistic interpretations (see [BKPS10]). In particular, the sloshing problem (1.1.1) could be also used to model the stationary heat distribution in Ω such that the walls \mathcal{W} are perfectly insulated and the heat flux through S is proportional to the temperature. If instead the walls \mathcal{W} are kept under zero temperature, one obtains the following mixed Steklov-Dirichlet problem:

$$\begin{cases} \Delta u = 0 & \text{in } \Omega, \\ u = 0 & \text{on } \mathcal{W}, \\ \frac{\partial u}{\partial n} = \lambda^D u & \text{on } S, \end{cases} \quad (1.3.1)$$

Let $0 < \lambda_1^D \leq \lambda_2^D \leq \dots \nearrow \infty$ be the eigenvalues of the problem (1.3.1). Similarly to Theorem 1.2.2 we obtain

Theorem 1.3.2. *Assume that the domain Ω and its boundary $\partial\Omega = S \sqcup \mathcal{W}$ satisfy the assumptions of Theorem 1.2.2. Let α and β be the interior angles between \mathcal{W} and S at the points A and B , resp., and assume $0 < \beta \leq \alpha < \pi/2$. Then the following asymptotic expansion holds as $k \rightarrow \infty$:*

$$\lambda_k^D L = \pi \left(k - \frac{1}{2} \right) + \frac{\pi^2}{8} \left(\frac{1}{\alpha} + \frac{1}{\beta} \right) + r^D(k), \quad \text{where } r^D(k) = o(1). \quad (1.3.3)$$

If, moreover, the walls \mathcal{W} are straight near the corner points A and B , then

$$r^D(k) = O \left(k^{1-\frac{\pi}{\alpha}} \right). \quad (1.3.4)$$

We also have the following analogue of Proposition 1.2.6.

Proposition 1.3.5. *In the notation of Theorem 1.3.2, let $\alpha = \pi/2 > \beta$, and assume that A satisfies a local John's condition.. Then*

$$\lambda_k^D L = \pi \left(k - \frac{1}{4} \right) + \frac{\pi^2}{8\beta} + r(k), \quad r(k) = o(1). \quad (1.3.6)$$

The same result holds if $\alpha = \beta = \pi/2$ and additionally B satisfies a local John's condition. If, moreover, the walls W are straight near both corners, then

$$r(k) = O\left(k^{1-\frac{\pi}{\beta}}\right).$$

provided $\beta < \pi/2$, and

$$r(k) = o\left(e^{-k/C}\right)$$

for some $C > 0$ if $\alpha = \beta = \pi/2$.

This result will be used for our subsequent analysis of polygonal domains, see Subsection 1.6. The analogue of Corollary 1.2.10 clearly holds in the Steklov-Dirichlet case as well.

Remark 1.3.7. We believe that asymptotic formulae (1.2.3) and (1.3.3) in fact hold for any angles $\alpha, \beta \leq \pi$. Note also that if the walls are straight near the corners, our method yields a slightly better remainder estimate for the Steklov-Dirichlet problem compared to the Steklov-Neumann one. We refer to the proof of Theorem 2.1.4 for details.

1.4 Outline of the approach

Let us sketch the main ideas of the proof of Theorem 1.2.2; modifications needed to obtain Theorem 1.3.2 are quite minor. First, we observe that using domain monotonicity for sloshing eigenvalues (see [BKPS10]), one can deduce the general asymptotic expansion (1.2.3) from the two-term asymptotics with the remainder (1.2.4) for domains with straight walls near the corners. Assuming that the walls are straight near the corner points, we construct the *quasimodes*, i.e. the approximate eigenfunctions of the problem (1.1.1). This is done by transplanting certain model solutions of the mixed Steklov-Neumann problem in an infinite angle (cf. [DW00] where a similar idea has been implemented at a physical level of rigour). These solutions are in fact of independent interest: they were used to describe “waves on a sloping beach” (see [Han26, Lew46, Pet50]). The approximate eigenvalues given by the first two terms on the right hand side of (1.2.3) are then found from a matching condition between the two model solutions transplanted to the corners A and B , respectively. Given that the model solutions decay rapidly away from the sloshing surface, it follows that the shape of the walls away from the corners does not matter for our approximations, so in fact the domain Ω can be viewed as a triangle with angles α and β at the sloshing surface S . From the standard quasimode analysis it follows that there exist a sequence of eigenvalues of the sloshing satisfying the asymptotics (1.2.3). A major remaining challenge is to show that this sequence is *asymptotically complete* (i.e. that there are no other sloshing eigenvalues that have not been accounted for, see subsection 4.6 for a formal definition), and that the enumeration of eigenvalues given by (1.2.3) is correct.

Our quasimode construction for arbitrary angles $\alpha, \beta < \pi/2$ is based on the model solutions obtained by Peters [Pet50]. These solutions are given in terms of some complex integrals, and while their asymptotic representations allow us to construct the quasimodes, they are not accurate enough to prove completeness. However, it turns out that for angles $\alpha = \beta = \pi/2q$, $q \in \mathbb{N}$, model solutions can be written down explicitly as linear combinations of certain complex exponentials. Moreover, it turns out that for such angles the model solutions can be used to approximate the eigenfunctions of a Sturm-Liouville type problem of order $2q$ with Neumann

boundary conditions (see Theorem 1.5.4), for which the completeness follows from the general theory of linear ODEs. The enumeration of the sloshing eigenvalues in (1.2.3) may then be verified by developing the approach outlined in [Nai67, Chapter 2]. Another important property used here is a peculiar duality between the Dirichlet and Neumann eigenvalues of the Sturm-Liouville problem, see Proposition 4.2.2.

Once we have proven completeness and established the enumeration of the sloshing eigenvalues in the case $\alpha = \beta = \pi/2q$, we use once again domain monotonicity together with continuous perturbation arguments to complete the proof of Theorem 1.2.2 for arbitrary angles.

1.5 An application to higher order Sturm-Liouville problems

Let us elaborate on the link between the sloshing problem with angles $\alpha = \beta = \pi/2q$ and the higher order Sturm-Liouville type ODEs mentioned in the previous section. For a given $q \in \mathbb{N}$, consider an eigenvalue problem on an interval $(A, B) \subset \mathbb{R}$ of length L :

$$(-1)^q U^{(2q)}(x) = \Lambda^{2q} U(x), \quad (1.5.1)$$

with either Dirichlet

$$U^{(m)}(A) = U^{(m)}(B) = 0, \quad m = 0, 1, \dots, q-1, \quad (1.5.2)$$

or Neumann

$$U^{(m)}(A) = U^{(m)}(B) = 0, \quad m = q, q+1, \dots, 2q-1, \quad (1.5.3)$$

boundary conditions. For $q = 1$ we obtain the classical Sturm-Liouville equation describing vibrations of either a fixed or a free string. The case $q = 2$ yields the vibrating beam equation, also with either fixed or free ends. It follows from general elliptic theory that the spectrum of the boundary value problems (1.5.2) or (1.5.3) for the equation (1.5.1) is discrete. It is easy to check that all Dirichlet eigenvalues are positive, while the Neumann spectrum contains an eigenvalue zero of multiplicity q ; the corresponding eigenspace is generated by the functions $1, x, \dots, x^{q-1}$. Let

$$\underbrace{0 = \dots = 0}_{q \text{ times}} < (\Lambda_{q+1})^{2q} \leq (\Lambda_{q+2})^{2q} \leq \dots \nearrow \infty$$

be the spectrum of the Neumann problem (1.5.3). Then, as shown in Proposition 4.2.2, $\Lambda_k^D = \Lambda_{k+q}$, $k = 1, 2, \dots$, where $(\Lambda_k^D)^{2q}$ are the eigenvalues of the Dirichlet problem (1.5.2). We have the following result:

Theorem 1.5.4. *For any $q \in \mathbb{N}$, the following asymptotic formula holds for the eigenvalues Λ_k :*

$$\Lambda_k L = \pi \left(k - \frac{1}{2} \right) - \frac{\pi q}{2} + O(e^{-k/C}) \quad (1.5.5)$$

where $C > 0$ is some positive constant. Moreover, let λ_k , $k = 1, 2, \dots$, be the eigenvalues of a sloshing problem (1.1.1) with straight walls near the corners making equal angles $\pi/2q$ at both corner points with the sloshing surface S of length L . Then

$$\lambda_k = \Lambda_k + O(e^{-k/C}), \quad (1.5.6)$$

and the eigenfunctions u_k of the sloshing problem decrease exponentially away from the sloshing boundary S .

Spectral asymptotics of Sturm-Liouville type problems of arbitrary order for general self-adjoint boundary conditions have been studied earlier, see [Nai67, Chapter II, section 9] and references therein. However, for the special case of problem (1.5.1) with Dirichlet or Neumann boundary conditions, formula (1.5.5) gives a more precise result. First, (1.5.5) gives the asymptotics for each Λ_k , while earlier results yield only an asymptotic form of the eigenvalues without specifying the correct numbering. Second, we get an exponential error estimate, which is an improvement upon a $O(1/k)$ that was known previously. Finally, and maybe most importantly, (1.5.6) provides a physical meaning to the Sturm-Liouville problem (1.5.1) for arbitrary order $q \geq 1$. We also note that for $q = 2$, the Sturm-Liouville eigenfunctions $U_k(x)$, $k = 1, 2, \dots$, are precisely the traces of the sloshing eigenfunctions $u_k|_S$ on an isosceles right triangle $\Omega = T$ such that the sloshing surface $S = (A, B)$ is its hypotenuse. This fact was already known to H. Lamb back in the nineteenth century (see [Lam79]), and Theorem 1.5.4 extends the connection between higher order Sturm-Liouville problems and the sloshing problem to $q \geq 2$.

1.6 Towards sharp asymptotics for Steklov eigenvalues on polygons

As was discussed in [GP17, Section 3], precise asymptotics of Steklov eigenvalues on polygons and on smooth planar domains are quite different. Moreover, the powerful pseudodifferential methods used in the smooth case can not be applied for polygons, since the Dirichlet-to-Neumann operator on the boundary of a non-smooth domain is not pseudodifferential. It turns out that the methods of the present paper may be developed in order to investigate the Steklov spectral asymptotics on polygonal domains. This is the subject of the subsequent paper [LPPS17]. While establishing sharp eigenvalue asymptotics for polygons requires a lot of further work, a sharp remainder estimate in Weyl's law follows immediately from Theorems 1.2.2 and 1.3.2.


Corollary 1.6.1. *Let $N_{\mathcal{P}}(\lambda) = \#\{\lambda_k < \lambda\}$ be the counting function of Steklov eigenvalues λ_k on a convex polygon \mathcal{P} of perimeter L . Then*

$$N_{\mathcal{P}}(\lambda) = \frac{L}{\pi}\lambda + O(1). \tag{1.6.2}$$

Remark 1.6.3. The asymptotic formula (1.6.2) improves upon the previously known error bound $o(\lambda)$ (see [San55, Agr06]). Note that the $O(1)$ estimate for the error term in Weyl's law is *optimal*, since the counting function is a step-function.

Proof. Given a convex n -gon \mathcal{P} , take an arbitrary point $O \in \mathcal{P}$. It can be connected with the vertices of \mathcal{P} by n smooth curves having only point O in common in such a way that at each vertex, the angles between the sides of the polygon and the corresponding curve are smaller than $\pi/2$. This is clearly possible since all the angles of a convex polygon are less than π . Let \mathcal{L} be the union of those curves. Consider two auxiliary spectral problems: in the first one, we impose Dirichlet conditions on \mathcal{L} and keep the Steklov condition on the boundary of the polygon, and in the second one we impose Neumann conditions on \mathcal{L} and keep the Steklov condition on the boundary. Let $N_1(\lambda)$ and $N_2(\lambda)$ be, respectively, the counting functions of the first and the second problem. By Dirichlet–Neumann bracketing (which works for the sloshing problems via the variational principle in the same way it does for the Laplacian) we get

$$N_1(\lambda) \leq N_{\mathcal{P}}(\lambda) \leq N_2(\lambda), \quad \lambda > 0.$$


 Figure 3: A polygon with auxiliary curves \mathcal{L}

The spectrum of the second auxiliary problem can be represented as the union of spectra of n sloshing problems, while simultaneously the spectrum of the first auxiliary problem can be represented as the union of spectra of n corresponding Steklov-Dirichlet problems. Applying Theorems 1.2.2 and 1.3.2 to those problems, and transforming the eigenvalue asymptotics into the asymptotics of counting functions, we obtain $N_2(\lambda) - N_1(\lambda) = O(1)$, which implies (1.6.2). This completes the proof of the corollary. \square

Let us conclude this subsection by a result in the spirit of Theorems 1.2.2 and 1.3.2 that will be used in [LPPS17] in the proof of the sharp asymptotics of Steklov eigenvalues on polygons.

Proposition 1.6.4. *Let $\Omega = \triangle ABZ$ be a triangle with angles $\alpha, \beta \leq \pi/2$ at the vertices A and B , respectively. Consider a mixed Steklov-Dirichlet-Neumann problem on this triangle with the Steklov condition imposed on AB , Dirichlet condition imposed on AZ and Neumann condition imposed on BZ . Assume that the side AB has length L . Then the eigenvalues λ_k of the mixed Steklov-Dirichlet-Neumann problem on $\triangle ABZ$ satisfy the asymptotics:*

$$\lambda_k L = \pi \left(k - \frac{1}{2} \right) + \frac{\pi^2}{8} \left(\frac{1}{\alpha} - \frac{1}{\beta} \right) + O \left(k^{1 - \frac{\pi}{\max(\alpha, 2\beta)}} \right). \quad (1.6.5)$$

Remark 1.6.6. Note that the Dirichlet condition near the vertex A yields a contribution $\frac{\pi^2}{8\alpha}$ (with a positive sign) to the eigenvalue asymptotics, while the Neumann condition near B contributes $-\frac{\pi^2}{8\beta}$ (with a negative sign). This is in good agreement with the intuition provided by Theorems 1.2.2 and 1.3.2.

Remark 1.6.7. In fact, we believe that a stronger statement than the one proposed in Remark 1.3.7 is true: formula (1.6.5) holds for any mixed Steklov-Dirichlet-Neumann problem on a domain

with a curved Steklov part AB of length L , curved walls \mathcal{W} , and angles $\alpha, \beta < \pi$ at A and B , such that the Dirichlet condition is imposed near A and Neumann condition is imposed near B .

1.7 Plan of the paper

In Section 2 we use the Peters solutions of the sloping beach problem [Pet50] to construct quasimodes for the sloshing and Steklov-Dirichlet problems on triangular domains. In Section 3 we consider the case of angles of the form $\pi/2q$ for some positive integer q . In Section 4 we first prove the completeness of this system of exponentially accurate quasimodes using a connection to higher order Sturm-Liouville eigenvalue problems. In particular, we prove Theorem 1.5.4. After that, we apply domain monotonicity arguments in order to prove Theorems 1.2.2 and 1.3.2, as well as Propositions 1.2.6, 1.3.5 and 1.6.4 for triangular domains. In section 5 we extend these results to domains with curvilinear walls: first, for domains with the walls that are straight near the corners, and then to domains with general curvilinear walls. In Appendix A we prove Theorem 2.1.4, which is essentially based on the ideas of [Pet50]. In Appendix B we prove an auxiliary Proposition B.1.2 which is needed to prove Theorem 1.5.4. This section draws heavily on the results of [Nai67, Chapter 2]. Some numerical examples are presented in Appendix C. A short list of notation is in Appendix D.

Acknowledgments

The authors are grateful to Lev Buhovski for suggesting the approach used in section 5.3, to Rinat Kashaev for proposing an alternative route to prove Theorem 3.1.12, as well as to Alexandre Girouard and Yakar Kannai for numerous useful discussions on this project. The research of L.P. was supported by by EPSRC grant EP/J016829/1. The research of I.P. was supported by NSERC, FRQNT, Canada Research Chairs program, as well as the Weston Visiting Professorship program at the Weizmann Institute of Science, where part of this work has been accomplished. The research of D.S. was supported in part by NSF EMSW21-RTG 1045119 and in part by a Faculty Summer Research Grant from DePaul University.

2 Construction of quasimodes for triangular domains

2.1 The sloping beach problem

Let (x, y) be Cartesian coordinates in \mathbb{R}^2 , let $z = x + iy \in \mathbb{C}$, and let (ρ, θ) denote the polar coordinates $z = \rho e^{i\theta}$. Let \mathfrak{S}_α be the planar sector $-\alpha \leq \theta \leq 0$. Consider the following mixed Robin-Neumann problem

$$\begin{cases} \Delta\phi = 0 & \text{in } \mathfrak{S}_\alpha, \\ \frac{\partial\phi}{\partial y} = \phi & \text{on } \mathfrak{S}_\alpha \cap \{\theta = 0\}, \\ \frac{\partial\phi}{\partial n} = 0 & \text{on } \mathfrak{S}_\alpha \cap \{\theta = -\alpha\}, \end{cases} \quad (2.1.1)$$

and the mixed Robin-Dirichlet problem

$$\begin{cases} \Delta\phi = 0 & \text{in } \mathfrak{S}_\alpha, \\ \frac{\partial\phi}{\partial y} = \phi & \text{on } \mathfrak{S}_\alpha \cap \{\theta = 0\}, \\ \phi = 0 & \text{on } \mathfrak{S}_\alpha \cap \{\theta = -\alpha\}. \end{cases} \quad (2.1.2)$$

Note that there is no spectral parameter in these problems (hence the boundary conditions are called Robin rather than Steklov). Our aim is to exhibit solutions of (2.1.1) and (2.1.2) decaying away from the horizontal part of the boundary with the property that $\phi(x, 0) \rightarrow \cos(x - \chi)$ for some fixed χ as $x \rightarrow \infty$.

This problem is known as the *sloping beach problem*, and has a long and storied history in hydromatics, see [Lew46] references therein. It turns out that the form of the solutions depends in a delicate way on the *arithmetic* properties of the angle α ; we will discuss this in more detail later on. Let

$$\mu_\alpha = \frac{\pi}{2\alpha}, \quad \chi_{\alpha,N} = \frac{\pi}{4}(1 - \mu_\alpha), \quad \chi_{\alpha,D} = \frac{\pi}{4}(1 + \mu_\alpha). \quad (2.1.3)$$

The following key result was essentially established by Peters [Pet50]:

Theorem 2.1.4. *For any $0 < \alpha < \pi/2$, there exist solutions $\phi_{\alpha,N}(x, y)$ and $\phi_{\alpha,D}(x, y)$ of (2.1.1) and (2.1.2), respectively, and a constant $C > 0$ such that:*

- $|\phi_{\alpha,N}(x, y)|$ and $|\phi_{\alpha,D}(x, y)|$ are bounded on the closed sector $\overline{\mathfrak{S}_\alpha}$;

-

$$\phi_{\alpha,N}(x, y) = e^y \cos(x - \chi_{\alpha,N}) + R_{\alpha,N}(x, y), \quad (2.1.5)$$

where

$$|R_{\alpha,N}(x, y)| \leq C\rho^{-\mu}, \quad |\nabla_{(x,y)} R_{\alpha,N}(x, y)| \leq C\rho^{-\mu-1};$$

-

$$\phi_{\alpha,D}(x, y) = e^y \cos(x - \chi_{\alpha,D}) + R_{\alpha,D}(x, y), \quad (2.1.6)$$

where

$$|R_{\alpha,D}(x, y)| \leq C\rho^{-2\mu}, \quad |\nabla_{(x,y)} R_{\alpha,D}(x, y)| \leq C\rho^{-2\mu-1}.$$

For $\alpha = \pi/2$ we have

$$\phi_{\frac{\pi}{2},N} = e^y \cos x, \quad \phi_{\frac{\pi}{2},D} = e^y \sin x. \quad (2.1.7)$$

Remark 2.1.8. Note that $R_{\alpha,N}(x, y)$ and $R_{\alpha,D}(x, y)$ are harmonic. Note also that our further analysis (specifically, the asymptotic completeness of quasimodes) will imply that the solutions $\phi_{\alpha,N}(x, y)$ and $\phi_{\alpha,D}(x, y)$ are unique in the sense that we cannot replace the particular constants $\chi_{\alpha,N}$ and $\chi_{\alpha,D}$ from (2.1.3) by any other value.

The construction of the solutions for both the Robin-Neumann problem and the Robin-Dirichlet problem is due to Peters [Pet50]. The approach is based on complex analysis, specifically the Wiener-Hopf method. The solution is written down explicitly as a complex integral, which allows us to analyse the asymptotics. We have reproduced this construction, taking special care with the remainder estimates that were not worked out in [Pet50]. The proof of Theorem 2.1.4 is quite technical and is deferred until Appendix A.

2.2 Quasimode analysis

In what follows, we focus on the proof of Theorem 1.2.2; minor modifications required for Theorem 1.3.2 will be discussed later. As such we suppress all N subscripts.

As was indicated in subsection 1.7, we split the proof of Theorem 1.2.2 into several steps. Our first objective is to prove the asymptotic expansion (1.2.3) with the remainder estimate (1.2.4) for triangular domains.

Convention: *From now on and until subsection 4.5 we assume that Ω is a triangle $T = \triangle ABZ$.*

The key starting idea is to glue together two scaled Peters solutions $\pm\phi_\alpha(\sigma x, \sigma y)$, one at each corner A and B , to construct quasimodes. These Peters solutions must match, meaning that the phases of the trigonometric functions in the asymptotics of both solutions must agree. Denoting those phases by $\chi_\alpha = \chi_{\alpha,N}$ and $\chi_\beta = \chi_{\beta,N}$, see (2.1.3), we require

$$\cos(\sigma x - \chi_\alpha) = \pm \cos((L - x)\sigma - \chi_\beta).$$

Solving this equation immediately gives $\sigma = \sigma_k$ for some integer k , where

$$\sigma_k L = \pi \left(k - \frac{1}{2} \right) - \frac{\pi^2}{8} \left(\frac{1}{\alpha} + \frac{1}{\beta} \right). \quad (2.2.1)$$

This could be viewed as the *quantization condition* resulting in the asymptotics (1.2.3).

Let us now make this precise. Let the plane wave $p_\sigma(x, y) = e^{\sigma y} \cos(\sigma x - \chi_\alpha)$, where σ is determined by the quantization condition (2.2.1). Letting $z = (x, y)$, we set

$$R_{A,\sigma}(z) := \phi_\alpha(\sigma z) - p_\sigma(z);$$

that is, $R_{A,\sigma}(z)$ is the difference between the scaled Peters solution in the sector of angle α with the vertex at A , and the scaled trigonometric function. Similarly, let

$$R_{B,\sigma}(z) := \phi_\beta(\sigma(L - \bar{z})) - p_\sigma(L - \bar{z})$$

be the difference between the scaled Peters solution in the sector of angle β with the vertex at B and a scaled trigonometric function. Note that $R_{A,\sigma}(z) = R_\alpha(z\sigma)$ and similarly for $R_{B,\sigma}(z)$. By the respective remainder estimates in Theorem 2.1.4,

$$\begin{aligned} |R_{A,\sigma}(z)| &\leq C\sigma^{-\mu_\alpha} |z - A|^{-\mu_\alpha}; & |R_{B,\sigma}(z)| &\leq C\sigma^{-\mu_\beta} |L - \bar{z} - B|^{-\mu_\beta}; \\ |\nabla R_{A,\sigma}(z)| &\leq C\sigma^{-\mu_\alpha} |z - A|^{-\mu_\alpha-1}; & |\nabla R_{B,\sigma}(z)| &\leq C\sigma^{-\mu_\beta} |L - \bar{z} - B|^{-\mu_\beta-1}. \end{aligned}$$

For $\sigma = \sigma_k$ satisfying the quantization condition (2.2.1), let us define a function v'_σ (note that v'_σ is not a derivative of v_σ but a new function, and we will follow this convention in the sequel):

$$v'_\sigma(z) := p_\sigma(z) + R_{A,\sigma}(z) + R_{B,\sigma}(z) \quad (2.2.2)$$

(indeed, this definition is meaningful only when σ satisfies (2.2.1)). This is our first attempt at a quasimode. The problem with this function is that, while it is harmonic, it does not satisfy the Neumann condition on \mathcal{W} . However, the error is small and we can correct it.

Indeed, in a neighbourhood of A , the function $p_\sigma + R_{A,\sigma}$ is the Peters solution and hence satisfies the Neumann condition on AZ , and $\nabla R_{B,\sigma}$ is of order $O(\sigma^{-\mu_\beta})$, so the normal derivative

of v'_σ is $O(\sigma^{-\mu_\beta})$. A similar analysis holds in a neighbourhood of B . Away from both A and B , all three terms have gradients of magnitude $O(\sigma^{-\mu_\alpha})$ (as $\alpha \geq \beta$). Therefore

$$\left| \frac{\partial v'_\sigma}{\partial n} \right| \leq C\sigma^{-\mu_\alpha} \quad \text{on } \mathcal{W}. \quad (2.2.3)$$

In order to correct this ‘‘Neumann defect’’, consider a function η_σ defined as a solution of the following system:

$$\begin{cases} \Delta \eta_\sigma = 0 & \text{in } \Omega; \\ \frac{\partial}{\partial n} \eta_\sigma = \frac{\partial}{\partial n} v'_\sigma & \text{on } \mathcal{W}; \\ \frac{\partial}{\partial n} \eta_\sigma = -\kappa_\sigma \psi & \text{on } S, \end{cases} \quad (2.2.4)$$

where $\psi \in C^\infty(S)$ is a fixed function, supported away from the vertices, with $\int_S \psi = 1$, and where

$$\kappa_\sigma = \int_{\mathcal{W}} \frac{\partial v'_\sigma}{\partial n}.$$

Note that the integral of the Neumann data over $\partial\Omega$ in (2.2.4) is zero; thus a solution η_σ to (2.2.4) exists and is uniquely defined up to an additive constant. It is well-known (see, for instance, [Ron15]) that the Neumann-to-Dirichlet map is a bounded operator on $L^2_*(\partial\Omega)$, the space of mean-zero L^2 functions on the boundary. Therefore, $\|\eta_\sigma\|_{L^2(\partial\Omega)} \leq C\sigma^{-\mu_\alpha}$, and hence

$$\|\eta_\sigma\|_{L^2(S)} \leq C\sigma^{-\mu_\alpha}. \quad (2.2.5)$$

With this auxiliary function, we define a corrected quasimode:

$$v_\sigma(z) := v'_\sigma(z) - \eta_\sigma(z) = p_\sigma(z) + R_{A,\sigma}(z) + R_{B,\sigma}(z) - \eta_\sigma(z).$$

Observe that v_σ is harmonic and satisfies the homogeneous Neumann boundary condition on \mathcal{W} .

The key property of our new quasimodes is the following

Lemma 2.2.6. *With the notation as above, there exists a constant C such that*

$$\|\mathcal{D}v_\sigma - \sigma v_\sigma\|_{L^2(S)} \leq C\sigma^{1-\mu_\alpha}, \quad (2.2.7)$$

where \mathcal{D} is the sloshing operator (1.1.2).

Remark 2.2.8. Note that $\mu_\alpha = \frac{\pi}{2\alpha} > 1$ for $\alpha < \pi/2$, and therefore we get quasimodes of order $O(\sigma^{-\delta})$ for $\delta := 1 - \mu_\alpha > 0$. For $\alpha \geq \pi/2$ one would need to modify our approach.

Proof of Lemma 2.2.6. Since v_σ is harmonic and satisfies the Neumann condition on \mathcal{W} , we have

$$\mathcal{D}(v_\sigma|_S) = \frac{\partial v_\sigma}{\partial n} \Big|_S.$$

Consider first a region away from the vertex A . In this region,

$$\frac{\partial(p_\sigma + R_{B,\sigma})}{\partial n} \Big|_S - \sigma(p_\sigma + R_{B,\sigma})|_S = 0$$

by Theorem 2.1.4. Moreover, due to the bounds on $R_{A,\sigma}$ in Theorem 2.1.4,

$$\left| \frac{\partial R_{A,\sigma}}{\partial n} - \sigma R_{A,\sigma} \right| \leq C\sigma^{1-\mu_\alpha} \quad \text{on } S$$

pointwise, and hence the same estimate holds in $L^2(S)$. Finally, by construction of η_σ , we know

$$\frac{\partial \eta_\sigma}{\partial n} \Big|_S = -c_\sigma \psi, \quad |c_\sigma| \leq C\sigma^{-\mu_\alpha};$$

combining this with (2.2.5) yields

$$\left\| \frac{\partial \eta_\sigma}{\partial n} - \sigma \eta_\sigma \right\|_{L^2(S)} \leq C\sigma^{1-\mu_\alpha}.$$

Putting everything together using the definition of v_σ gives us the required estimate away from A . A similar analysis shows that the same estimate holds away from B , completing the proof. \square

Remark 2.2.9. For the Dirichlet or mixed problems on triangles, it is also possible to construct $\eta_\sigma(z)$ harmonic and satisfying (2.2.5) such that $v_\sigma(z)$ satisfies the appropriate homogeneous boundary conditions on $\mathcal{W} := \mathcal{W}_1 \cup \mathcal{W}_2$. In this case, η_σ is a solution of the following problem:

$$\begin{cases} \Delta \eta_\sigma = 0 & \text{in } \Omega; \\ \frac{\partial}{\partial n} \eta_\sigma = \frac{\partial}{\partial n} v'_\sigma & \text{on any Neumann side } \mathcal{W}_1; \\ \eta_\sigma = v'_\sigma & \text{on any Dirichlet side } \mathcal{W}_2; \\ \frac{\partial}{\partial n} \eta_\sigma = 0 & \text{on } S. \end{cases} \quad (2.2.10)$$

Indeed, solutions to such mixed problems are known to exist even on a larger class of Lipschitz domains [Bro94], which includes our setting (see the discussion in [Bro94, Introduction]). By [Bro94, Theorem 1.5], the solution η_σ exists, is unique, and

$$\|\nabla \eta_\sigma\|_{L^2(\partial\Omega)}^2 + \|\eta_\sigma\|_{L^2(\partial\Omega)}^2 \leq C \left(\left\| \frac{\partial}{\partial n} v'_\sigma \right\|_{L^2(\mathcal{W}_1)}^2 + \|\nabla_{\tan} v'_\sigma\|_{L^2(\mathcal{W}_2)}^2 + \|v'_\sigma\|_{L^2(\mathcal{W}_2)}^2 \right),$$

where ∇_{\tan} denotes a tangential derivative along \mathcal{W} . By Theorem (2.1.4), the first two terms on the right hand side are actually bounded by $(C\sigma^{-\mu_\alpha-1})^2$, and the third term is bounded by $(C\sigma^{-\mu_\alpha})^2$. Overall, we obtain

$$\|\nabla \eta_\sigma\|_{L^2(\partial\Omega)}^2 + \|\eta_\sigma\|_{L^2(\partial\Omega)}^2 \leq (C\sigma^{-\mu_\alpha})^2,$$

so in particular

$$\|\eta_\sigma\|_{L^2(\partial\Omega)} \leq C\sigma^{-\mu_\alpha}. \quad (2.2.11)$$

The analysis then proceeds as above, and in particular the analogue of Lemma 2.2.6 holds by an identical proof.

Going back to the Neumann setting again, we let

$$\sigma_j = \frac{1}{L} \left(\pi \left(j - \frac{1}{2} \right) - \frac{\pi^2}{8} \left(\frac{1}{\alpha} + \frac{1}{\beta} \right) \right), \quad j = 1, 2, \dots, \quad (2.2.12)$$

as in (2.2.1). Abusing notation slightly, let $v_j := v_{\sigma_j}|_S$ be the traces of the corresponding quasimodes. Assume further that $\|v_j\|_{L^2(S)} = 1$. By (2.2.12), $\sigma_j \geq cj$ for some $c > 0$, and thus we have

$$\|\mathcal{D}v_j - \sigma_j v_j\|_{L^2(S)} \leq Cj^\delta, \quad \delta = 1 - \mu_\alpha > 0. \quad (2.2.13)$$

Now let $\{\varphi_k\}_{k=1}^\infty$ be an orthonormal basis of the eigenfunctions of the sloshing operator \mathcal{D} , with eigenvalues λ_j . By completeness and orthonormality of the $\{\varphi_k\}$, we have, for each j ,

$$v_j = \sum_{k=1}^{\infty} a_{jk} \varphi_k, \quad a_{jk} = (v_j, \varphi_k), \quad \sum_{k=1}^{\infty} a_{jk}^2 = 1. \quad (2.2.14)$$

Plugging in (2.2.13), we get

$$\left\| \sum_{k=1}^{\infty} a_{jk} (\lambda_k - \sigma_j) \varphi_k \right\|_{L^2(S)} \leq Cj^{-\delta}$$

and hence

$$\sum_{k=1}^{\infty} a_{jk}^2 (\lambda_k - \sigma_j)^2 \leq Cj^{-2\delta}.$$

Since $\sum_{k=1}^{\infty} a_{jk}^2 = 1$, it cannot be true that $|\lambda_k - \sigma_j| > Cj^{-\delta}$ for all k . Therefore the following Lemma holds.

Lemma 2.2.15. *For each $j \in \mathbb{N}_0$, there exists $k \in \mathbb{N}_0$ such that*

$$|\sigma_j - \lambda_k| \leq Cj^{-\delta}.$$

In other words, there exists a subsequence of the spectrum that behaves asymptotically as σ_j , up to an error which is $O(j^{-\delta})$. The key question that we now face is how to prove that $j = k$, i.e., that the sequence gives the *full* spectrum. In order to achieve this, we have to deal with several issues. First, we do not know whether the quasimodes $\{v_j\}$ form a basis for $L^2(S)$. Second, we do not have good control of the errors $R_{A,\sigma}$ and $R_{B,\sigma}$ near their respective corners. Therefore, some new ideas are needed; in particular, we need more accurate quasimodes. Luckily for us, as will be shown later, we can get away with constructing such quasimodes for some angles only.

3 Exponentially accurate quasimodes for angles of the form

$$\frac{\pi}{2q}$$

3.1 Hanson-Lewy solutions for angles $\pi/2q$

We will now construct quasimodes in the special case where both angles α and β are equal to $\pi/2q$ for some integer $q \geq 2$.

To do this, let us first go back to the study of waves on sloping beaches. Recall that \mathfrak{S}_α is the planar sector with $-\alpha \leq \theta \leq 0$, and let I_1 be the half-line $\theta = 0$ with I_2 the half-line $\theta = -\alpha$. Consider the complex variable z . Let q be a positive integer and let $\alpha = \pi/2q$,

$$\xi = \xi_q := e^{-i\pi/q}. \quad (3.1.1)$$

Proposition 3.1.2. *Suppose $g(z)$ is an arbitrary function. Set $(\mathcal{A}g)(z) := g(\xi\bar{z})$. Then*

$$(g - \mathcal{A}g)|_{I_2} = 0, \quad \frac{\partial}{\partial n}(g + \mathcal{A}g)|_{I_2} = 0.$$

Proof. This can be checked by direct computation. □

The following definition is useful throughout.

Definition 3.1.3. The *Steklov defect* of an arbitrary function $g(z)$ on \mathfrak{S}_α is

$$\text{SD}(g) := \left(\frac{\partial g}{\partial y} - g \right) \Big|_{I_1}.$$

Note that $g(z)$ satisfies the Steklov condition with parameter 1 if and only if $\text{SD}(g)$ is identically zero.

For setup purposes, consider functions of the form

$$g(z) = e^{p\bar{z}}, \quad h(z) = e^{pz}.$$

Differentiation immediately tells us that

$$\text{SD}(g) = (-ip - 1)g|_{I_1}; \quad \text{SD}(h) = (ip - 1)h|_{I_1}.$$

Given $g = e^{p\bar{z}}$, set

$$\mathcal{B}g(z) = \frac{ip + 1}{ip - 1} e^{pz}.$$

It is immediate that

Proposition 3.1.4. *For any $g = e^{p\bar{z}}$, with $p \in \mathbb{C}$,*

$$\text{SD}(g + \mathcal{B}g) = 0.$$

Now let $f(z) = e^{-iz}$, and consider

$$f_0(z) = f(z); \quad f_1(z) = \mathcal{A}f_0(z), \quad f_2 = \mathcal{B}f_1(z), \quad f_3 = \mathcal{A}f_2(z), \dots$$

We then have

$$f_1(z) = f(\xi\bar{z}), \quad f_2(z) = \eta(\xi)f(\xi z),$$

where

$$\eta(\xi) = \frac{i(-i\xi) + 1}{i(-i\xi) - 1} = \frac{\xi + 1}{\xi - 1}.$$

Further on,

$$f_3(z) = \eta(\xi)f(\xi^2\bar{z}), \quad f_4(z) = \eta(\xi)\eta(\xi^2)f(\xi^2z), \dots,$$

$$f_{2q-1}(z) = \eta(\xi)\eta(\xi^2) \dots \eta(\xi^{q-1})f(\xi^q\bar{z}) = \prod_{j=1}^{q-1} \eta(\xi^j)f(-\bar{z}).$$

Finally, set

$$v_\alpha(z) = \sum_{m=0}^{2q-1} f_m(z). \quad (3.1.5)$$

Theorem 3.1.6. *The function $v_\alpha(z)$ defined by (3.1.5) is harmonic, satisfies Neumann boundary conditions on I_2 , and $\text{SD}(v_\alpha) = 0$.*

Proof. Since $v_\alpha(z)$ is a sum of rotated plane waves, it is harmonic. It satisfies Neumann boundary conditions because we can write

$$\begin{aligned} v_\alpha &= (f_0 + f_1) + (f_2 + f_3) + \dots + (f_{2q-2} + f_{2q-1}) \\ &= (f_0 + \mathcal{A}f_0) + (f_2 + \mathcal{A}f_2) + \dots + (f_{2q-2} + \mathcal{A}f_{2q-2}) \end{aligned}$$

and use Proposition 3.1.2 on each term. To see that it satisfies $\text{SD}(v_\alpha) = 0$, write

$$\begin{aligned} v_\alpha &= f_0 + (f_1 + f_2) + \dots + (f_{2q-3} + f_{2q-2}) + f_{2q-1} \\ &= f_0 + (f_1 + \mathcal{B}f_1) + \dots + (f_{2q-3} + \mathcal{B}f_{2q-3}) + f_{2q-1}. \end{aligned}$$

Using Proposition 3.1.4 and the linearity of SD, we have

$$\text{SD}(v_\alpha) = \text{SD}(f_0) + \text{SD}(f_{2q-1}),$$

and all that remains is to show these two terms sum to zero. In fact both are separately zero, because $f_0(z) = e^{-iz} = e^{-ix}e^y$, and $\text{SD}(f_0)$ is zero by direct computation. Moreover,

$$f_{2q-1}(z) = \prod_{j=1}^{q-1} \eta(\xi^j)e^{i\bar{z}} = \prod_{j=1}^{q-1} \eta(\xi^j)e^{ix}e^y,$$

and the same direct computation shows $\text{SD}(f_{2q-1}) = 0$. This completes the proof. \square

We call $v_\alpha(z)$ the *Hanson-Lewy solution* for the sloping beach problem with angle $\alpha = \pi/2q$. It is helpful to introduce the notation

$$\gamma(\xi) := \prod_{j=1}^{q-1} \eta(\xi^j). \quad (3.1.7)$$

Lemma 3.1.8. *We have $\gamma(\xi) = e^{i\pi(q-1)/2}$. If q is even, $\gamma(\xi) = \pm i$, and if q is odd, $\gamma(\xi) = \pm 1$.*

Proof. We have

$$\gamma(\xi) = \prod_{j=1}^{q-1} \frac{e^{-i\pi j/q} + 1}{e^{-i\pi j/q} - 1} = \prod_{j=1}^{q-1} \frac{1 + e^{i\pi j/q}}{1 - e^{i\pi j/q}},$$

where we have multiplied numerator and denominator by $e^{i\pi j/q}$. Relabeling terms in the numerator by switching j for $q - j$ gives

$$\gamma(\xi) = \prod_{j=1}^{q-1} \frac{1 + e^{i\pi(q-j)/q}}{1 - e^{i\pi j/q}} = \prod_{j=1}^{q-1} \frac{1 - e^{-i\pi j/q}}{1 - e^{i\pi j/q}} = \prod_{j=1}^{q-1} e^{-i\pi j/q} \frac{e^{i\pi j/q} - 1}{1 - e^{i\pi j/q}} = \prod_{j=1}^{q-1} (-e^{-i\pi j/q}).$$

This may be rewritten as

$$\gamma(\xi) = \prod_{j=1}^{q-1} (e^{i\pi - i\pi j/q}) = \exp \left(i\pi \left(q - 1 - \sum_{j=1}^{q-1} \frac{j}{q} \right) \right) = \exp \left(i\pi \left(q - 1 - \frac{q-1}{2} \right) \right),$$

which yields the result. \square

Lemma 3.1.9. *On I_1 , the Hanson-Lewy solutions $v_\alpha(z)$ are of the form*

$$v_\alpha(x) = e^{-ix} + \gamma(\xi)e^{ix} + \text{decaying exponentials.}$$

Proof. Indeed, the first and last terms are $e^{-iz} + \gamma(\xi)e^{i\bar{z}}$, and all other exponentials in the sum defining v_α are of the form $e^{-i\xi^j z}$ or $e^{-i\xi^j \bar{z}}$ for $j = 1, \dots, q-1$. On I_1 , $z = \bar{z} = x$, so we have $e^{-i\operatorname{Re}(\xi^j)x} e^{\operatorname{Im}(\xi^j)x}$. But $\operatorname{Im}(\xi^j) = \sin(-\pi j/q) < 0$ for $j = 1, \dots, q-1$, and the exponential is thus decaying. \square

This may be strengthened:

Lemma 3.1.10. *The rescaled solution $v_\alpha(\sigma z)$ is $e^{i\sigma z} + \gamma(\xi)e^{i\sigma \bar{z}}$ plus a remainder which is exponentially decaying in σ as $\sigma \rightarrow \infty$ for each $z \in \mathfrak{S}_\alpha$. The exponential decay constant is uniform over all $z \in \mathfrak{S}_\alpha$ with $|z| = 1$.*

Proof. Each term in the sum defining $v_\alpha(z)$ other than the first and last terms is $e^{-i\xi^j z}$ or $e^{-i\xi^j \bar{z}}$ for $j = 1, \dots, q-1$. Observe that $|e^{-i\xi^j \sigma z}|$ decays exponentially in σ when $\xi^j z$ is in the negative half-plane, uniformly for z away from the real axis. Since $\arg \xi^j$ is $-\pi j/q$ for $1 \leq j \leq q-1$ and $\arg z$ is between 0 and $-\pi/2q$, we see that for each $z \in \mathfrak{S}_\alpha$,

$$-\frac{\pi}{q} \leq \arg(\xi^j z) \leq -\pi + \frac{\pi}{2q}.$$

Thus we have the required decay and it is uniform over the set of $z \in \mathfrak{S}_\alpha$ with $|z| = 1$. A similar calculation shows that

$$-\frac{\pi}{2q} \leq \arg(\xi^j \bar{z}) \leq -\pi + \frac{\pi}{q},$$

and the same argument applies, completing the proof. \square

Remark 3.1.11. A similar construction holds for the solutions of the mixed Steklov-Dirichlet problem. Set

$$\tilde{f}_0 = f; \tilde{f}_1 = -\mathcal{A}\tilde{f}_0; \tilde{f}_2 = \mathcal{B}\tilde{f}_1; \tilde{f}_3 = -\mathcal{A}\tilde{f}_2, \dots$$

Then set

$$\tilde{v}_\alpha(z) = \sum_{e=0}^{2q-1} \tilde{f}_e(z).$$

By Propositions 3.1.2 and 3.1.4, these satisfy $\tilde{v}_\alpha|_{L^2} = 0$ and $\text{SD}(u) = 0$. They also have the same exponential decay properties.

Here is a key, novel, observation about the Hanson-Lewy solutions:

Theorem 3.1.12. *For each $\alpha = \frac{\pi}{2q}$, the solutions $v_\alpha(z)$ and $\tilde{v}_\alpha(z)$ satisfy the following properties when restricted to I_1 :*

$$v_\alpha^{(m)}(0) = 0, \quad m = q, \dots, 2q-1; \quad \tilde{v}_\alpha^{(m)}(0) = 0, \quad m = 0, \dots, q-1.$$

Proof. Consider first the Neumann case. Set $y = 0$. We have

$$v_\alpha(x) = e^{-ix} + e^{-i\xi x} + \eta(\xi)e^{-i\xi x} + \eta(\xi)e^{-i\xi^2 x} + \dots$$

Therefore

$$v_\alpha^{(m)}(x) = (-i)^m (1 + \xi^m) \left(1 + \sum_{k=1}^{q-1} \xi^{km} \prod_{j=1}^k \eta(\xi^j) \right). \quad (3.1.13)$$

We now apply the Lemma in [Lew46, p.745]. In the notation of Lewy, $\epsilon = \xi = e^{-i\pi/q}$, and the product in the definition of $f(\xi)$ in [Lew46] is precisely the one appearing in (3.1.13) times $(-1)^k$. Hence, the right hand side of (3.1.13) vanishes precisely for $m = q, \dots, 2q-1$. Indeed, for $m = q$ this is obvious since $1 + \xi^q = 0$; for $m = q + r$, $r = 1, \dots, q-1$, we have $\xi^{k(q+r)} = \xi^r (\xi^q)^k = (-1)^k \xi^{kr}$, and the result follows from Lewy's lemma.

In the Dirichlet case we have

$$\tilde{v}_\alpha(x) = e^{-ix} - e^{-i\xi x} - \eta(\xi)e^{-i\xi x} + \eta(\xi)e^{-i\xi^2 x} + \dots$$

Therefore

$$\tilde{v}_\alpha^{(m)}(0) = (-i)^m (1 - \xi^m) \left(1 + \sum_{k=1}^{q-1} (-1)^k \xi^{km} \prod_{j=1}^k \eta(\xi^j) \right).$$

A similar use of Lewy's lemma shows this is zero for $m = 0, 1, \dots, q-1$, and the theorem is proved. \square

Remark 3.1.14. An alternative proof of this Theorem was provided to us by Rinat Kashaev [Kas16], based on combinatorial techniques used in [KMS93].

Remark 3.1.15. The Hanson-Lewy solutions are closely connected to a higher order Sturm-Liouville type ODE on the real line, specifically

$$(-1)^q U^{(2q)} = \Lambda^{2q} U. \quad (3.1.16)$$

It is immediate that the Hanson-Lewy solutions v_α and \tilde{v}_α satisfy this equation with $\Lambda = 1$. This observation together with Theorem 3.1.12 will be key for the next step of the argument.

3.2 Exponentially accurate quasimodes for angles of the form $\pi/2q$

Consider the sloshing problem on a triangle with top side of length $L = 1$ (for simplicity) and with angles $\alpha = \beta = \pi/2q$, $q \in \mathbb{N}$, q even. Suppose that σ satisfies the quantization condition (2.2.1). For such σ , let

$$g_\sigma(z) = v_\alpha(\sigma z) + v_\beta^d(\sigma(1 - \bar{z})),$$

where v_β^d is the sum of just the *decaying* exponents in v_β (see Lemma 3.1.9). Note that the choice of σ yields that the oscillating exponents in $(v_\alpha|_{I_1})(\sigma x)$ and $(v_\beta|_{I_1})(\sigma(1 - x))$ coincide, so there is a similar decomposition near B .

First we view the functions g_σ as quasimodes for the sloshing operator. We are in the same situation we were with the Peters solutions, but now the errors decay exponentially rather than polynomially. By an identical argument to that in subsection 2.2, there exist functions $\eta_\sigma(z)$ satisfying (2.2.11) such that if we set $v_\sigma(z) = g_\sigma(z) - \eta_\sigma(z)$, then we have the key quasimode estimate

$$\|\mathcal{D}(v_\sigma|_S) - \sigma v_\sigma\|_{L^2(S)} = O(e^{-\sigma/C}). \quad (3.2.1)$$

Now we view g_σ as quasimodes for an ODE problem. By Theorem 3.1.12, we see that $g_\sigma(x)$ is a solution of the ODE (3.1.16), and moreover one which satisfies up to an error $O(e^{-\sigma/C})$, for some $C > 0$, the self-adjoint boundary conditions

$$g_\sigma^{(m)}(0) = g_\sigma^{(m)}(1) = 0, \quad m = q, \dots, 2q - 1.$$

at the ends of the interval. Specifically, we claim that these functions $g_\sigma(x)$ are exponentially accurate quasimodes on $[0, 1]$ for the elliptic self-adjoint ODE eigenvalue problem

$$\begin{cases} (-1)^q U^{(2q)} = \Lambda^{2q} U \\ U^{(m)}(0) = U^{(m)}(1) = 0 \text{ for } m = q, \dots, 2q - 1. \end{cases} \quad (3.2.2)$$

In order to show this, we need to run a similar argument to the one we have run in subsection 2.2 to correct our quasimodes. First, we correct the boundary values of g_σ in order for the quasimode to lie in the domain of the operator. We do it as follows.

Let $\Phi_m(x)$ be a function equal to x^m near $x = 0$ and smoothly decaying so that it is identically zero whenever $x > 1/2$. Then for each $\sigma = \sigma_k$, there exists a function

$$\bar{\eta}_\sigma(x) = \sum_{m=q}^{2q-1} (a_m \Phi_m(x) + b_m \Phi_m(1 - x)),$$

where a_m, b_m are both $O(e^{-\sigma/C})$, and where $g_\sigma(x) + \bar{\eta}_\sigma(x)$ satisfy the boundary conditions in (3.2.2). In other words,

$$\bar{v}_\sigma(x) = g_\sigma(x) + \bar{\eta}_\sigma(x)$$

could be used as a quasimode for the eigenvalue problem (3.2.2). Indeed, it is easy to see that

$$\|(-1)^q \bar{v}_\sigma^{(2q)}(x) - \sigma^{2q} \bar{v}_\sigma(x)\|_{L^2([0,1])} = O(e^{-\sigma/C}). \quad (3.2.3)$$

The consequences of these quasimode estimates will be discussed in the next section. Note that by our estimates on η , the functions $\bar{\eta}$, v_σ , g_σ , and \bar{v}_σ are all within $O(e^{-\sigma/C})$ in $L^2(S)$ norm, and in particular

$$\|\bar{v}_\sigma - v_\sigma\|_{L^2(S)} \leq C e^{-\sigma/C}. \quad (3.2.4)$$

4 Completeness of quasimodes via higher order Sturm-Liouville equations

4.1 An abstract linear algebra result

In order to complete the proof of Theorem 1.2.2 for triangular domains, we need to show that the family of quasimodes constructed in subsection 2.2 is complete, and make sure that the numeration of the corresponding eigenvalues is precisely as in (1.2.3). Unlike the general case when we were using Peters solutions, we are well equipped to do that for angles $\pi/2q$ using exponentially accurate quasimodes. We will need the following strengthening of Lemma 2.2.15, formulated as an abstract linear algebra result below.

Theorem 4.1.1. *Let \mathcal{D} be a self-adjoint operator on an infinite-dimensional Hilbert space \mathbf{H} with a discrete spectrum $\{\lambda_j\}_{j=1}^{\infty}$ and a complete orthonormal basis of corresponding eigenvectors $\{\varphi_j\}$. Suppose $\{\mathbf{v}_j\}$ is a sequence of quasimodes such that*

$$\|\mathcal{D}\mathbf{v}_j - \sigma_j\mathbf{v}_j\|_{\mathbf{H}} \leq F(j), \quad (4.1.2)$$

where $\lim_{j \rightarrow \infty} F(j) = 0$. Then

- 1) For all j , there exists k such that $|\sigma_j - \lambda_k| \leq F(j)$;
- 2) For each j , there exists a vector \mathbf{w}_j with the following properties:
 - \mathbf{w}_j is a linear combination of eigenvectors of \mathcal{D} with eigenvalues in the interval $[\sigma_j - \sqrt{F(j)}, \sigma_j + \sqrt{F(j)}]$;
 - $\|\mathbf{w}_j\|_{\mathbf{H}} = 1$;
 - $\|\mathbf{w}_j - \mathbf{v}_j\|_{\mathbf{H}} \leq \sqrt{F(j)} + \sqrt{\frac{F(j)}{1-F(j)}} = 2\sqrt{F(j)}(1 + o(1))$ as $j \rightarrow \infty$.

Proof. Indeed, the first part of the statement is proved using precisely the same arguments as Lemma 2.2.15. Let us prove part 2) of the theorem. Set, similarly to (2.2.14),

$$\mathbf{v}_j = \sum_{k=1}^{\infty} \mathbf{a}_{jk} \varphi_k, \quad \mathbf{a}_{jk} = (\mathbf{v}_j, \varphi_k)_{\mathbf{H}}, \quad \sum_{k=0}^{\infty} \mathbf{a}_{jk}^2 = 1.$$

From the estimate (4.1.2),

$$F(j)^2 \geq \sum_{k: |\lambda_k - \sigma_j| > \sqrt{F(j)}} \mathbf{a}_{jk}^2 (\lambda_k - \sigma_j)^2 \geq F(j) \sum_{k: |\lambda_k - \sigma_j| > \sqrt{F(j)}} \mathbf{a}_{jk}^2, \quad (4.1.3)$$

hence

$$\sum_{k: |\lambda_k - \sigma_j| > \sqrt{F(j)}} \mathbf{a}_{jk}^2 \leq F(j) \Rightarrow 1 - F(j) \leq \sum_{k: |\lambda_k - \sigma_j| \leq \sqrt{F(j)}} \mathbf{a}_{jk}^2 \leq 1. \quad (4.1.4)$$

Letting

$$\mathbf{w}_j := \frac{\sum_{k: |\lambda_k - \sigma_j| \leq \sqrt{F(j)}} \mathbf{a}_{jk} \varphi_k}{\sum_{k: |\lambda_k - \sigma_j| \leq \sqrt{F(j)}} \mathbf{a}_{jk}^2}$$

completes the proof. □

As a consequence of (3.2.1) and (3.2.3), this Lemma may be applied to both the sloshing and ODE problems in the case where both α and β equal $\pi/2q$. In that case, for each j sufficiently large, there exist functions w_j and W_j which are both within $O(e^{-j/C})$ of v_j in L^2 norm (by (3.2.4)), and which are linear combinations of eigenfunctions for the sloshing and ODE problems, respectively, with eigenvalues within $O(e^{-j/C})$ of σ_j . Therefore there is an infinite subsequence, also denoted w_j , of linear combinations of sloshing eigenfunctions which are close to linear combinations of eigenfunctions W_j of the ODE. The problem now is to prove asymptotic completeness and to make sure that the enumeration lines up.

4.2 Eigenvalue asymptotics for higher order Sturm-Liouville problems

Our next goal is to understand the asymptotics of the eigenvalues of the higher order Sturm-Liouville problem (3.2.2) with Neumann boundary conditions. For Dirichlet boundary conditions, the corresponding eigenvalue problem is

$$\begin{cases} (-1)^q U^{(2q)} = \Lambda^{2q} U \\ U^{(m)}(0) = U^{(m)}(1) = 0, m = 0, \dots, q-1. \end{cases} \quad (4.2.1)$$

As was mentioned in subsection 1.5, the Dirichlet and Neumann spectra for the ODE above are related by the following proposition:

Proposition 4.2.2. *The nonzero eigenvalues of the Neumann ODE problem (3.2.2) are the same as those of the Dirichlet ODE problem (4.2.1), including multiplicity. However, the kernel of (4.2.1) is trivial, whereas the kernel of (3.2.2) consists of polynomials of degree less than q and therefore has dimension q .*

Proof. The kernel claims follow from direct computation. For the rest, the solutions with nonzero eigenvalue of the ODE problems (3.2.2) and (4.2.1) must be of the form

$$U(x) = \sum_{k=0}^{2q-1} c_k e^{\omega_k \Lambda x}, \quad (4.2.3)$$

where $\{\omega_1, \dots, \omega_n\}$ are the n th roots of -1 . We claim that a particular $U(x)$ satisfies the Dirichlet boundary conditions if and only if the function

$$\tilde{U}(x) := \sum_{k=0}^{2q-1} c_k \omega_k^q e^{\omega_k \Lambda x}$$

satisfies the Neumann boundary conditions. Indeed, this is obvious; since $\omega_k^{2q} = 1$, for any m and any x , $U^{(m)}(x) = 0$ if and only if $\tilde{U}^{(m+q)}(x) = 0$. This gives a one-to-one correspondence between Dirichlet and Neumann eigenfunctions which completes the proof. □

The following asymptotic result may be extracted, with some extra work (see Appendix B), from a book of M. Naimark [Nai67, Theorem 2, equations (45 a) and (45 b)]. Note that our differential operator has order $n = 2q$, so μ in the notation of [Nai67] is equal to q .

Proposition 4.2.4. *For each $q \in \mathbb{N}$, all sufficiently large eigenvalues of the boundary value problem (3.2.2) are given by the formula*

$$\left\{ \left(k - \frac{1}{2}\right)\pi + O\left(\frac{1}{k}\right) \right\}_{k=K}^{\infty}, \quad (4.2.5)$$

for some $K \in \mathbb{Z}$.

Corollary 4.2.6. *In particular, there exists a $J = J_q \in \mathbb{Z}$ so that for large enough j ,*

$$\Lambda_j \sim \sigma_{j+J} + O\left(\frac{1}{j}\right).$$

This corollary is immediate from the explicit formula for σ_j , with the appropriate values of $\alpha = \beta = \pi/2q$.

Remark 4.2.7. Proposition 4.2.4 implies that λ_j are simple and separated by nearly π for large enough j . However, it does not tell us right away that $J = 0$, and further work is needed to establish this.

The proof of Proposition 4.2.4 is given in Appendix B.

4.3 Connection to sloshing eigenvalues

Recall now that $\{w_j\}$ are linear combinations of sloshing eigenfunctions and $\{W_j\}$ are linear combinations of ODE eigenfunctions, each with eigenvalues in shrinking intervals around σ_j , and each exponentially close to a quasimode. As a consequence of the remark following Corollary 4.2.6, there are gaps between consecutive eigenvalues of the ODE and so for large enough j there is only one eigenvalue of the ODE in each of those intervals. This means that W_j , for sufficiently large j , must actually be an eigenfunction of the ODE, rather than a linear combination of eigenfunctions. By 4.2.6, we must have $W_j = U_{j-J}$, with eigenvalue Λ_{j-J} .

Now consider $\{w_j\}_{j \geq N}$ and $\{W_j\}_{j \geq N}$, and let their spans be \mathbf{X}_N and \mathbf{X}_N^* respectively. Pick N large enough so that W_j are eigenfunctions and so that the intervals in Theorem 4.1.1 are disjoint, and also large enough so that

$$\sum_{j \geq N} \|w_j - W_j\|_{L^2(S)}^2 \leq \sum_{j \geq N} (Ce^{-j/C})^2 < 1. \quad (4.3.1)$$

We know by ODE theory that $\{W_j\}$ form a complete orthonormal basis of $L^2(S)$ and that therefore

$$\dim(\mathbf{X}_N^*)^\perp = N - 1.$$

Lemma 4.3.2. *For sufficiently large N , the dimension $\dim \mathbf{X}_N^\perp = N - 1$ as well.*

Proof. This is essentially a version of the Bary-Krein lemma (see [Kaz71]) and our argument closely follows [Fi16]. Define $\mathbf{A} : \mathbf{X}_N^\perp \rightarrow (\mathbf{X}_N^*)^\perp$ and $\mathbf{B} : (\mathbf{X}_N^*)^\perp \rightarrow \mathbf{X}_N^\perp$ by

$$\mathbf{A}f = f - P_{\mathbf{X}_N^*}f; \quad \mathbf{B}f = f - P_{\mathbf{X}_N}f,$$

where $P_{X_N^*}$ and P_{X_N} are orthogonal projections. By (4.3.1) we have for any $f \in X_N^\perp$:

$$\|P_{X_N^*} f\|^2 = \sum_{j \geq N} |(f, w_j^*)|^2 = \sum_{j \geq N} |(f, w_j^* - w_j)|^2 \leq \epsilon \|f\|^2, \quad \epsilon < 1.$$

Similarly, for any $f \in (X_N^*)^\perp$, $\|P_{X_N} f\| < \epsilon^2 \|f\|^2$. Therefore, for any $f \in X_N^\perp$,

$$BAf = (f - P_{X_N^*} f) - P_{X_N} (f - P_{X_N^*} f) = f - (I - P_{X_N}) P_{X_N^*} f = f - P_{X_N^\perp} P_{X_N^*} f,$$

and hence

$$\|(I_{X_N^\perp} - BA)f\| = \|P_{X_N^\perp} P_{X_N^*} f\| \leq \|P_{X_N^*} f\| < \epsilon \|f\| \text{ for } \epsilon < 1.$$

Therefore, A must be injective, as otherwise $BAf = 0$ for some nonzero f and we get a contradiction. Hence $\dim X_N^\perp \leq \dim (X_N^*)^\perp = N - 1$. Repeating the same argument with X_N and X_N^* interchanged shows that $\dim (X_N^*)^\perp \leq \dim X_N^\perp$, and therefore that

$$\dim X_N^\perp = \dim (X_N^*)^\perp = N - 1,$$

as desired. □

One immediate consequence of this Lemma is that the sequence $\{w_j\}_{j \geq N}$ after a certain point contains only pure eigenfunctions. Indeed, if w_j is a linear combination of $k \geq 2$ eigenfunctions, there are $k - 1$ linearly independent functions generated by the same eigenfunctions, orthogonal to w_j (and all other $\{w_j\}$, since eigenfunctions are orthonormal). But since $\dim X_N^\perp < \infty$, $k = 1$ starting from some $j = N$. Without loss of generality we can pick N sufficiently large so that all w_j are simple for $j \geq N$.

Even more importantly, Lemma 4.3.2 tells us that the sequences $\{w_j\}_{j \geq N}$ and $\{W_j\}_{j \geq N}$ are missing the same number of eigenfunctions, namely $N - 1$. Since we know that W_j corresponds to eigenvalue $\Lambda_{j-J} = \sigma_j + O\left(\frac{1}{j}\right)$, we have proved the following

Proposition 4.3.3. *In the case $\alpha = \beta = \pi/2q$, there exists a constant $C > 0$ and an integer J_q such that both the sloshing eigenvalues λ_j and the ODE eigenvalues Λ_j satisfy*

$$\lambda_j L = \pi \left(j + J_q - \frac{1}{2} - \frac{q}{2} \right) + O(e^{-Cj}) = \Lambda_j L + O(e^{-Cj}), \quad (4.3.4)$$

Remark 4.3.5. Under the same assumptions, asymptotics (4.3.4) holds for the Steklov-Dirichlet eigenvalues λ_j^D and the eigenvalues Λ_j^D of the ODE with Dirichlet boundary conditions (4.2.1) with $-\frac{q}{2}$ being replaced by $+\frac{q}{2}$. Note that the shift J_q in the Dirichlet case (which a priori could be different) is the same as in the Neumann case, as immediately follows from Proposition 4.2.2.

4.4 Proof of Theorem 1.5.4 for triangular domains

We are now in a position to complete the proof of Theorem 1.5.4. Given Proposition 4.3.3, it remains to show that the shift $J_q = 0$ for all $q \in \mathbb{N}$. We prove this by induction. For $q = 1$, by computing the eigenvalues of the standard (second-order) Sturm-Liouville problem, we observe that $J_1 = 0$. In order to make the induction step we use the domain monotonicity properties of Steklov-Neumann and Steklov-Dirichlet eigenfunctions, namely:

Proposition 4.4.1. [BKPS10, section 3] *Suppose Ω_1 and Ω_2 are two domains with the same sloshing surface S , with $\Omega_1 \subseteq \Omega_2$. Then for all $k = 1, 2, \dots$, we have $\lambda_k(\Omega_1) \leq \lambda_k(\Omega_2)$ and $\lambda_k^D(\Omega_1) \geq \lambda_k^D(\Omega_2)$.*

Let Ω_q be an isosceles triangle with angles $\pi/2q$ at the base. We have $\Omega_{q+1} \subset \Omega_q$ for all $k \geq 1$. Assume now $J_{q+1} > J_q = 0$ for some $q > 0$. Then it immediately follows from (4.3.4) that $\lambda_j(\Omega_{q+1}) > \lambda_j(\Omega_q)$ for large j , which contradicts Proposition 4.4.1 for Neumann eigenvalues. Assuming instead that $J_{q+1} < J_q$, we also get a contradiction, this time with Proposition 4.4.1 for Dirichlet eigenvalues. Therefore, $J_q = 0$ for all $k = 1, 2, \dots$, and this completes the proof of Theorem 1.5.4 for triangular domains. \square

Remark 4.4.2. A surprising feature of this proof is that domain monotonicity for mixed Steklov eigenvalues implies new results for the eigenvalues of higher order Sturm-Liouville problems, which a priori are easier to investigate.

4.5 Completeness of quasimodes for arbitrary triangular domains

We now complete the proofs of Theorems 1.2.2 and 1.3.2, as well as Propositions 1.2.6 and 1.3.5, for triangular domains, by taking full advantage of the domain monotonicity properties in Proposition 4.4.1.

Let Ω_s , $s \in [0, 1]$, be a continuous family of sloshing domains (not necessarily triangles) sharing a common sloshing surface S with $L = 1$. Assume that each Ω_s is straight in a neighbourhood of the vertices, with angles $\alpha(s)$ and $\beta(s)$. Moreover, assume that Ω_s is a monotone family, i.e. that $s < t \Rightarrow \Omega_s \subseteq \Omega_t$, and assume that $\alpha(s)$ and $\beta(s)$ are both less than $\pi/2$ for all $0 \leq s < 1$ (possibly equaling $\pi/2$ for $s = 1$).

For the moment we specialize to the Neumann setting; denote the associated quasi-frequencies by $\sigma_j(s)$, and observe by the formula (2.2.1) that they are uniformly equicontinuous in s . By Lemma 2.2.15, for any s and any j , we know that there exist integers $k(j, s)$ such that for j sufficiently large,

$$|\sigma_j(s) - \lambda_{k(j,s)}(s)| = o(1) \tag{4.5.1}$$

By the work of Davis [Dav65], we have a similar bound if $\alpha(s) = \beta(s) = \pi/2$.

The completeness property we need to show translates to the statement that $k(j, s) = j$ for all sufficiently large j , for then the indices match and we have decaying bounds on $|\sigma_j - \lambda_j|$, rather than just $|\sigma_j - \lambda_k|$ for some unknown k .

The key lemma is the following.

Lemma 4.5.2. *Suppose that Ω_s , $s \in [0, 1]$, is a family of sloshing domains as above. Consider the Neumann case. Then*

1. *If $s < s'$ and $k(j, s') \geq j$ for all sufficiently large j , then there exists $N > 0$ so that for all $j \geq N$, $k(j, s) \geq j$.*
2. *If $s < s'$ and $k(j, s) \leq j$ for all sufficiently large j , then there exists $N > 0$ so that for all $j \geq N$, $k(j, s') \leq j$.*
3. *If both $k(j, 0) = j$ and $k(j, 1) = j$ for all sufficiently large j , then for each $s \in [0, 1]$, there exists $N > 0$ so that for all $j \geq N$, $k(j, s) = j$.*

In the Dirichlet case, the same result holds if we flip the inequalities in the conclusion of the first two statements.

Remark 4.5.3. Of course, $k(j, s)$ may not be well-defined for small j , as there may in some cases be more than one value that works, but if the hypotheses are satisfied for some choices of k , then the conclusions must be satisfied for all choices of k .

Proof of Lemma 4.5.2. The third statement is an immediate consequence of the first two (applied with $s' = 1$ and $s = 0$ respectively), so we need only to prove the first two. Since the second one is practically identical to the first one, we only prove the first one here.

First consider the case when s and s' are such that $|\sigma_j(s) - \sigma_j(s')|$ (which is independent of j) is less than π , specifically less than $\pi - \epsilon$ for some $\epsilon > 0$. Then for sufficiently large j , applying Lemma 2.2.15 as in the discussion above, we can arrange both

$$|\lambda_{k(j,s)}(s) - \lambda_{k(j,s')}(s')| < \pi - \epsilon/2 \text{ and } \lambda_{k(j-1,s')}(s') < \lambda_{k(j,s)}(s') - (\pi - \epsilon/2).$$

As a consequence,

$$\lambda_{k(j-1,s')}(s') < \lambda_{k(j,s)}(s).$$

But by domain monotonicity applied to $\lambda_{k(j-1,s')}$, this implies that

$$\lambda_{k(j-1,s')}(s) < \lambda_{k(j,s)}(s),$$

and therefore that $k(j, s) > k(j-1, s')$. Since $k(j-1, s') \geq j-1$ for sufficiently large j by assumption, we must have $k(j, s) > j-1$ and hence $k(j, s) \geq j$, since it is an integer. This is what we wanted.

In the case where s and s' are not such that $|\sigma_j(s) - \sigma_j(s')| < \pi$, simply do the proof in steps: first extend to some $s_1 < s$ with $|\sigma_j(s_1) - \sigma_j(s')| < \pi$, then to some $s_2 < s_1$, et cetera. Since in all cases $|\sigma_j(s) - \sigma_j(s')|$ is finite, this process can be set up to terminate in finitely many steps, completing the proof. \square

Now we can complete the proofs of Theorems 1.2.2 and 1.3.2 for triangular domains. It is possible to embed the triangle Ω as $\Omega_{1/2}$ in a continuous, nested family of sloshing domains $\{\Omega_s\}_{s \in [0,1]}$, where Ω_0 is an isosceles triangle with angles $\pi/2q$ for some large even q , and where Ω_1 is a *rectangle*. For Ω_1 , explicit calculations (see [BKPS10]) show that

$$\lambda_k L = \pi \left(k - \frac{1}{2} \right) - \frac{\pi^2}{8} \left(\frac{2}{\pi} + \frac{2}{\pi} \right) + o(1) = \pi(k-1) + o(1),$$

and therefore that $k(j, 1) = j$ for all sufficiently large j , which is our completeness property. However, we have already proved completeness for Ω_0 , in Theorem 1.5.4, so $k(j, 0) = j$ for all sufficiently large j . And we know we can construct quasimodes on $\Omega_{1/2}$, since it is a triangle, and therefore by Lemma 2.2.15 there exist integers $k(j, 1/2)$ such that

$$|\sigma_j - \lambda_{k(j,1/2)}| = O \left(k^{1 - \frac{\pi/2}{\max\{\alpha, \beta\}}} \right).$$

By statement (3) of Lemma 4.5.2, we have $k(j, 1/2) = j$ for all sufficiently large j . Thus for all sufficiently large k ,

$$\lambda_k L = \sigma_k L + O \left(k^{1 - \frac{\pi/2}{\max\{\alpha, \beta\}}} \right) = \pi \left(k - \frac{1}{2} \right) - \frac{\pi^2}{8} \left(\frac{1}{\alpha} + \frac{1}{\beta} \right) + O \left(k^{1 - \frac{\pi/2}{\max\{\alpha, \beta\}}} \right),$$

which proves Theorem 1.2.2. The proof of Theorem 1.3.2 is essentially identical, with domain monotonicity going in the opposite direction, and the estimate on the error term being modified according to Theorem 2.1.4.

The proofs of Propositions 1.2.6 and 1.3.5 for triangular domains also follow in the same way, by using the model solutions (2.1.7) in the $\mathfrak{S}_{\pi/2}$ in order to construct the quasimodes.

4.6 Completeness of quasi-frequencies: abstract setting

Here, we would like to formulate the results of the previous subsection on the abstract level. We will not need these results in the present paper, but we will need them in the subsequent paper [LPPS17]. The proofs of these results are very similar to the proofs above, and we will omit them.

Let $\Sigma := \{\sigma_j\}$ and $\Lambda := \{\lambda_j\}$, $j = 1, 2, \dots$ be two non-decreasing sequences of real numbers tending to infinity, which are called *quasi-frequencies* and *eigenvalues*, respectively. Suppose that the following “quasi-frequency gap” condition is satisfied: there exists a constant $C > 0$ such that

$$\sigma_{j+1} - \sigma_j > C \tag{4.6.1}$$

for sufficiently large j .

Definition 4.6.2. We say that Σ is a *system of quasi-frequencies approximating eigenvalues* Λ if there exists a mapping $k : \mathbb{N} \rightarrow \mathbb{N}$ such that:

- (i) $|\sigma_j - \lambda_{k(j)}| \rightarrow 0$;
- (ii) $k(j_1) \neq k(j_2)$ for sufficiently large and distinct j_1, j_2 .

Under assumptions (i-ii) and (4.6.1) we obviously have that for j large enough, $\{k(j)\}$ is a strictly increasing sequence of natural numbers. Therefore, if we denote $m(j) := k(j) - j$, then the sequence $\{m(j)\}$ is non-decreasing for j large enough, and, therefore, it is converging (possibly, to $+\infty$). Put

$$M = M(\Sigma, \Lambda) := \lim_{j \rightarrow \infty} m(j).$$

Definition 4.6.3. We say that the system of quasi-frequencies Σ is *asymptotically complete* in Λ if $M \neq +\infty$.

Now we assume that we have one-parameter families $\Sigma(s) = \{\sigma_j(s)\}$ and $\Lambda(s) = \{\lambda_j(s)\}$, $s \in [0, 1]$ of quasi-frequencies and eigenvalues.

Lemma 4.6.4. *Suppose the family $\Sigma(s)$ is equicontinuous and the family $\Lambda(s)$ is monotone increasing (i.e. each $\lambda_j(s)$ is increasing). Then for sufficiently large j (uniformly in s) the function $m(j) = m(j, s)$ is non-increasing in s .*

Proof. The proof is the same as the proof of statement (i) of Lemma 4.5.2. □

Under the conditions of the previous Lemma, the function $M = M(s)$ is non-increasing. Therefore, the following alternative formulation of statement (iii) of Lemma 4.5.2 can be obtained as an immediate consequence:

Corollary 4.6.5. *If $\Sigma(0)$ is asymptotically complete in $\Lambda(0)$, then $\Sigma(1)$ is asymptotically complete in $\Lambda(1)$.*

Remark 4.6.6. With some extra work the “quasi-frequency” gap” assumption (4.6.1) could be weakened, and it is sufficient to require that the number of quasi-frequencies σ_j inside any interval of length one is bounded. More details on that will be provided in [LPPS17].

4.7 Proof of Proposition 1.6.4

Using a similar strategy, let us now prove the asymptotics (1.6.5) for the mixed Steklov-Dirichlet-Neumann problem on triangles.

First, assume that $\alpha = \pi/2$. Consider a sloshing problem on the doubled isosceles triangle $\triangle B'BZ$, where B' is symmetric to B with respect to A . Using the odd-even decomposition of eigenfunctions with respect to symmetry across AZ , we see that the spectrum of the sloshing problem on $\triangle B'BZ$ is the union (counting multiplicity) of the eigenvalues of the sloshing problem and of the mixed Steklov-Dirichlet-Neumann problem on $\triangle ABZ$. Given that we have already computed the asymptotics of the two former problems, it is easy to check that the spectral asymptotics of the latter problem satisfies (1.6.5). A similar reflection argument works in the case when $\beta = \pi/2$ and α is arbitrary. This proves the proposition when either of the angles is equal to $\pi/2$.

Let now $0 < \alpha, \beta \leq \pi/2$ be arbitrary. Choose two additional points X and Y such that X lies on the continuation of the segment BZ , Y lies on the continuation of the segment AZ , and $\angle XAB = \angle YBA = \pi/2$. Consider a family of triangles $\Omega_s = ABP_s$, $0 \leq s \leq 1$, such that $P_0 = X$, $P_1 = Y$, and as s changes, the point P_s continuously moves along the union of segments $[X, Z] \cup [Z, Y]$. In particular, $X = P_0$, $Y = P_1$, and $Z = P_{s_0}$ for some $0 < s_0 < 1$. We impose the Dirichlet condition on $[A, P_s]$ and the Neumann condition on $[B, P_s]$. The Steklov condition on (A, B) does not change.




Figure 4: Construction of auxiliary triangles in the case $\alpha = \pi/2$ (left) and $\alpha < \pi/2$ (right)

For each of the triangles Ω_s one can construct a system of quasimodes similarly to the sloshing or Steklov-Dirichlet case; the only difference is that in this case we use the Dirichlet Peters solutions in one angle and the Neumann Peters solutions in the other.

It remains to show that this system is complete. In order to do that we use the same approach as in Lemma 4.5.2. A simple modification of the argument for the domain monotonicity of the sloshing and Steklov-Dirichlet eigenvalues (see [BKPS10]) yields that for all $k \geq 1$, $\lambda_k(\Omega_s) \leq \lambda_k(\Omega_{s'})$ if $s \leq s'$. Indeed, if $P_s \in [X, Z]$ we can continue any trial function on a smaller triangle by zero across AP_s to get a trial function on a larger triangle, which means that the eigenvalues increase with s . Similarly, if $P_s \in [Z, Y]$, we can always restrict a trial function on a larger triangle to a smaller triangle, and this procedure decreases the Dirichlet energy without changing the denominator in the Rayleigh quotient. Hence, the eigenvalues increase with s in this case as well.

Since we have already proved (1.6.5) for right triangles, in the notation of Lemma 4.5.2 it means that $k(j, 0) = k(j, 1) = j$ for all sufficiently large j . Following the same logic as in this lemma we get that $k(j, s) = s$ for any s , in particular, for $s = s_0$. This completes the proof of the proposition. \square

5 Domains with curvilinear boundary

5.1 Domains which are straight near the surface

Suppose Ω is a sloshing domain whose sides are straight in a neighbourhood of A and B . Let T be the sloshing triangle with the same angles α and β as Ω ; from the results in the previous Sections we understand sloshing eigenvalues and eigenfunctions on T very well. As before, denote the sloshing eigenfunctions on T by $\{u_j(z)\}$, with eigenvalues λ_j . Further, denote the original *uncorrected* quasimodes on T , as in (2.2.2), by $\{v'_{\sigma_j}(z)\}$, with quasi-eigenvalues σ_j . Our strategy will be to use the quasimodes v'_{σ_j} on T , cut off appropriately and modified slightly, as quasimodes for sloshing on Ω .

Indeed, let $\chi(z)$ be a cut-off function on Ω with the following properties, which are easy to arrange:

- $\chi(z)$ is equal to 1 in a neighbourhood of S ;
- $\chi(z)$ is supported on $\Omega \cap T$ and in particular is zero wherever the boundaries of Ω are not straight;
- At every point $z \in \partial\Omega$, $\nabla\chi(z)$ is orthogonal to the normal $\mathbf{n}(z)$ to $\partial\Omega$.

Now define a set of functions on Ω , by

$$\tilde{v}'_{\sigma_j}(z) := \chi(z)v'_{\sigma_j}(z). \quad (5.1.1)$$

The functions $\tilde{v}'_{\sigma_j}(z)$ do not quite satisfy Neumann conditions on the boundary, for the same reason the $v_{\sigma_j}(z)$ do not; we will have to correct them. Note, however, that since $\nabla\chi(z) \cdot \mathbf{n}(z) = 0$,

$$\frac{\partial}{\partial n} \tilde{v}'_{\sigma_j}(z) = \chi(z) \frac{\partial}{\partial n} v'_{\sigma_j}(z), \quad z \in \mathcal{W},$$

and hence, analogously to (2.2.3),

$$\left\| \frac{\partial}{\partial n} \tilde{v}'_{\sigma_j} \right\|_{C^0(\mathcal{W})} \leq C\sigma_j^{-\mu} \quad \text{on } \mathcal{W}.$$

We correct these in precisely the same way as before, by adding a function $\tilde{\eta}_{\sigma_j}$ which is harmonic and which has Neumann data bounded everywhere in C^0 norm by $C\sigma_j^{-\mu}$. As before, since the Neumann-to-Dirichlet map is bounded on $L^2_*(S)$, we have

$$\left\| \frac{\partial}{\partial y} \tilde{\eta}_{\sigma_j} \right\|_{L^2(S)} + \|\tilde{\eta}_{\sigma_j}\|_{L^2(S)} \leq C\sigma_j^{-\mu}.$$

Then we define corrected quasimodes

$$\tilde{v}_{\sigma_j}(z) := \tilde{v}'_{\sigma_j}(z) - \tilde{\eta}_{\sigma_j}(z),$$

which now satisfy Neumann boundary conditions.

We may also compute their Laplacian. Since $v'_{\sigma_j}(z)$ are harmonic functions and $\tilde{\eta}_{\sigma_j}(z)$ are also harmonic, we have

$$\Delta \tilde{v}_{\sigma_j}(z) = \Delta \tilde{v}'_{\sigma_j}(z) = (\Delta \chi(z))v'_{\sigma_j}(z) + 2\nabla \chi(z) \cdot \nabla v'_{\sigma_j}(z). \quad (5.1.2)$$

Therefore $\Delta \tilde{v}_{\sigma_j}(z)$ is supported only where $\nabla \chi(z)$ is nonzero. For later use, we need to estimate the L^2 norm of $\Delta \tilde{v}_{\sigma_j}(z)$. It is immediate from the formula (5.1.2) that for some universal constant C depending on the cutoff function,

$$\|\Delta \tilde{v}_{\sigma_j}(z)\|_{L^2(\Omega)} \leq C\|v'_{\sigma_j}(z)\|_{H^1(T \cap \text{supp}(\nabla \chi))}.$$

The discussion before (2.2.3) shows that the C^1 norm of $v'_{\sigma_j}(z)$ is bounded by $C\sigma_j^{-\mu}$, and therefore obviously

$$\|v'_{\sigma_j}(z)\|_{H^1(T \cap \text{supp}(\nabla \chi))} \leq C\sigma_j^{-\mu}. \quad (5.1.3)$$

Thus

$$\|\Delta \tilde{v}_{\sigma_j}(z)\|_{L^2(\Omega)} \leq C\sigma_j^{-\mu}. \quad (5.1.4)$$

We would like to use $\{\tilde{v}_{\sigma_j}|_S\}$ as our quasimodes. However, since \tilde{v}_{σ_j} are not harmonic functions in the interior, we do not have $\mathcal{D}(\tilde{v}_{\sigma_j}|_S) = (\frac{\partial}{\partial y} \tilde{v}_{\sigma_j})|_S$. We need to correct \tilde{v} to be harmonic. To do this, we must solve the mixed boundary value problem

$$\begin{cases} \Delta \tilde{\varphi}_{\sigma_j} = -\Delta \tilde{v}_{\sigma_j} & \text{on } \Omega; \\ \frac{\partial \tilde{\varphi}_{\sigma_j}}{\partial n} = 0 & \text{on } \mathcal{W}; \\ \tilde{\varphi}_{\sigma_j} = 0 & \text{on } S, \end{cases} \quad (5.1.5)$$

as then we can use

$$\bar{v}_{\sigma_j} := \tilde{v}_{\sigma_j} + \tilde{\varphi}_{\sigma_j}$$

as our quasimodes. The key estimate we want is given by

Lemma 5.1.6. *If $\tilde{\varphi}_{\sigma_j}$ solves (5.1.5), then for some constant C independent of σ_j ,*

$$\left\| \frac{\partial}{\partial n} \tilde{\varphi}_{\sigma_j} \right\|_{L^2(S)} + \|\tilde{\varphi}_{\sigma_j}\|_{L^2(S)} \leq C\sigma^{-\mu}.$$

Proof. The theory of mixed boundary value problems on domains with corners is required here; originally due to Kondratiev [Kon67], it has been developed nicely by Grisvard [Gri85] in the setting of exact polygons. We will therefore transfer our problem to an exact polygon, namely T , via a conformal map. Let $\Phi : \Omega \rightarrow T$ be the conformal map which preserves the vertices and takes an arbitrary point Z' on \mathcal{W} to the vertex Z of the triangle $\triangle ABZ$. Conformal maps preserve Dirichlet and Neumann boundary conditions, so the problem (5.1.5) becomes, with $\varphi = \Phi_*(\tilde{\varphi})$,

$$\begin{cases} \Delta\varphi_{\sigma_j} = -|\Phi'(z)|^2\Delta\tilde{v}_{\sigma_j}(z) & \text{on } T; \\ \frac{\partial\varphi_{\sigma_j}}{\partial n} = 0 & \text{on } \mathcal{W}_T; \\ \varphi_{\sigma_j} = 0 & \text{on } S_T. \end{cases} \quad (5.1.7)$$

By [Gri85, Lemma 4.4.3.1], there is indeed a unique solution $\varphi_{\sigma_j}(z) \in H^1(T)$ to (5.1.7). Moreover, in this setting, the solution is actually in $H^2(T)$. Indeed, since all of the angles of T are less than $\pi/2$ and in particular we do not have any corners of angle exactly $\pi/2$, [Gri85, Theorem 4.4.3.13] applies. It tells us that $\varphi_{\sigma_j}(z)$ equals an element of $H^2(T)$ plus a linear combination of explicit separated-variables solutions at the corners, which Grisvard denotes $S_{j,m}$. However, consulting the definition of $S_{j,m}$ in of [Gri85, equation (4.4.3.7)], it is immediate that each $S_{j,m}$ is itself in $H^2(T)$ (see also [Gri85, Lemma 4.4.3.5]) and therefore $\varphi_{\sigma_j}(z) \in H^2(T)$.

Now that (5.1.7) has a solution $\varphi_{\sigma_j}(z) \in H^2(T)$, we may apply the *a priori* estimate [Gri85, (4.1.2)]. It tells us that

$$\|\varphi_{\sigma_j}\|_{H^2(T)} \leq C (\|\Delta\varphi_{\sigma_j}\|_{L^2(T)} + \|\varphi_{\sigma_j}\|_{L^2(T)}) \leq C\|\Delta\varphi_{\sigma_j}\|_{L^2(T)},$$

where the second inequality follows from the fact that the first eigenvalue of of the Laplacian on T with Dirichlet conditions on S_T and Neumann conditions on \mathcal{W}_T is positive. But since $\varphi_{\sigma_j}(z)$ solves (5.1.7),

$$\|\varphi_{\sigma_j}\|_{H^2(T)} \leq C \|\Phi'(z)|\Delta\tilde{v}_{\sigma_j}(z)\|_{L^2(T)}.$$

Now observe that $|\Phi'(z)|$ is certainly smooth and bounded on the support of $\Delta\tilde{v}_{\sigma_j}(z)$, since that support is away from the vertices. Therefore, by (5.1.4),

$$\|\varphi_{\sigma_j}(z)\|_{H^2(T)} \leq C\sigma^{-\mu}. \quad (5.1.8)$$

This can now be used to complete the proof. By (5.1.8), using the definition of Sobolev norms and the trace restriction Theorem,

$$\left\| \frac{\partial}{\partial y} \varphi_{\sigma_j} \right\|_{L^2(S)} \leq \left\| \frac{\partial}{\partial y} \varphi_{\sigma_j} \right\|_{H^{1/2}(S)} \leq C \left\| \frac{\partial}{\partial y} \varphi_{\sigma_j} \right\|_{H^1(T)} \leq C \|\varphi_{\sigma_j}\|_{H^2(T)} \leq C\sigma^{-\mu}.$$

All that remains to prove Lemma 5.1.6 is to undo the conformal map. By [PP14, Lemma 4.3] (technically the local version, but the proof is local), since Φ preserves the angles at A and B , Φ is $C^{1,\alpha}$ for any α , and hence C^∞ , in a neighbourhood of S , and Φ' is nonzero. Note that Φ may not be C^∞ near Z' in particular, but we do not care about that region. Since Φ is smooth in a neighbourhood of S and Φ' is nonzero, both the measure on S and the magnitude of the normal derivative $\frac{\partial}{\partial y}$ change only up to a constant bounded above and below, and hence, as desired,

$$\left\| \frac{\partial}{\partial n} \tilde{\varphi}_{\sigma_j} \right\|_{L^2(S)} \leq C\sigma_j^{-\mu}.$$

An identical argument without taking $\frac{\partial}{\partial y}$ shows that

$$\|\tilde{\varphi}_{\sigma_j}\|_{L^2(S)} \leq C\sigma_j^{-\mu};$$

in fact, the bound is actually on the $H^{3/2}(S)$ norm. Hence the sum is bounded by $C\sigma_j^{-\mu}$ as well, which proves Lemma 5.1.6. \square

Now we claim our putative quasimodes \bar{v}_{σ_j} on Ω satisfy a quasimode estimate:

Lemma 5.1.9. *There is a constant C such that*

$$\|\mathcal{D}_\Omega(\bar{v}_{\sigma_j}|_S) - \sigma_j\bar{v}_{\sigma_j}\|_{L^2(S)} \leq C\sigma_j^{1-\mu}. \quad (5.1.10)$$

Indeed, \bar{v}_{σ_j} are harmonic and satisfy Neumann conditions on \mathcal{W} , so we know that

$$\mathcal{D}_\Omega(\bar{v}_{\sigma_j}|_S) = \frac{\partial}{\partial y}\bar{v}_{\sigma_j}\Big|_S.$$

As a result,

$$\mathcal{D}_\Omega(\bar{v}_{\sigma_j}|_S) - \sigma_j\bar{v}_{\sigma_j} = \frac{\partial}{\partial y}\tilde{v}_{\sigma_j}\Big|_S - \sigma_j\tilde{v}_{\sigma_j}|_S + \frac{\partial}{\partial y}\tilde{\varphi}_{\sigma_j}\Big|_S - \sigma_j\tilde{\varphi}_{\sigma_j}|_S,$$

which by definition of $\tilde{v}_j\sigma_j$ is

$$\frac{\partial}{\partial y}\tilde{v}'_{\sigma_j}\Big|_S - \sigma_j\tilde{v}'_{\sigma_j}|_S - \frac{\partial}{\partial y}\tilde{\eta}_{\sigma_j}\Big|_S + \sigma_j\tilde{\eta}_{\sigma_j}|_S + \frac{\partial}{\partial y}\tilde{\varphi}_{\sigma_j}\Big|_S - \sigma_j\tilde{\varphi}_{\sigma_j}|_S.$$

The last two terms have L^2 norms which are bounded by $C\sigma^{1-\mu}$, by Lemma 5.1.6. The third and fourth terms satisfy the same bound as a consequence of (5.1.3). For the first two terms, recall that $\tilde{v}'_{\sigma_j} = \chi v'_{\sigma_j}$ and $\chi = 1$ near S , so we need to estimate the L^2 norm of

$$\frac{\partial}{\partial y}v'_{\sigma_j}\Big|_S - \sigma_jv'_{\sigma_j}|_S.$$

However, v'_{σ_j} are the explicit uncorrected quasimodes from (2.2.2). By the proof of Lemma (2.2.6), these two terms both have L^2 norms bounded by $C\sigma_j^{1-\mu}$, completing the proof of (5.1.10).

Now Theorem 4.1.1 applies immediately to tell us that near each sufficiently large quasi-frequency σ_j , there exists at least one sloshing eigenvalue $\lambda_k(\Omega)$. The problem, again, is to prove completeness.

5.2 Completeness

We use the same strategy that we did for our original quasimode problem. Since the strategy is the same, we omit some details.

First we prove completeness when $\alpha = \beta = \pi/2q$. To do this, we instead use Hanson-Lewy quasimodes. All the analysis we have done in the previous subsection goes through with $Ce^{-\sigma/C}$ replacing $C\sigma^{-\mu}$ everywhere, and we get (5.1.10) with a right-hand side of $Ce^{-j/C}$. So Theorem

4.1.1 applies. It shows in particular that there are (linear combinations of) sloshing eigenfunctions on Ω , which we call $w_{j,\Omega}$, with eigenvalues exponentially close to σ_j and with the property that

$$\|w_{j,\Omega} - \bar{v}_{\sigma_j}\|_{L^2} \leq C e^{-j/C} \text{ as } j \rightarrow \infty.$$

However $\|\bar{v}_{\sigma_j} - \tilde{v}_{\sigma_j}\|_{L^2(S)}$, $\|\tilde{v}_{\sigma_j} - v'_{\sigma_j}\|_{L^2(S)}$, $\|v'_{\sigma_j} - v_{\sigma_j}\|_{L^2(S)}$, and $\|v_{\sigma_j} - u_j\|_{L^2(S)}$ are all bounded by $C e^{-j/C}$ themselves in this case, so we have in all that

$$\|w_{j,\Omega} - u_j\|_{L^2} \leq C e^{-j/C} \text{ as } j \rightarrow \infty.$$

We know that $\{u_j\}$ are complete. By the same argument as in Lemma 4.3.2, the sequence $\{w_{j,\Omega}\}$ must at some point contain only pure eigenfunctions, and for suitably large N the sequences $\{w_{j,\Omega}\}_{j \geq N}$ and $\{u_j\}_{j \geq N}$ are missing the same number of eigenfunctions, which shows that

$$\lambda_{j,\Omega} = \lambda_{j,T} + O(e^{-j/C}).$$

This proves completeness, along with full exponentially accurate eigenvalue asymptotics, in the $\alpha = \beta = \pi/2q$ case.

Finally, we use domain monotonicity to deal with general α and β ; the argument runs exactly as in subsection 4.5. The result of Davis [Dav65] shows completeness in the $\pi/2$ case, and we can sandwich our domain Ω as $\Omega_{1/2}$ in a continuous, nested family of sloshing domains $\{\Omega_s\}$ that satisfy the hypotheses of Lemma 4.5.2. Note that Lemma 4.5.2 is stated in such a way as to apply here as well.

5.3 General curvilinear domains

We now generalize further, to domains with curvilinear boundary. The authors are grateful to Lev Buhovski for suggesting the approach in this subsection [Buh17].

Suppose that Ω is a sloshing domain which satisfies the following conditions:

- (C1) Ω is simply connected;
- (C2) The walls \mathcal{W} are Lipschitz;
- (C3) For any $\epsilon > 0$, there exist sloshing domains Ω_- and Ω_+ with piecewise smooth boundary, with $\Omega_- \subset \Omega \subset \Omega_+$, and with the additional property that Ω_- and Ω_+ are straight lines in a δ -neighbourhood of A and B with vertex angles in $[\alpha - \epsilon, \min\{\alpha + \epsilon, \pi/2\}]$.

The point of this third condition is that we have already proved sloshing eigenvalue asymptotics for Ω_- and Ω_+ , which we will need in the subsequent domain monotonicity argument. This third condition is satisfied, for example, if the boundary S is C^1 in any small neighbourhood of the vertices A and B and the angles are strictly less than $\pi/2$. It is also satisfied under the ‘‘local John’s condition’’ (see Propositions 1.2.6 and 1.3.5) if one or both angles equals $\pi/2$, and makes clear why that condition is necessary: we have not proved sloshing asymptotics in the case where one or both angles are greater than $\pi/2$.

Under these conditions, we claim the asymptotics (1.2.3), (1.3.3), as well as (1.2.7) and (1.3.6) for domains satisfying the additional ‘‘local John’s condition’’, and thereby complete the proofs of Theorems 1.2.2 and 1.3.2 as well as Propositions 1.2.6 and 1.3.5.

Proof. Assume $L = 1$ for simplicity. Pick any $\gamma > 0$. By continuity, there exists a sufficiently small ϵ such that for any $\epsilon' \in [-\epsilon, \epsilon]$,

$$\left| \frac{1}{\alpha + \epsilon'} - \frac{1}{\alpha} \right| + \left| \frac{1}{\beta + \epsilon'} - \frac{1}{\beta} \right| < \frac{8\gamma}{2\pi^2}.$$

By condition (C3) above, there exist $\Omega_- \subset \Omega$ and $\Omega_+ \supset \Omega$ as described. By the definition of ϵ , and the explicit form of σ_j ,

$$|\sigma_j(\Omega_+) - \sigma_j(\Omega)| < \gamma/2 \text{ and } |\sigma_j(\Omega_-) - \sigma_j(\Omega)| < \gamma/2 \text{ for all } j.$$

But by our previous work, we have eigenvalue asymptotics for Ω_+ and Ω_- . In particular, there exists N so that for all $j \geq N$,

$$|\lambda_j(\Omega_+) - \sigma_j(\Omega_+)| < \gamma/2 \text{ and } |\lambda_j(\Omega_-) - \sigma_j(\Omega_-)| < \gamma/2.$$

And by domain monotonicity (for Neumann — inequalities reverse for the Dirichlet case),

$$\lambda_j(\Omega_-) \leq \lambda_j(\Omega) \leq \lambda_j(\Omega_+) \text{ for all } j.$$

Putting these ingredients together, for $j \geq N$,

$$\lambda_j(\Omega) \leq \lambda_j(\Omega_+) < \sigma_j(\Omega_+) + \frac{\gamma}{2} \leq \sigma_j(\Omega) + \gamma.$$

Similarly,

$$\lambda_j(\Omega) \geq \lambda_j(\Omega_-) > \sigma_j(\Omega_-) - \frac{\gamma}{2} \geq \sigma_j(\Omega) - \gamma.$$

We conclude that for $j \geq N$, $|\lambda_j(\Omega) - \sigma_j(\Omega)| < \gamma$. Since $\gamma > 0$ was arbitrary, we have the asymptotics with $o(1)$ error that we want. \square

A Proof of Theorem 2.1.4

In this section, we prove Theorem 2.1.4. The proof follows [Pet50] for the most part, doing the extra work needed to prove the careful remainder estimates. There is a key difference of notation: throughout, we use $\mu_\alpha = \mu = \pi/(2\alpha)$, where Peters uses $\mu = \pi/\alpha$.

A.1 Robin-Neumann problem

Following Peters, we complexify our problem by setting $z = \rho e^{i\theta}$ and look for an analytic function $f(z)$ in \mathfrak{S}_α with $\phi = \text{Re}(f)$. The boundary conditions must be rewritten in terms of f . Using the Cauchy-Riemann equations we have $\phi_y = \text{Re}(if'(z))$ and $\phi_x = \text{Re}(f'(z))$. After some algebraic transformations, which we skip for brevity, the problem (2.1.1) becomes

$$\begin{cases} f(z) & \text{is analytic in } \mathfrak{S}_\alpha, \\ \text{Re}(if'(z)) = \text{Re}(f(z)) & \text{for } z \in \mathfrak{S}_\alpha \cap \{\theta = 0\}, \\ \text{Re}(ie^{-i\alpha} f'(z)) = 0 & \text{for } z \in \mathfrak{S}_\alpha \cap \{\theta = -\alpha\}. \end{cases} \quad (\text{A.1.1})$$

A.2 Peters solution

We now write down Peters solution, in a form due to Alker [Alk77]. Note that Alker's y -axis points in the opposite direction from Peters and so we have modified the expression in [Alk77] accordingly. First define the auxiliary function

$$I_\alpha(\zeta) := \frac{1}{\pi} \int_0^{\zeta^\infty} \log(1 + v^{-\pi/\alpha}) \frac{\zeta}{v^2 + \zeta^2} dv. \quad (\text{A.2.1})$$

Note that $I_\alpha(\zeta)$ is defined for $\arg(\zeta) \in (-\alpha, \alpha)$, which includes the real axis. As in the appendix of [Pet50], $I_\alpha(\zeta)$ has a meromorphic continuation, with finitely many branch points, to the entire complex plane, which we also call $I_\alpha(\zeta)$. Each branch point is logarithmic; there is one at the origin, and others on the unit circle in the negative real half-plane. We then let

$$g_\alpha(\zeta) = \exp\left(-I_\alpha(\zeta e^{-i\alpha}) + \log\left(\frac{\zeta + i}{\zeta}\right)\right). \quad (\text{A.2.2})$$

The function $g_\alpha(\zeta)$ is originally defined for $\arg(\zeta) \in (0, 2\alpha)$, but has a meromorphic continuation to the entire complex plane minus a single branch cut from the origin, with singularities along the portion of the unit circle outside the sector $-\pi/2 - \alpha \leq \arg(\zeta) \leq \pi/2 + \alpha$. For $\text{Re}(\zeta) > 0$ we have the representation formula [Pet50, (4.8)]

$$g_\alpha(\zeta) = \exp\left(-\frac{1}{\pi} \int_0^\infty \log\left(\frac{1 - t^{-2\mu}}{1 - t^{-2}}\right) \frac{\zeta}{t^2 + \zeta^2} dt\right). \quad (\text{A.2.3})$$

Finally, let P be a keyhole path, consisting of the union of a nearly full circle of radius 2, traversed counterclockwise, and two linear paths, one on each side of the angle $\theta = \pi + \alpha/2$. Then Peters solution is given by

$$f(z) = \frac{\mu^{1/2}}{i\pi} \int_P \frac{g_\alpha(\zeta)}{\zeta + i} e^{z\zeta} d\zeta. \quad (\text{A.2.4})$$

A.3 Verification of Peters solution

Let us verify that Peters solution, which is obviously analytic, is actually a solution by checking the boundary conditions in (A.1.1), beginning with the one along the real axis. This follows [Pet50, section 6].

By [Pet50, p. 335], $g_\alpha(\zeta) \rightarrow 1$ as $\zeta \rightarrow \infty$ (see Proposition A.5.1 for a rigorous proof), and since $e^{z\zeta}$ is very small as $\zeta \rightarrow \infty$ along P , differentiation under the integral sign in (A.2.4) is justified. We get

$$if'(z) - f(z) = \frac{\mu^{1/2}}{i\pi} \int_P \frac{g_\alpha(\zeta)(i\zeta - 1)}{\zeta + i} e^{z\zeta} d\zeta = \frac{\mu^{1/2}}{\pi} \int_P g_\alpha(\zeta) e^{z\zeta} d\zeta. \quad (\text{A.3.1})$$

We claim this integral is pure imaginary whenever z is on the positive real axis. Indeed, we may shift the branch cut of $g_\alpha(\zeta)$ so that it lies along the negative real axis, and shift P so that it is symmetric with respect to the real axis. Observe that $g_\alpha(\zeta)$ is real whenever ζ is real and positive. By reflection, this implies that $g_\alpha(\bar{\zeta}) = \overline{g_\alpha(\zeta)}$ for all ζ . The analogous statement is true

for $e^{z\zeta}$ and hence for $g_\alpha(\zeta)e^{z\zeta}$, and it follows immediately from symmetry of P that (A.3.1) is purely imaginary. Thus $\operatorname{Re}(if'(z) - f(z)) = 0$, as desired.

For the other boundary condition, we compute

$$ie^{-i\alpha}f'(z) = \frac{\mu^{1/2}}{\pi} \int_P \frac{g_\alpha(\zeta)e^{-i\alpha}\zeta}{\zeta + i} e^{z\zeta} d\zeta, \quad (\text{A.3.2})$$

and claim that this is pure imaginary when $\arg(z) = -\alpha$ - i.e. when $z = \rho e^{-i\alpha}$ for some $\rho > 0$. Shifting the branch cut and contour of integration, and letting $w = \zeta e^{-i\alpha}$, the integral (A.3.2) becomes

$$\frac{\mu^{1/2}}{\pi} \int_{P_1} \frac{g_\alpha(we^{i\alpha})we^{i\alpha}}{we^{i\alpha} + i} e^{\rho w} dw = \frac{\mu^{1/2}}{\pi} \int_{P_1} \exp\{-I_\alpha(w)\} e^{\rho w} dw, \quad (\text{A.3.3})$$

where we have chosen P_1 so that it is symmetric with respect to the real axis and the branch cut is along the negative real axis. The boundary condition follows immediately as above, since from (A.2.1) $I_\alpha(w)$ is positive for w real and positive.

A.4 Asymptotics of Peters solution as $z \rightarrow \infty$

Now we study the asymptotics as $z \rightarrow \infty$ of (A.2.4), following [Pet50, section 7] but with more rigour. Let, as before,

$$\chi = \chi_{\alpha,N} = \frac{\pi}{4}(1 - \mu) = \frac{\pi}{4} \left(1 - \frac{\pi}{2\alpha}\right).$$

Theorem A.4.1. *There exist a function $E : (0, \pi/2) \rightarrow \mathbb{R}$ and a complex-valued function $R_\alpha(z)$ depending on α such that*

$$f(z) = e^{E(\alpha)}e^{-i(z-\chi)} + R_\alpha(z), \quad (\text{A.4.2})$$

where for any fixed $\alpha \in (0, \pi/2)$ there exists a constant C such that for all $z \in \mathfrak{S}_\alpha$,

$$|R_\alpha(z)| \leq Cz^{-\mu}, \quad |\nabla_z R_\alpha(z)| \leq Cz^{-\mu-1}. \quad (\text{A.4.3})$$

Remark A.4.4. A slightly stronger version of this theorem (identifying $E(\alpha)$) is claimed in [Alk77]. However, the proof is not given and the extraction does not seem obvious, although it is numerically clear. So we prove this version, which is all we need since we may scale by an overall constant anyway.

Remark A.4.5. For the solution $\phi = \operatorname{Re}(f(z))$ to our original problem, we see that

$$\phi(x, 0) = e^{E(\alpha)} \cos(z - \chi) + R_\alpha(x + 0i), \quad (\text{A.4.6})$$

which is the radiation condition we wanted in the first place.

Proof. The portion of the integral (A.2.4) along the infinite line segments decays exponentially in $|z|$ (note that $\arg(z\zeta)$ is between $\pi - \alpha/2$ and $\pi + \alpha/2$). So we may deform the contour P to a contour P' consisting of the union of a circle of radius $1/2$ and line segments along each side of the branch cut. This deformation passes through singularities of $g_\alpha(\zeta)/(\zeta + i)$, one at

$\zeta = -i$ and others at $\zeta = \lambda_1, \dots, \lambda_m$ along the unit circle in the negative real half-plane, with $\arg(\lambda_j) \notin (-\pi/2 - \alpha, \pi/2 + \alpha)$. We see that

$$\begin{aligned} f(z) &= 2\mu^{1/2}g_\alpha(-i)e^{-iz} + \frac{\mu^{1/2}}{i\pi} \int_{P'} \frac{g_\alpha(\zeta)}{\zeta + i} e^{z\zeta} d\zeta \\ &\quad + \sum_{j=1}^m e^{z\lambda_j} 2\mu^{1/2} \operatorname{Res}_{\zeta=\lambda_j} (g_\alpha(\zeta)/(\zeta + i)). \end{aligned} \tag{A.4.7}$$

Let

$$\begin{aligned} \tilde{R}_\alpha(z) &:= \frac{\mu^{1/2}}{i\pi} \int_{P'} \frac{g_\alpha(\zeta)}{\zeta + i} e^{z\zeta} d\zeta; \\ R_\alpha(z) &:= \tilde{R}_\alpha(z) + \sum_{j=1}^m e^{z\lambda_j} 2\mu^{1/2} \operatorname{Res}_{\zeta=\lambda_j} (g_\alpha(\zeta)/(\zeta + i)). \end{aligned} \tag{A.4.8}$$

To prove Theorem A.4.1 we must first prove the error estimates for $R_\alpha(z)$ and then compute $g_\alpha(-i)$.

A.4.1 Error estimates for $R_\alpha(z)$

Consider $R_\alpha(z)$. Since $\arg(z) \in (-\alpha, 0)$, the finite sum of exponentials and its gradient decay exponentially as $|z| \rightarrow \infty$. The decay is actually uniform in α for α in any interval $(\epsilon, \pi - \epsilon)$ with $\epsilon > 0$ fixed, as the number of residues is bounded in such an interval as well. Only $\tilde{R}_\alpha(z)$ remains. Using (A.2.1), we may write

$$\tilde{R}_\alpha(z) = \frac{\mu^{1/2}}{i\pi} \int_{P'} e^{z\zeta} \exp(-I_\alpha(\zeta e^{-i\alpha})) \frac{d\zeta}{\zeta}. \tag{A.4.9}$$

The only singularity inside P' is along the branch cut, including the branch point 0. We see that we must understand the asymptotics of $I_\alpha(\zeta)$ as $\zeta \rightarrow 0$. Peters identifies the leading order term of $I_\alpha(\zeta)$ at $\zeta = 0$ as $-\mu \log \zeta$, but we need to understand the remainder and prove bounds on it and its gradient. Throughout, we choose to suppress the α subscripts. The key is the following lemma:

Lemma A.4.10. *For ζ in a small neighbourhood of zero, with argument up to the branch cut on either side, we have*

$$I(\zeta) = -\mu \log \zeta + p(\zeta) + R(\zeta),$$

where $p(\zeta)$ is a polynomial in ζ with $p(0) = 0$, and where there is a constant C such that

$$|R(\zeta)| \leq C|\zeta|^{-2\mu}.$$

Proof. The proof proceeds in two steps. First we prove this for all ζ with argument in a compact subset of $(-\pi/2, \pi/2)$. Then we use a functional relation satisfied by $I(\zeta)$ to extend to ζ with arguments up to and into the negative half-plane.

For the first step, we use the representation for $I(\zeta)$ on p. 353 of [Pet50]. From the last paragraph of this page, we have

$$I(\zeta) = -\mu \log \zeta + \int_0^{1/2} \ln(1 + v^{2\mu}) \frac{\zeta}{v^2 + \zeta^2} dv + \int_{1/2}^{\infty} \ln(1 + v^{2\mu}) \frac{\zeta}{v^2 + \zeta^2} dv.$$

The last term is convergent, as $\ln(1 + v^{2\mu}) \sim 2\mu \ln v$ as $v \rightarrow \infty$, and differentiation in ζ under the integral sign is easily justified. So the last term is holomorphic, and by direct substitution it is zero at $\zeta = 0$. For the second term, we use a Taylor expansion for $\ln(1 + v^{2\mu})$, which is convergent when $v < 1$, and obtain

$$\int_0^{1/2} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{v^{2\mu n}}{n} \frac{\zeta}{v^2 + \zeta^2} dv.$$

Change variables in the integral to $w = \zeta v$; we end up with

$$\int_0^{1/2\zeta} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n} \zeta^{2\mu n} w^{2\mu n} \frac{1}{w^2 + 1} dw. \quad (\text{A.4.11})$$

Now further break up the integral (A.4.11), at $w = 2$. The integral from 0 to 2 is bounded in absolute value by

$$\int_0^2 \sum_{n=1}^{\infty} |\zeta|^{2\mu n} \frac{2^{2\mu n}}{n} dw = 2 \sum_{n=1}^{\infty} |\zeta|^{2\mu n} \frac{2^{2\mu n}}{n} \leq 2 \sum_{n=1}^{\infty} |2\zeta|^{2\mu n} = \frac{|4\zeta|^{2\mu}}{1 - |2\zeta|^{2\mu}},$$

which is bounded by $C|\zeta|^{2\mu}$ for some universal constant C and sufficiently small $|\zeta|$. For the remainder of the integral, we use the Taylor expansion of $(w^2 + 1)^{-1}$ about infinity, which is valid for $w > 1$: it is $w^{-2} - w^{-4} + w^{-6} + \dots$, and we end up with

$$\int_2^{1/2\zeta} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n} \zeta^{2\mu n} w^{2\mu n} \left(\sum_{m=1}^{\infty} (-1)^{m+1} w^{-2m} \right) dw.$$

Now observe that $|\zeta w| \leq \frac{1}{2}$ on this interval and $w > 2$, so both the sums in n and m are absolutely convergent and in fact the double sum is absolutely convergent. This justifies all rearrangements as well as term-by-term integration. Our last remaining piece thus becomes

$$\sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{(-1)^{n+m}}{n} \zeta^{2\mu n} \int_2^{1/2\zeta} w^{2\mu n - 2m} dw,$$

which equals

$$\begin{aligned} & \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{(-1)^{n+1+m}}{n} \zeta^{2\mu n} \left(\frac{w^{2\mu n - 2m + 1}}{2\mu n - 2m + 1} \right) \Big|_2^{1/2\zeta} \\ &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{(-1)^{n+1+m}}{n(2\mu n - 2m + 1)} \zeta^{2\mu n} (\zeta^{-2\mu n + 2m - 1} 2^{-2\mu n + 2m - 1} - 2^{2\mu n - 2m + 1}) \\ &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{(-1)^{n+1+m}}{n(2\mu n - 2m + 1)} (\zeta^{2m-1} 2^{-2\mu n + 2m - 1} - \zeta^{2\mu n} 2^{2\mu n - 2m + 1}). \end{aligned}$$

The sum now has two terms. The second term, with a $\zeta^{2\mu n}$, can be summed first in m (obviously) and then in n , and the whole thing is bounded by a multiple of the $n = 1$ term, namely $C|\zeta|^{2\mu}$. The first term can be summed first in n , at which point it becomes an expression of the form $\sum_{m=1}^{\infty} a_m \zeta^{2m-1}$. Since the a_m do not grow too fast as $m \rightarrow \infty$ - in fact they grow as 2^{2m} - this expression represents a holomorphic function in a disk of sufficiently small radius about the origin. Moreover this holomorphic function is zero at zero.

We have now shown that $I(\zeta)$ is the sum of $-\mu \log \zeta$, a holomorphic function zero at the origin, and a term bounded in absolute value by $C|\zeta|^{2\mu}$. Taking the Taylor series of the holomorphic function, separating out the finitely many terms which are not $O(|\zeta|^{2\mu})$, and calling them $p(\zeta)$ completes the proof.

It remains to extend the argument outside the positive real half-plane. Let $h(\zeta) = I(\zeta) + \mu \log \zeta$. Then we know that $h(\zeta)$ is holomorphic in a disk with the exception of a branch cut, and it continues across the branch cut because $I(\zeta)$ does. We also know, from the first line on p. 353 of [Pet50], that for all ζ away from the branch cut,

$$I(\zeta e^{2i\alpha}) = I(\zeta) \frac{\zeta e^{i\alpha} + i}{\zeta e^{i\alpha} - i}.$$

Applying this gives an equivalent relation for $h(\zeta)$, noting that $e^{2i\alpha\mu} = e^{i\pi} = -1$:

$$h(\zeta e^{2i\alpha}) = -h(\zeta) \frac{\zeta e^{i\alpha} + i}{\zeta e^{i\alpha} - i}.$$

The function $H(\zeta) := -(\zeta e^{i\alpha} + i)/(\zeta e^{i\alpha} - i)$ is holomorphic in the disk, and we have for all nonzero ζ :

$$h(\zeta e^{2i\alpha}) = h(\zeta)H(\zeta).$$

Recall that in the positive real half-plane (in a sector bounded away from the real axis),

$$h(\zeta) = p(\zeta) + R(\zeta).$$

Plugging in the relation for $h(\zeta)$, we see that for $-\pi/2 + \epsilon < \arg \zeta < \pi/2 - \epsilon$,

$$h(\zeta e^{2i\alpha}) = p(\zeta)H(\zeta) + R(\zeta)H(\zeta).$$

However, we are assuming that $\alpha < \pi/2$. Therefore, there is a range of ζ , namely $-\pi/2 + \epsilon < \arg \zeta < \pi/2 - \epsilon - 2\alpha$, where $\zeta e^{2i\alpha}$ is in the positive half-plane and thus we have a second representation:

$$h(\zeta e^{2i\alpha}) = p(\zeta e^{2i\alpha}) + R(\zeta e^{2i\alpha}).$$

Setting the previous two equations equal, we see that for this small range of ζ :

$$p(\zeta e^{2i\alpha}) - p(\zeta)H(\zeta) = R(\zeta)H(\zeta) - R(\zeta e^{2i\alpha}).$$

The left-hand side is holomorphic in a disk. The right-hand side is $O(|\zeta|^{2\mu})$. We conclude that the left-hand side must have a zero of order at least 2μ at $\zeta = 0$. Therefore, there is a C such that for *all* ζ in the disk,

$$|p(\zeta e^{2i\alpha}) - p(\zeta)H(\zeta)| \leq C|\zeta|^{2\mu}.$$

Therefore, for $-\pi/2 + \epsilon < \arg \zeta < \pi/2 - \epsilon$,

$$|p(\zeta e^{2i\alpha}) - h(\zeta e^{2i\alpha}) + R(\zeta)H(\zeta)| \leq C|\zeta|^{2\mu},$$

and therefore

$$h(\zeta e^{2i\alpha}) = p(\zeta e^{2i\alpha}) + O(|\zeta|^{2\mu}).$$

This proves the lemma for ζ with arguments now up to $\pi/2 - \epsilon + 2\alpha$. Continuing this procedure, one step of size 2α at a time, gives the result for all ζ with arguments in a neighbourhood of $[-\pi, \pi]$, completing the proof. \square

Now we return to estimating the remainder. Using the Lemma and recalling $h(\zeta) = I(\zeta) + \mu \log \zeta$, we have

$$\tilde{R}_\alpha(z) = -\frac{\mu^{1/2}}{\pi} \int_{P'} e^{z\zeta} \zeta^\mu e^{-h(\zeta)} \frac{d\zeta}{\zeta}. \quad (\text{A.4.12})$$

Change variables to let $w = z\zeta$; for $|z| > 1$, the contour deforms smoothly back to $\arg(z)P'$ (which is P' rotated by $\arg(z)$), as there are no singularities remaining inside P' . We get

$$\tilde{R}_\alpha(z) = -\frac{\mu^{1/2} z^{-\mu}}{\pi} \int_{\arg(z)P'} e^w w^{\mu-1} e^{-h(w/z)} dw. \quad (\text{A.4.13})$$

The contour can be deformed to two straight lines, one on either side of the branch cut. Since $h(\cdot)$ grows at most logarithmically along these lines (note that $\arg(w/z) = \pm\pi + \alpha/2$ on $\arg(z)P'$), the integral is bounded as $z \rightarrow \infty$. In fact, since $h(0) = 0$, it converges to the corresponding integral with $e^{-h(w/z)}$ replaced by 1. Thus $|\tilde{R}_\alpha(z)| \leq Cz^{-\mu}$ as desired.

We must also estimate $|\nabla_z \tilde{R}_\alpha(z)|$. However, $\tilde{R}_\alpha(z)$ is holomorphic, so we just need to consider $\partial_z \tilde{R}_\alpha(z)$. The differentiation brings down a factor of ζ , which becomes w/z . The extra power of w is absorbed, and the extra power of z moves outside the integral, yielding decay of the form $z^{-\mu-1}$ instead of $z^{-\mu}$. This proves the remainder estimation part of Theorem A.4.1. It remains only to evaluate $g_\alpha(-i)$.

A.4.2 Evaluation of $g_\alpha(-i)$

We cannot simply use the formula (A.2.3) to evaluate $g_\alpha(-i)$, because that formula is only valid for ζ in the positive real half-plane. We must instead use the fact that

$$\log g_\alpha(\zeta) = \log\left(\frac{\zeta + i}{\zeta}\right) - I(\zeta e^{-i\alpha}), \quad (\text{A.4.14})$$

so we are interested in understanding the behavior of $I(\zeta)$ near $\zeta = -ie^{-i\alpha}$.

First we need a good representation for $I(\zeta)$ in this region. Considering the last equation on p. 350 of [Pet50], we see that for ζ with $-\pi/2 - 2\alpha < \arg(\zeta) < \pi/2 + 2\alpha$,

$$\begin{aligned} I(\zeta e^{-i\alpha}) &= \frac{1}{2\pi i} \left(\int_{I_1} \log(1 + (-iu)^{-2\mu}) \frac{du}{u - \zeta e^{-i\alpha}} + \int_{M_2} \log(1 + (iu)^{-2\mu}) \frac{du}{u - \zeta e^{-i\alpha}} \right) \\ &\quad + \log \frac{\zeta + i}{\zeta}. \end{aligned}$$

Here I_1 may be chosen to be the incoming path along $\arg(\zeta) = \pi/2$ and M_2 the outgoing path along $\arg(\zeta) = -\pi/2 - 2\alpha$ (each can be moved by up to α in either direction before running into a singularity of the integrand, but these are the most convenient choices). We would prefer that I_1 and M_2 be equal and opposite paths, but they are not. So we shift the path I_1 to the path L_0 which is incoming along the ray $\arg(\zeta) = \pi/2 - 2\alpha$. This is done for other reasons on p. 351 of [Pet50], and we pick up a contribution from the branch cut of the integrand. Overall we get

$$I(\zeta e^{-i\alpha}) = \frac{1}{2\pi i} \left(\int_{L_0} \log(1 + (-iu)^{-2\mu}) \frac{du}{u - \zeta e^{-i\alpha}} + \int_{M_2} \log(1 + (iu)^{-2\mu}) \frac{du}{u - \zeta e^{-i\alpha}} \right) + \log \frac{\zeta + i}{\zeta} - \log \frac{\zeta - i}{\zeta}. \quad (\text{A.4.15})$$

Using (A.4.14), then plugging in $\zeta = -i$, shows that

$$-\log(g_\alpha(-i)) = \frac{1}{2\pi i} \left(\int_{L_0} \log(1 + (-iu)^{-2\mu}) \frac{du}{u + ie^{-i\alpha}} + \int_{M_2} \log(1 + (iu)^{-2\mu}) \frac{du}{u + ie^{-i\alpha}} \right) - \log 2. \quad (\text{A.4.16})$$

Now we parametrise these integrals. For the first, let $u = tie^{-2i\alpha}$, and for the second, let $u = -tie^{-2i\alpha}$. With the appropriate signs, we find that

$$\begin{aligned} -\log(g_\alpha(-i)) &= -\log 2 + \frac{1}{2\pi i} \int_0^\infty \log(1 + t^{-2\mu}) \left(\frac{1}{t - e^{i\alpha}} - \frac{1}{t + e^{i\alpha}} \right) dt \\ &= -\log 2 + \frac{1}{\pi i} \int_0^\infty \log(1 + t^{-2\mu}) \frac{e^{i\pi/2\mu}}{t^2 - e^{i\pi/\mu}} dt. \end{aligned} \quad (\text{A.4.17})$$

We have now written $g_\alpha(-i)$ in terms of a convergent integral, that on the right of (A.4.17). The usual Taylor series expansions show that this integral is smooth in α for $\alpha \in (\epsilon, \pi - \epsilon)$. This integral seems nontrivial to evaluate and is not obviously in any common table of integrals. Nevertheless, we can find the imaginary part of $\log(g_\alpha(-i))$ by using the following lemma.

Lemma A.4.18. *For $\mu \in \mathbb{R}$, let*

$$J(\mu) := \int_0^\infty \log(1 + t^{-2\mu}) \frac{e^{i\pi/2\mu}}{t^2 - e^{i\pi/\mu}} dt. \quad (\text{A.4.19})$$

Then for each $\mu > 1/2$, which corresponds exactly to $\alpha < \pi/2$,

$$\operatorname{Re}(J(\mu)) = \frac{\pi^2}{4}(1 - \mu). \quad (\text{A.4.20})$$

Proof. Change variables in $J(\mu)$ by letting $t = r^{-1}$. Then $dt = -r^{-2}dr$, and we have

$$J(\mu) = \int_0^\infty \log(1 + r^{2\mu}) \frac{e^{i\pi/2\mu}}{1 - r^2 e^{i\pi/\mu}} dr = - \int_0^\infty \log(1 + r^{2\mu}) \frac{e^{-i\pi/2\mu}}{r^2 - e^{-i\pi/\mu}} dr. \quad (\text{A.4.21})$$

Breaking up the logarithm by bringing out $2\mu \log r$ gives

$$J(\mu) = - \int_0^\infty \log(1 + r^{-2\mu}) \frac{e^{-i\pi/2\mu}}{r^2 - e^{-i\pi/\mu}} dr - 2\mu \int_0^\infty \log(r) \frac{e^{-i\pi/2\mu}}{r^2 - e^{-i\pi/\mu}} dr. \quad (\text{A.4.22})$$

But we now recognise the first integral as the complex conjugate of $J(\mu)$. Therefore

$$\operatorname{Re}(J(\mu)) = \frac{1}{2}(J(\mu) + \overline{J(\mu)}) = -\mu \int_0^\infty \log(r) \frac{e^{-i\pi/2\mu}}{r^2 - e^{-i\pi/\mu}} dr. \quad (\text{A.4.23})$$

The integral (A.4.23) may now be evaluated explicitly. It is the integral along the positive real axis of the function $\log(z) \frac{e^{-i\pi/2\mu}}{z^2 - e^{-i\pi/\mu}}$, which has two poles (at $\pm e^{-i\pi/2\mu}$) and a branch cut at $z = 0$ which we take along the negative real axis. The decay at infinity and logarithmic growth at the origin enables us to use Cauchy's theorem to move the integral to an integral along the ray $\theta = (\pi/2 - \pi/2\mu)$; since $\mu > 1/2$, there are no singularities in the intervening region. Along this ray $z = i\rho e^{-i\pi/2\mu}$, so we have

$$\operatorname{Re}(J(\mu)) = \mu \int_0^\infty (\log \rho + i\frac{\pi}{2}(1 - 1/\mu)) \frac{i}{\rho^2 + 1} d\rho. \quad (\text{A.4.24})$$

The imaginary part of this integral is, fortunately, zero, and the real part is $-\pi^2(\mu - 1)/4$, completing the proof. \square

As an immediate consequence of the Lemma, we have that for some function $E(\alpha)$,

$$g_\alpha(-i) = e^{E(\alpha)} e^{i(1-\mu)\pi/4}. \quad (\text{A.4.25})$$

This completes the proof of Theorem A.4.1. \square

A.5 Asymptotics of Peters solution as $z \rightarrow 0$

This analysis is substantially easier. First we have a small proposition:

Proposition A.5.1. $g_\alpha(\zeta) \rightarrow 1$ as $|\zeta| \rightarrow \infty$, and the convergence is uniform in any sector away from the negative real axis.

Proof. This is claimed on p. 329 of [Pet50] and is immediate from (A.2.3) when $\operatorname{Re}(\zeta) > 0$, but is not so clear for other values of $\arg(\zeta)$, so we rewrite the proof here.

From the definition of $g_\alpha(\zeta)$ it suffices to show that $I_\alpha(\zeta) \rightarrow 0$ as $|\zeta| \rightarrow \infty$. This is immediately clear when $\arg(\zeta) \in (-\alpha, \alpha)$, from the definition of $I_\alpha(\zeta)$ and a change of variables. For other values of $\arg(\zeta)$ we can use other representations of $I_\alpha(\zeta)$. For example, the representation (A.4.15) is good beyond $\zeta = \pm\pi/2$, and each term approaches zero as $|\zeta| \rightarrow \infty$ as long as $\zeta e^{-i\alpha}$ is not on M_2 or L_0 . The rate of convergence depends only on the distance of $\zeta/|\zeta|$ from M_2 and L_0 . Continuing to move the contours as in the appendix of [Pet50], we can show that $I_\alpha(\zeta)$ goes to zero as $|\zeta| \rightarrow \infty$ as long as ζ is not on the branch cut. This completes the proof. \square

Now recall that

$$f(z) = i \int_P \frac{g_\alpha(\zeta)}{\zeta + i} e^{z\zeta} d\zeta.$$

Assume $|z| < 1$. Change variables to $w = z\zeta$, then deform the contour back to $\arg(z)P$ (this works since all singularities of the integrand are inside P). We get

$$f(z) = i \int_{\arg(z)P} \frac{g_\alpha(w/z)}{w + iz} e^w dw.$$

As $|z| \rightarrow 0$, $g_\alpha(w/z)$ converges uniformly to 1, and $(w + iz)^{-1}$ converges uniformly to w^{-1} . So as $|z| \rightarrow 0$,

$$f(z) \rightarrow i \int_{\arg(z)P} w^{-1} e^w dw = -2\pi.$$

Thus we have established that $\lim_{z \rightarrow 0} f(z)$ exists and equals -2π . This completes the proof of Theorem 2.1.4 in the Robin-Neumann case.

A.6 Robin-Dirichlet problem

This is also in [Pet50], again with some details but not others. We again look for a function $f(z)$ analytic in \mathfrak{S}_α . The condition at $z \in \mathfrak{S}_\alpha \cap \{\theta = -\alpha\}$ is now $\operatorname{Re}(f(z)) = 0$. Following Peters, using the same notation for $g_\alpha(\zeta)$ as before, a solution is

$$f(z) = i \int_P \frac{g_\alpha(\zeta)}{\zeta + i} \zeta^{-\mu} e^{z\zeta} d\zeta, \quad k \in \mathbb{Z}. \quad (\text{A.6.1})$$

The only difference between the solutions is the multiplication by $\zeta^{-\mu}$ inside the integral.

We verify that (A.6.1) satisfies the properties we need. Indeed it is obviously analytic in \mathfrak{S}_α (at least away from the corner point). The boundary condition on the real axis is verified in precisely the same way as for the Robin-Neumann problem, since $\zeta^{-\mu}$ is real when ζ is real. For the Dirichlet boundary condition, we again change variables to $w = \zeta e^{-i\alpha}$, when (A.6.1) becomes

$$f(z) = \int_{P_1} \frac{w^{-\mu} e^{i\alpha} g_\alpha(w e^{i\alpha})}{w e^{i\alpha} + i} e^{zw} dw = \int_{P_1} w^{-\mu-1} \exp\{-I_\alpha(w)\} e^{\rho w} dw, \quad (\text{A.6.2})$$

where P_1 is symmetric with respect to the real axis and the branch cut is along the negative real axis. As for (A.3.3), this has zero real part, showing that (A.6.2) is a solution.

Now we analyze the asymptotics. Let

$$\chi_{D,\alpha} = \chi + \frac{\pi^2}{4\alpha} = \frac{\pi}{4} \left(1 + \frac{\pi}{2\alpha} \right).$$

Analogously to the Robin-Neumann case, we have:

Theorem A.6.3. *With $E(\alpha) : (0, \pi/2) \rightarrow \mathbb{R}$ as in Theorem A.4.1, there exists $R'_\alpha(z) : W \times (0, \pi/2) \rightarrow \mathbb{C}$ such that*

$$f(z) = e^{E(\alpha)} e^{-i(z-\chi_{D,\alpha})} + R'_\alpha(z), \quad (\text{A.6.4})$$

where for any fixed $\alpha \in (0, \pi/2)$, there exists a constant $C > 0$ such that for all $z \in \mathfrak{S}_\alpha$,

$$|\operatorname{Re}(R'_\alpha(z))| \leq C|z|^{-2\mu}, \quad |\nabla_z(\operatorname{Re}(R'_\alpha(z)))| \leq C|z|^{-2\mu-1}. \quad (\text{A.6.5})$$

Proof. We mimic the proof of Theorem A.4.1. By the same contour-deformation process, we have

$$\begin{aligned} f(z) &= 2\mu^{1/2}g_\alpha(-i)e^{-iz}e^{i\pi\mu/2} + \frac{\mu^{1/2}}{i\pi} \int_{P'} \frac{g_\alpha(\zeta)\zeta^{-\mu}}{\zeta+i} e^{z\zeta} d\zeta \\ &+ \sum_{j=1}^m e^{z\lambda_j} 2\mu^{1/2} \operatorname{Res}_{\zeta=\lambda_j} (g_\alpha(\zeta)\zeta^{-\mu}/(\zeta+i)). \end{aligned} \quad (\text{A.6.6})$$

From the evaluation of $g_\alpha(-i)$ in the proof of Theorem A.4.1, we immediately see that the first term is $e^{E(\alpha)}e^{-i(z-\chi_{D,\alpha})}$. The sum of residues in (A.6.6) has the needed decay estimates. And if we call the second term $\tilde{R}'_\alpha(z)$, the methods of the proof of Theorem A.4.1 give

$$\tilde{R}'_\alpha(z) = -\frac{\mu^{1/2}}{\pi} \int_{P'} e^{z\zeta} e^{-h(\zeta)} \frac{d\zeta}{\zeta}, \quad (\text{A.6.7})$$

which has an extra factor of $\zeta^{-\mu}$ compared with (A.4.12). Changing variables as before gives

$$\tilde{R}'_\alpha(z) = -\frac{\mu^{1/2}}{\pi} \int_{\arg(z)P'} e^w e^{-h(w/z)} \frac{dw}{w}. \quad (\text{A.6.8})$$

Plugging in $h(\cdot) = p(\cdot) + R(\cdot)$ gives

$$\tilde{R}'_\alpha(z) = -\frac{\mu^{1/2}}{\pi} \int_{\arg(z)P'} e^w e^{-p(w/z)} e^{-R(w/z)} \frac{dw}{w}. \quad (\text{A.6.9})$$

Using the estimates on $R(w/z)$, we have

$$\tilde{R}'_\alpha(z) = -\frac{\mu^{1/2}}{\pi} \int_{\arg(z)P'} e^w e^{-p(w/z)} (1 + O(|w/z|^{2\mu})) \frac{dw}{w}. \quad (\text{A.6.10})$$

For the part of the integral corresponding to the first term 1, $p(w/z)$ is holomorphic in the full disk with no branch cut, so the contour may be deformed to a small disk around $w = 0$ (plus a portion around any branch cut along the negative real axis starting from $w = -1$), then evaluated using residues. We see that the first term is equal to

$$-2i\mu^{1/2} + O(e^{-|z|/C}), \quad (\text{A.6.11})$$

and therefore its real part is $O(e^{-|z|/C})$, even better than needed. On the other hand, the exact same analysis as in the Robin-Neumann case shows that the second term is $O(|z|^{-2\mu})$, which is what we wanted.

The gradient estimate follows as before, since differentiating (A.6.7) just brings down an extra power of ζ , which becomes w/z . □

The analysis of $f(z)$ as $z \rightarrow 0$ follows as before. We can show that

$$f(z) \sim z^\mu \int_{\arg(z)P} w^{-\mu-1} e^w dw.$$

Since $\mu > 0$, we have $\lim_{z \rightarrow 0} f(z) = 0$, and moreover $f(z) = O(|z|^\mu)$, which is what we need to complete the proof of Theorem 2.1.4 in the Robin-Dirichlet case.

Remark A.6.12. Using a similar approach, one can show that both in the Robin-Neumann and the Robin-Dirichlet cases, the solution $f(z)$ is smooth down to $z = 0$, and for each $k > 0$, $\partial_z^k f(z) = O(|z|^\mu)$ as $z \rightarrow 0$.

B Proof of Proposition 4.2.4.

B.1 Plan of the argument

Here we prove Proposition 4.2.4 by using the analysis of [Nai67]. As we will see, it is a special case of Theorem 2 in [Nai67].

Remark B.1.1. The argument presented in [Nai67] is slightly different for q even and q odd. In what follows we assume that q is even, which is in fact sufficient for our purposes. The proof for odd q is analogous and is left to the interested reader. Throughout we let $n = 2q$ and note that $n = 0 \pmod{4}$.

We first claim the following technical lemma:

Lemma B.1.2. *Suppose $q \in \mathbb{N}$ is even. Then the boundary conditions in (3.2.2) and (4.2.1) are regular in the sense of [Nai67, section 4.8]. Moreover, using the notation of [Nai67, equations (41)–(42)], $\theta_0 = 0$, and $\theta_1 = \theta_{-1} \neq 0$ in both cases.*

Assuming this lemma, we obtain asymptotics for the eigenvalues. Indeed, let ξ' and ξ'' be the roots of the equation $\theta_1 \xi^2 + \theta_0 \xi + \theta_{-1} = 0$; by our calculation, these are the roots of $\xi^2 + 1 = 0$:

$$\xi' = -i, \xi'' = i.$$

From [Nai67, Theorem 2, equations (45a) and (45b)], using the fact that $n = 0 \pmod{4}$, all sufficiently large eigenvalues form two sequences:

$$\begin{aligned} (\lambda'_k)^{2q} &= (2k\pi)^{2q} \left(1 - \frac{q \ln_0 \xi'}{k\pi i} + O\left(\frac{1}{k^2}\right) \right); \\ (\lambda''_k)^{2q} &= (2k\pi)^{2q} \left(1 - \frac{q \ln_0 \xi''}{k\pi i} + O\left(\frac{1}{k^2}\right) \right). \end{aligned}$$

Taking $2q$ -th roots and using Taylor series, and plugging in ξ' and ξ'' :

$$\begin{aligned} \lambda'_k &= 2k\pi \left(1 - \frac{\ln_0 \xi'}{2k\pi i} + O\left(\frac{1}{k^2}\right) \right) = 2k\pi + \frac{\pi}{2} + O\left(\frac{1}{k}\right); \\ \lambda''_k &= 2k\pi \left(1 - \frac{\ln_0 \xi''}{2k\pi i} + O\left(\frac{1}{k^2}\right) \right) = 2k\pi - \frac{\pi}{2} + O\left(\frac{1}{k}\right). \end{aligned}$$

Combining the two sequences yields Proposition 4.2.4.

Remark B.1.3. Theorem 2 in [Nai67], in some editions and English translations, only states that there are sequences of eigenvalues of the desired form, rather than stating that *all* sufficiently large eigenvalues form those sequences. However, the original Russian edition claims the stronger version, and it follows immediately from the proof in [Nai67] anyway.

B.2 Proof of Lemma B.1.2

To find θ_1 , θ_0 , and θ_{-1} , we need to analyze the notation of [Nai67]. The boundary conditions will be seen to be regular conditions of Sturm type, hence falling under the analysis of [Nai67, section 4.8], in particular giving $\theta = 0 = 0$ and θ_1 and θ_{-1} defined by (41)–(42) in [Nai67]. To analyze these determinants we need to identify the numbers ω_j , k_j , and k'_j . We do this first.

To analyze the ω_j , see the discussion before [Nai67, (38)]. Note that the sector S_0 referred to in the statement of theorem 2 is the sector of the complex plane with argument between 0 and $\frac{\pi}{n}$ (see [Nai67, (3)]). We then let $\omega_1, \dots, \omega_n$ be the n th roots of -1 , arranged so that for $\rho \in S_0$,

$$\operatorname{Re}(\rho\omega_1) \leq \operatorname{Re}(\rho\omega_2) \leq \dots \leq \operatorname{Re}(\rho\omega_n).$$

To figure out this ordering, let

$$\omega := e^{i\pi/n}.$$

Then, in order, the set $\{\omega_1, \dots, \omega_n\}$ is

$$\{\omega^{2q-1}, \omega^{-(2q-1)}, \dots, \omega^3, \omega^{-3}, \omega^1, \omega^{-1}\}.$$

Note further that $\omega_\mu = \omega_q$; since q is even,

$$\omega_\mu = \omega^{-(q+1)}; \quad \omega_{\mu+1} = \omega^{q-1}.$$

In particular we actually have $\omega_{\mu+1} = -\omega_\mu$. This will be useful.

Now we need to understand the k_j and k'_j , which is done via a direct comparison with [Nai67, (40)]. The sums all vanish, and only the first terms remain. We see that we always have $k'_j = k_j$ in both the Neumann and Dirichlet settings, and for Neumann we have

$$k_1 = 2q - 1, \quad k_2 = 2q - 2, \dots, \quad k_q = q.$$

For Dirichlet we have

$$k_1 = q - 1, \quad k_2 = q - 2, \dots, \quad k_q = 0.$$

So these conditions are of Sturm type.

Since the conditions are Sturm type we immediately have $\theta_0 = 0$ as well as the formulas [Nai67, (41)–(42)] for θ_{-1} and θ_1 . Assume for the moment we are working in the Neumann setting. Define the matrices

$$A_q := \begin{bmatrix} \omega_1^{k_1} & \dots & \omega_\mu^{k_1} \\ \vdots & \ddots & \vdots \\ \omega_1^{k_\mu} & \dots & \omega_\mu^{k_\mu} \end{bmatrix} = \begin{bmatrix} \omega^{(2q-1)(2q-1)} & \omega^{-(2q-1)(2q-1)} & \dots & \omega^{-(q+1)(2q-1)} \\ \vdots & \vdots & \dots & \vdots \\ \omega^{(2q-1)(q)} & \omega^{-(2q-1)(q)} & \dots & \omega^{-(q+1)(q)} \end{bmatrix},$$

$$B_q := \begin{bmatrix} \omega_{\mu+1}^{k_1} & \dots & \omega_n^{k_1} \\ \vdots & \ddots & \vdots \\ \omega_{\mu+1}^{k_\mu} & \dots & \omega_n^{k_\mu} \end{bmatrix} = \begin{bmatrix} \omega^{(q-1)(2q-1)} & \omega^{-(q-1)(2q-1)} & \dots & \omega^{-1(2q-1)} \\ \vdots & \vdots & \dots & \vdots \\ \omega^{(q-1)(q)} & \omega^{-(q-1)(q)} & \dots & \omega^{-1(q)} \end{bmatrix}.$$

Let A'_q be A_q with the last column replaced by the first column of B_q , and let B'_q be B_q with the first column replaced by the last column of A_q . Note also that since $\omega_{\mu+1} = -\omega_\mu$, those

switched columns are the same except that the entry in each odd row is multiplied by -1 . In particular their last entries are the same, since q is even. Then [Nai67, (41)–(42)] give

$$\theta_{-1} = \pm \det A_q \det B_q; \quad \theta_1 = \pm \det A'_q \det B'_q, \quad (\text{B.2.1})$$

where the signs are the same (they both come from the same equation before (41) and (42) in [Nai67]). Thus it suffices to show that

$$\frac{\det A'_q}{\det A_q} = \frac{\det B'_q}{\det B_q}. \quad (\text{B.2.2})$$

The trick will be to reduce these to Vandermonde determinants by taking a factor out of each column; the factor we take out will always be the *last* entry in each column, which is simplified by the fact that $w^q = i$. For example,

$$\det B_q = i^{(q-1)} i^{-(q-1)} i^{(q-3)} i^{-(q-3)} \dots i^1 i^{-1} \det D_q = \det D_q,$$

where

$$D_q = \begin{bmatrix} \omega^{(q-1)(q-1)} & \omega^{-(q-1)(q-1)} & \dots & \omega^{-1(q-1)} \\ \vdots & \vdots & \dots & \vdots \\ \omega^{(q-1)(1)} & \omega^{-(q-1)(1)} & \dots & \omega^{-1(1)} \\ 1 & 1 & \dots & 1 \end{bmatrix}.$$

Similarly, $\det A_q = \det C_q$, where C_q is obtained as with D_q . As for B'_q and A'_q , note that the last entry of the last column of A_q is the same as the last entry of the first column of B_q , so all the pre-factors still cancel and we have $\det A'_q = \det C'_q$, $\det B'_q = \det D'_q$, with D'_q and C'_q obtained by switching the first column of D_q for the last column of C_q .

Switching the order of all of the rows introduces a factor of $(-1)^{q/2}$ in each determinant and brings us to a Vandermonde matrix in each case:

$$\det B_q = (-1)^{q/2} \det V(\omega^{q-1}, \omega^{-(q-1)}, \dots, \omega^1, \omega^{-1});$$

$$\det B'_q = (-1)^{q/2} \det V(\omega^{-(q+1)}, \omega^{-(q-1)}, \dots, \omega^1, \omega^{-1});$$

$$\det A_q = (-1)^{q/2} \det V(\omega^{2q-1}, \omega^{-(2q-1)}, \dots, \omega^{q+1}, \omega^{-(q+1)});$$

$$\det A'_q = (-1)^{q/2} \det V(\omega^{2q-1}, \omega^{-(2q-1)}, \dots, \omega^{q+1}, \omega^{q-1}).$$

Each of these Vandermonde determinants can be computed explicitly via the well-known pairwise difference formula. Considering $\det B'_q / \det B_q$, all the pairwise differences cancel except for those involving the first element. The first element just changes by an overall sign from B_q to B'_q , and we get

$$\frac{\det B'_q}{\det B_q} = \frac{(\omega^{-1} + \omega^{q-1})(\omega^1 + \omega^{q-1}) \dots (\omega^{-(q-1)} + \omega^{q-1})}{(\omega^{-1} - \omega^{q-1})(\omega^1 - \omega^{q-1}) \dots (\omega^{-(q-1)} - \omega^{q-1})}.$$

Similarly,

$$\frac{\det A'_q}{\det A_q} = \frac{(-\omega^{q-1} - \omega^{q+1}) \dots (-\omega^{q-1} - \omega^{-(2q-1)})(-\omega^{q-1} - \omega^{2q-1})}{(\omega^{q-1} - \omega^{q+1}) \dots (\omega^{q-1} - \omega^{-(2q-1)})(\omega^{q-1} - \omega^{2q-1})}.$$

Pulling a minus sign out of each term on the top and the bottom and reversing the order of the multiplication gives

$$\frac{\det A'_q}{\det A_q} = \frac{(\omega^{q-1} + \omega^{2q-1})(\omega^{q-1} + \omega^{-(2q-1)}) \dots (\omega^{q-1} + \omega^{q+1})}{(-\omega^{q-1} + \omega^{2q-1})(-\omega^{q-1} + \omega^{-(2q-1)}) \dots (-\omega^{q-1} + \omega^{q+1})}.$$

Flipping the order of addition within each term:

$$\frac{\det A'_q}{\det A_q} = \frac{(\omega^{2q-1} + \omega^{q-1})(\omega^{-(2q-1)} + \omega^{q-1}) \dots (\omega^{q+1} + \omega^{q-1})}{(\omega^{2q-1} - \omega^{q-1})(\omega^{-(2q-1)} - \omega^{q-1}) \dots (\omega^{q+1} - \omega^{q-1})}.$$

Using the fact that $\omega^{2q} = 1$ to simplify one element in each term, we see that this fraction is precisely $\frac{\det B'_q}{\det B_q}$, completing the proof of Lemma B.1.2 for the Neumann case.

The Dirichlet case is very similar, in fact easier; we have k_1 through k_q each decreasing by q . In particular the last entries of each column are 1, so the matrices corresponding to A_q, B_q, A'_q, B'_q are already Vandermonde. In fact, they are exactly equal to $C_q, D_q, C'_q,$ and D'_q respectively, so our previous computation completes the proof in this case as well. \square

C Numerical examples

In this Appendix we present some numerical results. All computations have been performed using the package FreeFem++ [Hec12].

C.1 Example illustrating Theorems 1.2.2 and 1.3.2

Let Ω be a triangle with $L = 1$, $\alpha = \frac{2\pi}{5}$ and $\beta = \frac{\pi}{6}$. We consider sloshing with Neumann or Dirichlet conditions on Ω . The eigenvalues and quasi-frequencies are given in the table below, and we see that the error is indeed quite small:




Figure 5: Triangle from Example C.1

k	λ_k	σ_k	$\left \frac{\sigma_k}{\lambda_k} - 1 \right $	λ_k^D	σ_k^D	$\left \frac{\sigma_k^D}{\lambda_k^D} - 1 \right $
1	0.	-0.88357		2.43592	2.45437	7.58×10^{-3}
2	0.85626	0.68722	1.97×10^{-1}	4.02389	4.02517	3.17×10^{-4}
3	2.28840	2.2580	1.33×10^{-2}	5.59623	5.59596	4.83×10^{-5}
4	3.82292	3.8288	1.54×10^{-3}	7.16681	7.16676	7.11×10^{-6}
5	5.39779	5.3996	3.37×10^{-4}	8.73757	8.73755	1.81×10^{-6}
6	6.96977	6.9704	9.22×10^{-5}	10.3084	10.3084	1.44×10^{-6}
7	8.54086	8.5412	4.00×10^{-5}	11.8792	11.8791	1.74×10^{-6}
8	10.1118	10.112	2.03×10^{-5}	13.4500	13.4499	2.36×10^{-6}
9	11.6827	11.683	1.11×10^{-5}	15.0208	15.0207	3.26×10^{-6}
10	13.2535	13.254	5.90×10^{-6}	16.5916	16.5915	4.47×10^{-6}

C.2 Example illustrating Remark 1.6.7

We define two domains Ω_{\pm} by setting a *curved* sloshing surface

$$S_{\pm} = \left\{ \left(x, \pm \frac{1}{2\pi} \sin(2\pi x) \right) \mid 0 < x < 1 \right\},$$

so that the length of the sloshing surface is

$$L = \int_0^1 \sqrt{1 + \cos^2(2\pi x)} dx = \frac{2\sqrt{2}}{\pi} E\left(\frac{1}{2}\right) \approx 1.21601$$

(where E denotes a complete elliptic integral of the second kind),

$$\mathcal{W}_1 = \mathcal{W}_{1,\pm} = \{z : |z - 1/2| = 1/2, -\pi \leq \arg(z) \leq -\pi/2\},$$

$$\mathcal{W}_2 = \mathcal{W}_{2,\pm} = \{z : |z - 1/2| = 1/2, -\pi/2 \leq \arg(z) \leq 0\},$$

so that

$$\alpha_+ = \beta_- = \frac{3\pi}{4}, \quad \alpha_- = \beta_+ = \frac{\pi}{4},$$

and impose in both cases the Dirichlet boundary condition on \mathcal{W}_1 and the Neumann one on \mathcal{W}_2 .

The quasifrequencies are then given by (1.6.5), which after omitting O -terms simplifies to

$$\sigma_{+,k} = \frac{\pi}{L} \left(k - \frac{1}{6} \right), \quad \sigma_{-,k} = \frac{\pi}{L} \left(k - \frac{5}{6} \right).$$

The comparisons between the numerically calculated eigenvalues and the quasifrequencies are




Figure 6: Domains Ω_+ (left) and Ω_- (right)

below. Note that there is very good agreement even for low k .

k	$\lambda_{+,k}$	$\sigma_{+,k}$	$\left \frac{\sigma_{+,k}}{\lambda_{+,k}} - 1 \right $	$\lambda_{-,k}$	$\sigma_{-,k}$	$\left \frac{\sigma_{-,k}}{\lambda_{-,k}} - 1 \right $
1	1.02371	2.15294	1.10	1.24543	0.430589	6.54×10^{-1}
2	5.65749	4.73648	1.63×10^{-1}	2.63524	3.01412	1.44×10^{-1}
3	8.13194	7.32001	9.98×10^{-2}	5.55627	5.59765	7.45×10^{-3}
4	10.3085	9.90354	3.93×10^{-2}	8.22122	8.18119	4.87×10^{-3}
5	12.8138	12.4871	2.55×10^{-2}	10.6845	10.7647	7.51×10^{-3}
6	15.3856	15.0706	2.05×10^{-2}	13.1122	13.3483	1.80×10^{-2}
7	17.9151	17.6541	1.46×10^{-2}	15.7600	15.9318	1.09×10^{-2}
8	20.4310	20.2377	9.46×10^{-3}	18.4111	18.5153	5.66×10^{-3}
9	22.9800	22.8212	6.91×10^{-3}	21.0011	21.0988	4.66×10^{-3}
10	25.5511	25.4047	5.73×10^{-3}	23.5873	23.6824	4.03×10^{-3}

D List of recurring notation

Notation	Meaning	Section
Ω	Sloshing domain	1
S, \mathcal{W}	Sloshing surface, free boundary	1
A, B	Vertices of Ω	1
L	Length of sloshing surface S	1
α, β	Interior angles of sloshing domain at A, B	1
$0 = \lambda_1 < \lambda_2 \dots$	Sloshing eigenvalues	1
$\{u_k\}$	Sloshing eigenfunctions	1
\mathcal{D}	Sloshing operator	1

\mathcal{H}	Harmonic extension operator from S to Ω with given boundary conditions on \mathcal{W}	1
$\{\lambda_k^D, u_k^D\}$	Eigenvalues and eigenfunctions for Steklov-Dirichlet problem	1
q	A positive integer, used often when $\alpha = \pi/2q$	1
$\{\Lambda_k, U_k\}$	Eigenvalues and eigenfunctions for associated ODE	1
$T = ABZ$	Triangle, used when Ω is triangular	1
$(x, y), (\rho, \theta)$	Cartesian and polar coordinates for \mathbb{R}^2	2
z	Associated complex coordinate $z = x + iy$	2
\mathfrak{S}_α	Planar sector with $-\alpha \leq \theta \leq 0$	2
$\chi_{\alpha, N}$ etc.	Phase shifts for various angles and boundary conditions	2
μ_α (μ for short)	Equals $\pi/(2\alpha)$	2
$\phi_{\alpha, N}$ etc.	Peters solutions in a planar sector	2
$\{\sigma_j\}, \sigma$	Quasi-frequencies	2
$p_\sigma(x, y)$	Plane wave corresponding to scaled Peters solution	2
v'_σ	Uncorrected quasimode (prime is NOT derivative)	2
ψ	Fixed function in $C_0^\infty(S)$	2
η_σ	Neumann correction function	2
v_σ, v_{σ_j}	Corrected quasimode, j th corrected quasimode	2
$\{v_j\}$	Restrictions of $\{v_{\sigma_j}\}$ to S	2
$\{\varphi_k\}$	Orthonormal basis of eigenfunctions of \mathcal{D} ; equal to $\{u_k _S\}$	2
ξ_q (ξ for short)	Equals $e^{-i\pi/q}$	3
$g(z)$	Arbitrary complex-valued function	3
\mathcal{A}, \mathcal{B}	Operators on complex-valued functions	3
SD	Steklov defect of a function	3
$f(z),$ variants $f_j(z)$	$f(z) = e^{-iz}$, variants obtained by applying \mathcal{A} and \mathcal{B} repeatedly	3
$f(z),$ variants	Analogues of $f(z)$ for Dirichlet case	3
$\eta(\xi)$	Equals $(\xi + 1)/(\xi - 1)$	3
$v_\alpha(z)$	Hanson-Lewy solution for angle α	3
$\gamma(\xi)$	Product involving $\eta(\xi)$	3
v_β^d	Sum of decaying exponents in v_β	3
$g_\sigma(z)$	Quasimodes on triangles constructed with Hanson-Lewy solutions	3
Boldface	Denotes abstract linear algebra context	4
λ_j, σ_j	Eigenvalues and quasi-eigenvalues in linear algebra lemma	4

$F(j)$	Error in linear algebra lemma	4
$\{w_j\}$	Linear combinations of sloshing eigenfunctions on S	4
$\{W_j\}$	Linear combinations of ODE eigenfunctions	4
n	$2q$	4
$\{\omega_k\}$	n th roots of -1	4
J_q	Integer representing asymptotic shift for angles $\pi/2q$	4
$\chi(z)$	Cutoff function	5
Tildes	Denotes curvilinear domain setting	5
\tilde{v}'_σ	Uncorrected quasimode, curvilinear case	5
$\tilde{\eta}_\sigma$	Neumann correction, curvilinear case	5
\tilde{v}_σ	Corrected quasimode (but not yet harmonic), curvilinear case	5
\bar{v}_σ	Further-corrected quasimode, curvilinear case (now harmonic)	5
$\tilde{\varphi}_\sigma$	Solution to a mixed boundary value problem on Ω	5
Φ	Conformal map from Ω to T	5
φ_σ	Solution to corresponding mixed problem on T	5
$f(z)$	Complex-valued Peters solution	A
ζ	New complex variable	A
$I_\alpha(\zeta)$, or $I(\zeta)$	Auxiliary function for Peters solution	A
$g_\alpha(\zeta)$	Exponentiated and modified version of $I(\zeta)$	A
P, P_1, P' , etc.	Complex contours in ζ plane	A
$R_\alpha(z)$	Remainder term: difference of Peters solution and plane wave	A
$\lambda_1, \dots, \lambda_m$	Residues of g_α	A
$\tilde{R}_\alpha(z)$	Remainder term without the residues	A
ω	Equals $e^{i\pi/n}$	B
$\theta_0, \theta_1, \theta_{-1}$	Notation from Naimark	B

References

- [Agr06] M.S. Agranovich, *On a mixed Poincare-Steklov type spectral problem in a Lipschitz domain*, Russ. J. Math. Phys. **13**(3), (2006), 281–290.
- [Alk77] G. Alker, *The scattering of short waves by a cylinder*, J. Fluid Mech. **82** (1977), part 4, 673–686.

- [BKPS10] R. Bañuelos, T. Kulczycki, I. Polterovich, and B. Siudeja, *Eigenvalue inequalities for mixed Steklov problems*. In Operator theory and its applications, volume **231** of Amer. Math. Soc. Transl. Ser. 2, pages 19-34. Amer. Math. Soc., Providence, RI, 2010.
- [Bro94] R. Brown, *The mixed problem for Laplace's equation in a class of Lipschitz domains*, Comm. PDE. (1994), **19** (1994).
- [Buh17] L. Buhovski, *Private communication*, 2017.
- [Dav65] A. M. J. Davis, *Two-dimensional oscillations in a canal of arbitrary cross-section*, Proc. Cam. Phil. Soc, **61** (1965), 827–846.
- [Dav74] A.M.J. Davis, *Short surface waves in a canal; dependence of frequency on the curvatures and their derivatives*, Q. J. Mech. Appl. Math. **27** (1974), 523–535.
- [DW00] A.M.J. Davis and P.D. Weidman, *Asymptotic estimates for two-dimensional sloshing modes*, Phys. Fluids **12** (2000), no. 5, 971–978.
- [FK83] D. W. Fox and J. R. Kuttler, *Sloshing frequencies*, Zeitschrift für angewandte Mathematik und Physik, **34** (1983), no. 5, 668–696.
- [Fi16] D. Fischer, <https://math.stackexchange.com/users/83702/daniel-fischer>, Proving that an orthonormal system close to a basis is also a basis, URL (version: 2016-09-29): <https://math.stackexchange.com/q/1365967>.
- [GP17] A. Girouard and I. Polterovich, *Spectral geometry of the Steklov problem*, J. Spectral Theory **7** (2017), no. 2, 321–359.
- [GP17a] A. Girouard and I. Polterovich, *The Steklov eigenvalue problem: some open questions*, CMS Notes **48** (2016), no. 3, 16–17.
- [Gre87] A. G. Greenhill, *Wave motion in hydrodynamics*, Amer. J. Math. **9** (1887), 62–112.
- [Gri85] P. Grisvard, *Elliptic problems in nonsmooth domains*. London, 1985.
- [Han26] E.T. Hanson, *The theory of ship waves*, Proc. Roy. Soc. London. Ser. A **111** (1926), 491–529.
- [Hec12] F. Hecht, *New development in FreeFem++*, J. Numer. Math. **20** (2012), no. 3-4, 251–265.
- [Kas16] R. Kashaev, *Private communication*, 2016.
- [KMS93] R. Kashaev, V. Mangazeev and Yu. Stroganov, *Star-square and tetrahedron equations in the Baxter-Bazhanov model*, Int. J. Mod. Phys. A **8** (1993), 1399–1409.
- [Kaz71] J. Kazdan, *Perturbation of complete orthonormal sets and eigenfunction expansions*, Proc. Amer. Math. Soc. **27** (1971), no. 3, 506-510.
- [Kon67] V. A. Kondrat'ev, *Boundary value problems for elliptic equations in domains with conical or angular points*, Tr. Mosk. Mat. Obs. **16** (1967), MSU, M., 209–292.

- [KKM04] V. Kozlov, N. Kuznetsov, O. Motygin, *On the two-dimensional sloshing problem*, Proc. Roy. Soc. London A. **460** (2004), 2587–2603.
- [Lam79] H. Lamb, *Hydrodynamics*, 1st ed., Cambridge, 1879, 6th ed., 1932. Reprinted by Dover, New York, 1945.
- [LPPS17] M. Levitin, L. Parnowski, I. Polterovich and D. Sher, *Sloshing, Steklov and Corners II: Steklov spectral asymptotics on polygons*, in preparation.
- [Lew46] Hans Lewy, *Water waves on sloping beaches*, Bull. Amer. Math. Soc. **52** (1946), no. 9, 737–775.
- [Nai67] M. A. Naimark, *Linear differential operators*. New York, 1967–1968.
- [PP14] K.-M. Perfekt, M. Putinar, *Spectral bounds for the Neumann-Poincaré operator on planar domains with corners*, J. Analyse Math. **124** (2014), no. 1, 39–57.
- [Pet50] A. S. Peters, *The effect of a floating mat on water waves*, Comm. Pure and Appl. Math., **3** (1950) no. 4, 319–354.
- [Ron15] L. Rondi, *Continuity properties of Neumann-to-Dirichlet maps with respect to the H-convergence of the coefficient matrices*, Inverse Problems **31** (2015), 045002.
- [San55] L. Sandgren, *A Vibration Problem*, Medd. Lunds Univ., Math. Sem., 1–84 (1955).
- [Ur74] F. Ursell, *Short surface waves in a canal: dependence of frequency on curvature*, J. Fluid Mech. **63** (1974), no. 1, 177–181.