

First order analytic difference equations and integrable quantum systems

S. N. M. Ruijsenaars

*Centre for Mathematics and Computer Science, P.O. Box 4079,
1009 AB Amsterdam, The Netherlands*

(Received 3 January 1996; accepted for publication 26 July 1996)

We present a new solution method for a class of first order analytic difference equations. The method yields explicit “minimal” solutions that are essentially unique. Special difference equations give rise to minimal solutions that may be viewed as generalized gamma functions of hyperbolic, trigonometric and elliptic type—Euler’s gamma function being of rational type. We study these generalized gamma functions in considerable detail. The scattering and weight functions (u - and w -functions) associated to various integrable quantum systems can be expressed in terms of our generalized gamma functions. We obtain detailed information on these u - and w -functions, exploiting the difference equations they satisfy.
© 1997 American Institute of Physics. [S0022-2488(97)01701-5]

TABLE OF CONTENTS

I. INTRODUCTION.....	1069
II. GENERAL RESULTS ON ANALYTIC DIFFERENCE EQUATIONS.....	1076
A. Preliminaries.....	1076
B. Fourier Transform Solutions.....	1079
C. Fourier Series Solutions.....	1087
III. GENERALIZED GAMMA FUNCTIONS.....	1091
A. The Hyperbolic Case.....	1091
B. The Elliptic Case.....	1101
C. The Trigonometric Case.....	1107
IV. SCATTERING FUNCTIONS.....	1112
A. The Hyperbolic Case.....	1112
B. The Elliptic Case.....	1121
C. The Trigonometric Case.....	1125
V. WEIGHT FUNCTIONS.....	1128
A. The Hyperbolic Case.....	1128
B. The Elliptic Case.....	1132
C. The Trigonometric Case.....	1135
APPENDICES.....	1137
A. THE GAMMA FUNCTION AND RELATED INTEGRALS.....	1137
B. UNIFORM ESTIMATES.....	1142

I. INTRODUCTION

This paper is concerned both with the general theory of first order analytic difference equations (from now on Δ Es) and with certain special functions that arise as solutions to Δ Es of a quite restricted type. As announced and partly detailed in our survey¹ and lectures,² among these special functions are the weight functions and scattering amplitudes associated with relativistic quantum integrable systems of Calogero-Moser type—which, in turn, for special parameter

choices reduce to functions occurring in various well-known infinite-dimensional integrable systems, such as the sine-Gordon theory, the XYZ chain and the eight-vertex model.

The first part of the paper (Sections II and III) does not involve integrable systems. To describe the scope of the results obtained therein, we start from two quite elementary first order AΔEs, namely,

$$M(w+1) = cM(w), \quad w \in \mathbb{C}, \quad c \in \mathbb{C}^*, \quad (1.1)$$

$$M(w+1) = wM(w), \quad w \in \mathbb{C}. \quad (1.2)$$

Obviously, the first one is solved by the function $\exp(w \ln c)$ and the second one by Euler's gamma function $\Gamma(w)$. These functions can be used as building blocks for solving AΔEs of the form

$$M(w+1) = Q(w)M(w), \quad w \in \mathbb{C}, \quad (1.3)$$

where $Q(w)$ is a rational function of w . Indeed, any function of the form

$$M(w) = e^{aw} \frac{\prod_{j=1}^M \Gamma(w - b_j)}{\prod_{k=1}^N \Gamma(w - c_k)}, \quad a, b_j, c_k \in \mathbb{C}, \quad (1.4)$$

satisfies (1.3) with $Q(w)$ rational, and varying the parameters a, M, N, b_j, c_k , yields all rational functions.

Suppose now that one can find meromorphic solutions to the AΔE (1.3) for $Q(w)$ equal to the Weierstrass σ -function $\sigma(w; \omega, \omega')$ with $\omega, -i\omega' \in (0, \infty)$, and its trigonometric ($-i\omega' = \infty$) and hyperbolic ($\omega = \infty$) degenerations—the sine and sinh-functions. (The additional factor $c \exp(aw^2)$ in the degenerate σ -functions is easily taken into account—one need only include a factor $\exp(P(w))$ with $P(w)$ a third order polynomial.) Then the respective solutions $M_{\text{ell}}(w), M_{\text{trig}}(w)$ and $M_{\text{hyp}}(w)$ can be used as building blocks to solve the AΔE (1.3) with $Q(w)$ any elliptic function with periods $2\omega, 2\omega'$ or its trigonometric and hyperbolic counterparts, resp. Indeed, any elliptic function $Q(w)$ can be written in the form (1.4), with the exponential replaced by a constant and $\Gamma(w)$ by $\sigma(w)$, so a corresponding meromorphic solution $M(w)$ to (1.3) is obtained by taking $\Gamma \rightarrow M_{\text{ell}}$ in (1.4).

Among other things, this paper presents and studies special functions generalizing the gamma function, which can be used as building blocks to solve AΔEs of the three types just described. In one case the pertinent function is not really new—up to a constant and an exponential it amounts to Thomae's q -gamma function.^{3,4} For the other two cases, however, the corresponding generalized gamma functions are new, and turn out to have some quite remarkable properties. The comprehensive study of these functions (to be found in Section III) constitutes one of the principal results of this paper.

In order to sketch the setting from which our generalized gamma functions emerge, we begin by pointing out that even when one restricts attention to functions $Q(w)$ and solutions $M(w)$ that are meromorphic (as we do), there is an enormous ambiguity in the solution. Indeed, assuming $M(w)$ is a solution and $m(w)$ any meromorphic function with period 1, it is obvious that the function $m(w)M(w)$ is a solution as well. The importance of singling out solutions with *special* properties is therefore evident.

In previous literature, the class of AΔEs to be studied—that is, the class of meromorphic functions $Q(w)$ —has been narrowed down by insisting that $Q(w)$ have a special asymptotics for $\text{Re } w \rightarrow \infty$. In particular, Nörlund in his comprehensive monograph⁵ uses this prescribed asymptotics to construct the uniquely determined solution he refers to as the “Hauptlösung” (see also Refs. 6–8).

By contrast, the key requirement on $Q(w)$ and $M(w)$ we impose is a special asymptotics for $|\operatorname{Im} w| \rightarrow \infty$, satisfied in particular for functions $Q(w)$ that are periodic in the imaginary direction. As will transpire below, this leads to essentially the same solutions only for rational and hyperbolic $Q(w)$, whereas Nörlund's methods do not apply to the trigonometric and elliptic cases.

As a matter of fact, we have opted for a shift in the imaginary direction—in contrast to the shift by 1 in the AΔE (1.3). This corresponds to the applications to integrable systems, and is also convenient in view of our different requirements concerning asymptotics. Moreover, we shall treat the step size as a variable, and we do not single out the positive or negative imaginary direction. Thus our starting point is the AΔE

$$F(z + ia/2) = \Phi(z)F(z - ia/2), \quad (1.5)$$

where $\Phi(z)$ is meromorphic, and where the step size a is an arbitrary positive number. Of course, this AΔE is related by a scaling and a shift over half the step size to the AΔE (1.3), so all results can be rephrased for (1.3)—at the expense, however, of cumbersome notation, which moreover hides some symmetries that naturally emerge when the second convention is used.

We are now prepared to describe the organization and results of the paper in more detail. Section II contains our general results on first order AΔEs. In Subsection II A we set the stage by delineating the class of functions $\Phi(z)$ allowed in (1.5). As a first requirement, we insist on $\Phi(z)$ being free of zeros and poles in a strip $|\operatorname{Im} z| < s$, $s > 0$. We denote such AΔEs as regular AΔEs, and refer to solutions that are free of zeros and poles in the strip $|\operatorname{Im} z| < s + a/2$ as regular solutions. The poles and zeros of a regular solution $F(z)$ outside $|\operatorname{Im} z| < s + a/2$ are completely determined by the poles and zeros of $\Phi(z)$ outside $|\operatorname{Im} z| < s$, as easily follows from (1.5).

Regular AΔEs can be rewritten in the additive form

$$f(z + ia/2) - f(z - ia/2) = \phi(z), \quad |\operatorname{Im} z| < s, \quad (1.6)$$

where $\phi(z)$ denotes (a suitable branch of) $\ln \Phi(z)$. Thus the search for regular solutions to (1.5) is reduced to finding solutions $f(z)$ to (1.6) that are analytic for $|\operatorname{Im} z| < s + a/2$. Using well-known properties of the partial differential operator $\partial/\partial \bar{z} = \partial_x + i\partial_y$ and Runge's approximation theorem, it can be proved that such solutions exist. We shall not detail this, however, since the existence arguments yield no information on the solution thus obtained. (An existence proof can be assembled from Ref. 9, for example.)

By contrast, the extra requirements we impose on $\Phi(z)$ (or equivalently $\phi(z)$) enable us to construct *explicit* solutions, with special properties that render them essentially unique. Roughly speaking, we require that $\phi(z)$ have at worst polynomial increase as $|\operatorname{Re} z| \rightarrow \infty$, and construct solutions $f(z)$ with the same property, which are moreover regular (i.e., analytic for $|\operatorname{Im} z| < s + a/2$). We refer to such solutions as *minimal* solutions: both their singularities and their asymptotics for $|\operatorname{Re} z| \rightarrow \infty$ are “best possible”—being enforced by the singularities and asymptotics of $\phi(z)$. Among other things, Theorem II.1 entails the uniqueness up to a constant of minimal solutions to the additive AΔE (1.6)—assuming they exist.

In Subsections II B and II C we study two classes of AΔEs that do admit minimal solutions—as is shown by exhibiting a minimal solution via explicit formulas involving $\phi(x)$, $x \in \mathbb{R}$. The key results are Theorem II.2 and II.5, resp. Theorem II.2 presupposes that $\phi(x)$ is an $L^1(\mathbb{R})$ -function, whose Fourier transform $\hat{\phi}(y)$ is in $L^1(\mathbb{R})$, too, and satisfies $\hat{\phi}(y) = O(y)$ for $y \rightarrow 0$; its corollary Theorem II.3 handles functions that have these properties after taking a certain number of x -derivatives. In Theorem II.5 it is assumed that $\phi(x)$ has period π/r , $r > 0$, and its zeroth Fourier coefficient vanishes; then Theorem II.6 handles functions $\phi(x)$ for which $\phi^{(k)}(x)$, $k \in \mathbb{N}^*$, has these properties.

The arbitrary additive constant in a minimal solution to the AΔE (1.6) can and will be fixed in the Fourier transform setting of Theorem II.2 by requiring that the solution go to 0 for $x \rightarrow \infty$; in the Fourier series setting of Theorem II.5 it is fixed by requiring that the minimal solution

(which is shown to be π/r -periodic) have vanishing zeroth Fourier coefficient. The unique solution $f(a; z)$ thus obtained is given by (2.26) and (2.106), resp. From the identity (2.38) it then follows that $f(a; z)$ satisfies the addition formula (2.28) in both settings.

The solution $f(a; z)$ has another illuminating feature: In both cases it satisfies

$$\lim_{a \downarrow 0} iaf(a; z) = \psi(z), \quad |\operatorname{Im} z| < s, \quad (1.7)$$

where $\psi(z)$ is a primitive of $\phi(z)$. Therefore, $iaf(a; z)$ may be viewed as a “generalized primitive” of $\phi(z)$. It should be noted that this feature is obviously compatible with the AΔE (1.6), but not *a priori* implied by it: In view of the huge multiplier ambiguity already discussed, the pertinent limit typically does not exist for more general solutions.

Theorems II.4 and II.7 are concerned with the $a \downarrow 0$ limit of minimal solutions to the AΔE (1.6) when ϕ is allowed to have a suitable a -dependence. At first sight, the assumptions may appear very restrictive, but they can in fact be verified for the applications occurring in Section III. The limit (1.7) may be viewed as a quite special consequence of these zero step size limit theorems.

In Appendix A we derive various results that involve Euler’s gamma function, not only as a concrete illustration of the theory developed in Subsections II A and II B, but also to prepare the ground for Section III, which is devoted to a study of generalized gamma functions. Below (1.4) we have already delineated the three cases that will be considered in Section III. Since we employ the AΔE (1.5) and not the AΔE (1.3), however, the trigonometric case turns into the hyperbolic case and vice versa. Moreover, the Weierstrass σ -function and its degenerations are traded for close relatives, to which the theory of Section II applies. The resulting minimal solutions (rendered unique in obvious ways) will be dubbed G -functions.

More specifically, Subsection III A deals with the hyperbolic G -function—the unique minimal solution to the AΔE

$$G(z + ia/2) = 2\operatorname{ch}(\pi z/b)G(z - ia/2), \quad b > 0, \quad (1.8)$$

that satisfies $G(0) = 1$ and $|G(x)| = 1$ for real x . Now it is evident that any solution $G(z)$ to (1.8) has the property that the quotient $G(z + ib/2)/G(z - ib/2)$ is an ia -antiperiodic function. It is not at all obvious, though, that a solution exists for which this quotient equals $2\operatorname{ch}(\pi z/a)$. The hyperbolic G -function does have this striking property: It is given by

$$G_{\text{hyp}}(a, b; z) = \exp\left(i \int_0^\infty \frac{dy}{y} \left(\frac{\sin 2yz}{2\operatorname{sh} y \operatorname{sh} by} - \frac{z}{aby} \right)\right), \quad |\operatorname{Im} 2z| < a + b, \quad (1.9)$$

and hence is manifestly symmetric under $a \leftrightarrow b$.

We present our results on the hyperbolic G -function in seven propositions. Proposition III.1 deals with the three elementary AΔEs to which G is a minimal solution, and Prop. III.2 details various automorphy properties. As already noted above, the poles and zeros of a regular solution to (1.5) readily follow from those of $\Phi(z)$; similarly, residues at simple poles can be determined in terms of $\Phi(z)$. This is worked out for G_{hyp} in Prop. III.3. An important dichotomy first emerges here: When a/b is an irrational number, all poles and zeros are simple, whereas for rational a/b this is not the case.

Since $G_{\text{hyp}}(z)$ is a minimal solution, its logarithm is polynomially bounded for $|\operatorname{Re} z| \rightarrow \infty$ and $|\operatorname{Im} z| \leq a/2$. For the case at hand, the precise asymptotics can be explicitly determined by comparison to the case $a = b$. (This case has special features that render it more accessible.) Proposition III.4 presents the details; the restriction on $|\operatorname{Im} z|$ is readily lifted by exploiting the AΔE (1.8).

From the representation (1.9) it is already clear that for fixed z in the strip $|\operatorname{Im} 2z| < a + b$ the G -function is real-analytic on $(0, \infty)$ in the parameters a and b . In Prop. III.5 we prove that G

actually extends to a function that is meromorphic in a, b and z , as long as the quotient b/a stays away from the negative real axis. This readily follows from a representation for the G -function in terms of an infinite product of gamma functions. To control the convergence of this product, some estimates on Laplace transforms assembled in Appendix B are crucial.

The latter estimates are also exploited in proving that a renormalized version of the hyperbolic G -function converges to the gamma function when one takes $a=1$ and $|b| \rightarrow 0$ in any sector $|\operatorname{Arg} b| \leq \chi, \chi \in [0, \pi)$. This is detailed in Prop. III.6. Two more zero step size limits are obtained in Prop. III.7. In the latter context the limit has branch cuts on the imaginary axis that arise from a confluence of zeros and poles.

Before turning to a sketch of Subsection III B, we would like to mention that G_{hyp} is not only the key building block for the hyperbolic scattering and weight functions of Subsections IVA and VA, but also for our recent generalization of Gauss' hypergeometric function ${}_2F_1$. In this context G_{hyp} plays the role of the gamma function in the Barnes representation for ${}_2F_1$ —except that the generalization is far more symmetric. For ${}_2F_1$ the symmetry is broken, since a step size is taken to zero that leads to the two quite different limiting functions of Propositions III.6 and III.7 (cf. Ref. 2, Subsection 6.3, and papers to appear).

In Subsection III B we study the elliptic G -function, which is given by

$$G_{\text{ell}}(r, a, b; z) = \exp \left(i \sum_{n=1}^{\infty} \frac{\sin 2nrz}{2n \operatorname{sh} nr \operatorname{sh} nr b} \right), \quad |\operatorname{Im} 2z| < a + b, \quad (1.10)$$

along the same lines as its hyperbolic counterpart (1.9). It is not obvious, but true that G_{ell} is a minimal solution to an AΔE of the form

$$\frac{G(z + ia/2)}{G(z - ia/2)} = \exp(c_0 + c_1 z + c_2 z^2) \sigma(z + ib/2; \pi/2r, ib/2), \quad (1.11)$$

where σ denotes the Weierstrass σ -function. Thus it can be used as a building block to solve the AΔE (1.5) with $\Phi(z)$ an elliptic function—as already discussed above.

As it turns out, it is quite convenient to trade the σ -function $\sigma(z; \pi/2r, ia/2)$ for a closely related function $s(r, a; z)$ (2.89). The latter function is odd and π/r -antiperiodic in z , and has limits $r^{-1} \sin rz$ and $\pi a^{-1} \operatorname{sh} \pi a^{-1} z$ for $a \uparrow \infty$ and $r \downarrow 0$, cf. (2.90) and (2.92), resp. Similarly, the function arising on the rhs of (1.11) will be denoted $R(r, b; z)$. In view of (1.10) it is given explicitly by

$$R(r, b; z) = \exp \left(- \sum_{n=1}^{\infty} \frac{\cos 2nrz}{n \operatorname{sh} nr b} \right), \quad |\operatorname{Im} 2z| < b, \quad (1.12)$$

so it is even and π/r -periodic in z . Most of the propositions in Subsection III B may be viewed as generalizations of hyperbolic counterparts, since one has

$$\lim_{r \downarrow 0} \exp(\pi^2/6rb) R(r, b; z) = 2 \operatorname{ch}(\pi z/b) \quad (1.13)$$

and

$$\lim_{r \downarrow 0} \exp(\pi^2 z/6irab) G_{\text{ell}}(r, a, b; z) = G_{\text{hyp}}(a, b; z), \quad (1.14)$$

cf. Prop. III.12.

Subsection III C concerns the trigonometric case. Our trigonometric G -function is given by

$$G_{\text{trig}}(r, a; z) = \exp\left(\sum_{n=1}^{\infty} \frac{e^{2inrz}}{2n \operatorname{sh} nra}\right), \quad \operatorname{Im} 2z > -a, \quad (1.15)$$

and can be obtained as a limit of the elliptic G -function, viz.,

$$G_{\text{trig}}(r, a; z) = \lim_{b \uparrow \infty} G_{\text{ell}}(r, a, b; z - ib/2). \quad (1.16)$$

In this case the elementary AΔE satisfied by the G -function reads

$$\frac{G(z + ia/2)}{G(z - ia/2)} = 1 - e^{2irz}. \quad (1.17)$$

Since the rhs has zeros on the real axis, this is not a regular AΔE. However, any shift $z \rightarrow z + ip, p > 0$, yields a regular AΔE, to which the (shifted) G -function is a minimal solution.

Propositions III.14–III.19 concern various properties of the G -function that are quite easily obtained from the series representation (1.15) or the product representation

$$G_{\text{trig}}(r, a; z) = \prod_{m=1}^{\infty} \frac{1}{1 - \exp(2irz - (2m-1)ar)}. \quad (1.18)$$

Proposition III.20, however, involves more work. Here, we prove that a renormalized version of G_{trig} converges to the gamma function for $r \downarrow 0$.

Fixing $a > 0$, it is clear from (1.18) that $G_{\text{trig}}(r, a; z)$ extends to a meromorphic function of r and z , as long as r stays in the right half plane. But one cannot solve the hyperbolic AΔE, obtained from (1.17) upon taking $r \rightarrow i\pi/b, b > 0$, by making use of the trigonometric G -function. By contrast, one is allowed to take $b \rightarrow i\pi/r, r > 0$, in the hyperbolic G -function, yielding the trigonometric function $2\cos rz$ on the rhs of (1.8). Accordingly, the quotient of the renormalized versions of $G_{\text{hyp}}(1, i\pi/r; z)$ and $G_{\text{trig}}(r, 1; z)$ (both of which converge to the gamma function as $r \downarrow 0$) is a quite nontrivial i -periodic function, cf. (3.171)–(3.173).

Just as in Subsections III A and III B, the last proposition of Subsection III C deals with two zero step size limits; once again, a confluence of zeros and poles gives rise to branch cuts. The subsection is concluded by detailing the relation of our trigonometric G -function to the q -gamma function.

We continue by sketching the physical setting in which the scattering and weight functions $u(z)$ and $w(z)$ of Sections IV and V, resp., arise. These functions are associated to relativistically invariant integrable generalizations^{10,11} of the nonrelativistic Calogero-Moser N -particle quantum systems.¹² The dynamics of these relativistic systems belongs to a commutative algebra generated by N independent commuting analytic difference operators. The step size in these difference operators is inversely proportional to the speed of light c , and for $c \rightarrow \infty$ the commuting difference operators converge to commuting differential operators.

Now a factorized product of u -functions is expected to encode the asymptotics of the diagonalizing joint eigenfunction transform, whereas a factorized product of w -functions can be used to transform the difference operators and eigenfunctions to an especially convenient form. In particular, in the trigonometric case the transformed eigenfunctions amount to Macdonald's q -Jacobi multivariable A_{N-1} polynomials, and the product of weight functions yields the function with respect to which the polynomials are orthogonal (cf. Ref. 2, Subsection 6.2 and references given there). (This is why $w(z)$ is referred to as a “weight function.”)

The key point is now that $u(z)$ and $w(z)$ solve first order AΔEs to which the theory developed in Sections II and III applies. In fact, in suitable parameter regimes $u(z)$ can be characterized as the unique minimal solution satisfying $u(0) = 1$ and $|u(x)| = 1$ for real x , whereas a reduced weight function $w_r(z)$ (closely related to $w(z)$) can be characterized in a similar way. It would

take us too far afield to explain here how these AΔEs (which are specified in Sections IV and V) emerge from the difference operators and their eigenfunctions. Instead, we refer to Ref. 1, p. 187, and Ref. 2, Subsection 4.3, for a derivation of the AΔEs satisfied by $u(z)$ and $w(z)$, resp. (See also our forthcoming paper.¹³)

From the viewpoint of special function theory, the u - and w -functions are just simple combinations of the G -functions from Section III: Both functions are of the form $G(\cdots)G(\cdots)/G(\cdots)G(\cdots)$. The pertinent combinations, however, turn out to have quite remarkable properties, which reflect their origins in the context of analytic difference operators and eigenfunction transforms.

We study the functions $u(z)$ and $w(z)$ along similar lines, once more handling the hyperbolic, elliptic and trigonometric cases successively. In each case we first define the relevant function in terms of G -functions, read off some automorphy properties, and introduce some associated functions and/or parameter regimes. Then we study the functions in relation to the elementary AΔEs they obey. As it happens, there is an additional elementary AΔE pertaining to a parameter (essentially the coupling constant in the integrable system picture), which makes it possible to express $u(z)$ and $w(z)$ in terms of products of s -functions (i.e., $\text{sh}(\cdot)$, $s(\cdot)$ and $\sin(\cdot)$, resp.) for certain parameter values. In the hyperbolic and elliptic cases, these values are in fact *dense* in the parameter space.

After obtaining these elementary representations for special parameters, we return to the general case and derive various representations of a different character. At the end of each subsection we obtain a number of limits, whose existence is suggested by the formal limiting behavior of the difference Hamiltonians. Quite a few of these limits may be physically interpreted as nonrelativistic limits. For the scattering functions we also derive limits that may be viewed as classical limits. The zero step size results of Sections II and III are the main tools in controlling most of the limits—in particular the classical limits.

To conclude this introduction, we would like to point out that our results entail a great many nontrivial identities. As a rule, these identities are not spelled out: they follow from different representations for the same function. To be sure, quite a few of these formulas can be assembled via elementary identities—one may even assert that this is precisely what we have done in this paper. But this hindsight wisdom obscures what we view as the basic reason underlying most of the identities, namely, the uniqueness of minimal solutions to first order AΔEs that admit such solutions.

To render the previous paragraph more concrete, we add an example. The sine-Gordon specialization of the u -function from Subsection IV A has been known in terms of the integral (4.30) for almost two decades (cf. Ref. 14 and references given there). Specifically, using our conventions, this S -matrix element reads

$$u(\pi, \alpha, \pi/2; z) = \exp \left(i \int_0^\infty \frac{dy}{y} \frac{\text{sh}(\alpha - \pi/2)y}{\text{ch}(\pi y/2) \text{sh} \alpha y} \sin 2yz \right), \quad |\text{Im} 2z| < d, \quad (1.19)$$

with d given by (4.32). (In point of fact, the integral occurred even earlier as a partition function of the six-vertex model, cf. Ref. 15.) Nevertheless, the result (4.28), expressing (1.19) as an elementary function for the dense set (4.27) of α -values, is new. For $\alpha < \pi$ the resulting identity can be verified directly by noting that the rhs of (4.28) is a minimal solution to the AΔE (4.6) with $\delta = -$, $a_+ = \pi$ and $a_- = \alpha$, which moreover has value 1 and modulus 1 for $z = 0$ and z real, resp., just as (1.19).

II. GENERAL RESULTS ON ANALYTIC DIFFERENCE EQUATIONS

A. Preliminaries

As announced in the Introduction, we are concerned with AΔEs of the form

$$\frac{F(z+ia/2)}{F(z-ia/2)} = \Phi(z), \quad a > 0, \quad (2.1)$$

where $\Phi(z)$ is a function that is meromorphic in \mathbb{C} (briefly: meromorphic). We shall call a function $F(z)$ a *solution* to (2.1) if and only if $F(z)$ is meromorphic in a strip $|\operatorname{Im} z| < s + a/2$, $s \in (0, \infty)$, and $F(z)$ satisfies (2.1) for $|\operatorname{Im} z| < s$.

The first thing to note is that any solution thus defined extends to a meromorphic function. Indeed, one can extend $F(z)$ upwards strip by strip via

$$F(z+ika) \equiv \prod_{j=1}^k \Phi(z+(j-1/2)ia) \cdot F(z), \quad |\operatorname{Im} z| \leq a/2, \quad (2.2)$$

and downwards via

$$F(z-ika) \equiv \prod_{j=1}^k \frac{1}{\Phi(z-(j-1/2)ia)} \cdot F(z), \quad |\operatorname{Im} z| \leq a/2. \quad (2.3)$$

Clearly, the quotient of two solutions to (2.1) is an ia -periodic meromorphic function.

Whenever $\Phi(x+iy)$, $x, y \in \mathbb{R}$, converges to 1 for $y \rightarrow \infty$, uniformly for x varying over arbitrary compact subsets of \mathbb{R} and sufficiently fast, the infinite product

$$F_+(z) \equiv \prod_{j=1}^{\infty} \frac{1}{\Phi(z+(j-1/2)ia)} \quad (2.4)$$

defines a solution to (2.1). We shall refer to F_+ as the upward iteration solution. It is readily seen that it is the only solution satisfying $F(x+iy) \rightarrow 1$ for $y \rightarrow \infty$. Similarly, the downward iteration solution

$$F_-(z) \equiv \prod_{j=1}^{\infty} \Phi(z-(j-1/2)ia) \quad (2.5)$$

exists provided $\Phi(x+iy) \rightarrow 1$ for $y \rightarrow -\infty$ (uniformly on x -compacts and sufficiently fast), and is the unique solution satisfying $F(x+iy) \rightarrow 1$ for $y \rightarrow -\infty$.

Consider, for example, the AΔEs with right-hand sides

$$\Phi_1(z) = \operatorname{ch} z, \quad \Phi_2(z) = 1 - \exp(iz - s), \quad \Phi_3(z) = 1 - \exp(iz + s), \quad s > 0. \quad (2.6)$$

In the first case no iteration solution exists, whereas in the second and third cases F_+ exists, but F_- does not.

Our main interest is in AΔEs (or, equivalently, meromorphic functions $\Phi(z)$) that admit solutions with special properties in the strip $|\operatorname{Im} z| \leq a/2$. Specifically, we shall restrict attention from now on to meromorphic functions $\Phi(z)$ that have no poles and zeros in a strip $|\operatorname{Im} z| < s$. Such functions and the associated AΔEs (2.1) will be called *regular*. A solution to a regular AΔE will be called *regular* iff it has no poles and zeros in $|\operatorname{Im} z| \leq a/2$. In view of (2.2) and (2.3) it then actually has no poles and zeros in $|\operatorname{Im} z| < s + a/2$. Clearly, the quotient of two regular solutions is

an ia -periodic nowhere vanishing entire function. Note that the three AΔEs defined by (2.6) are all regular; in the second case F_+ is regular, whilst in the third case F_+ is not (it has a pole in the set $ia/2[-1, 1]$).

It should be noticed that a regular solution is “maximally analytic,” in the sense that it is free of poles and zeros in the strip $|\operatorname{Im} z| \leq a/2$; its poles and zeros outside the latter strip are then determined by the AΔE (2.1), and can be read off from (2.2) and (2.3), whenever the poles and zeros of $\Phi(z)$ are known. We shall be primarily concerned with a restricted type of AΔE, which admits regular solutions that are “minimal.” To define this notion, we observe that a regular solution $F(z)$ to (2.1) admits a one-valued analytic logarithm in $|\operatorname{Im} z| < s + a/2$. We call F a *minimal* solution iff $\ln F(z)$ is polynomially bounded in $|\operatorname{Im} z| \leq a/2$. That is, there exist $c, d > 0$ and $k \in \mathbb{N}$ such that

$$|\ln F(z)| < c + d|z|^k, \quad \forall z \in \{|\operatorname{Im} z| \leq a/2\}. \quad (2.7)$$

Taking $z = x \in \mathbb{R}$ in (2.1), we deduce

$$|\Phi(x)|^\delta < \exp(2c + 2d|x|^k), \quad \forall x \in \mathbb{R}, \quad \delta = \pm 1. \quad (2.8)$$

Thus, $\Phi(z)$ must satisfy (2.8) for minimal solutions to exist.

To show that AΔEs admitting minimal solutions are by no means exceptional, let $g(z)$ be any meromorphic function that is analytic in $|\operatorname{Im} z| < s + a/2$ and polynomially bounded in $|\operatorname{Im} z| \leq a/2$. Then the AΔE with rhs $\Phi(z) \equiv \exp(g(z + ia/2) - g(z - ia/2))$ admits a minimal solution, viz., $F(z) = \exp(g(z))$. It is also to be noted that the right-hand side functions $\Phi(z)$ of (2.1) that admit minimal solutions form a group: If $F(z)$ is a minimal solution to (2.1), then $1/F(z)$ is a minimal solution to (2.1) with $\Phi \rightarrow 1/\Phi$, and if F_1, F_2 are minimal solutions to AΔEs (2.1) with rhs Φ_1, Φ_2 , resp., then $F(z) = F_1(z)F_2(z)$ is a minimal solution to (2.1) with $\Phi(z) = \Phi_1(z)\Phi_2(z)$.

A minimal solution is not only maximally analytic (since it is regular by definition), but also has the slowest increase to ∞ and/or decrease to 0 for $\operatorname{Re} z \rightarrow \pm \infty$ in the strip $|\operatorname{Im} z| \leq a/2$ that is compatible with (2.1). This will be clear from the following theorem, which shows, moreover, that minimal solutions have “minimal ambiguity.”

Theorem II.1: Assume that the meromorphic function $\Phi(z)$ is regular and satisfies (2.8). Let $F_1(z)$ and $F_2(z)$ be minimal solutions to the AΔE (2.1). Then there exist $C \in \mathbb{C}^*$ and $l \in \mathbb{Z}$ such that

$$F_1(z)/F_2(z) = C \exp(2\pi l z/a). \quad (2.9)$$

If $F_1(z)$ and $F_2(z)$ are bounded away from 0 and ∞ on \mathbb{R} , then one has $l = 0$ in (2.9). If $\Phi(z)$ is even, then for all minimal solutions $F(z)$ the function $F(z)F(-z)$ is constant. If $\Phi(0) = 1$ and the function $\Phi(z)\Phi(-z)$ equals 1, then for any minimal solution $F(z)$ there exists $k \in \mathbb{Z}$ such that $\exp(2\pi k z/a)F(z)$ is an even minimal solution.

Proof: Since F_1 and F_2 are minimal, they are *a fortiori* regular. Therefore, $F_1(z)/F_2(z)$ is an ia -periodic entire function $q(z)$ without zeros. Hence there exists $l \in \mathbb{Z}$ such that the function $q_0(z) \equiv q(z)\exp(-2\pi l z/a)$ has zero winding number around 0 as z goes from z_0 to $z_0 + ia$.

To prove that $q_0(z)$ is constant, we note that it can be written $\exp[r(z)]$, with $r(z)$ an ia -periodic entire function. Since F_1 and F_2 are minimal, $r(z)$ is polynomially bounded:

$$|r(z)| \leq C_1 + C_2|z|^k, \quad |\operatorname{Im} z| \leq a/2. \quad (2.10)$$

It is not hard to see that this entails constancy of $r(z)$. (Indeed, we can, for instance, argue as follows. Since $r(z)$ is ia -periodic and entire, it can be written $\sum_{n \in \mathbb{Z}} c_n w^n \equiv s(w)$, where

$w \equiv \exp(2\pi z/a)$, and where the series converges for $w \in \mathbb{C}^*$. In view of the bound (2.10), the function $ws(w)$ has limit 0 for $w \rightarrow 0$, so it is analytic at $w=0$. Hence, $c_n=0$ for $n<0$. Similarly, since (2.10) entails $s(w)/w \rightarrow 0$ for $w \rightarrow \infty$, we deduce $c_n=0$ for $n>0$.)

We have now proved the first assertion (2.9). The second one is then clear from (2.9). Now assume $\Phi(z)$ is even and $F(z)$ is a minimal solution. Consider the function $G(z) \equiv 1/F(-z)$. It satisfies

$$\frac{G(z+ia/2)}{G(z-ia/2)} = \frac{F(-z+ia/2)}{F(-z-ia/2)} = \Phi(-z) = \Phi(z), \quad (2.11)$$

so it is a solution, too. From minimality of F one easily deduces minimality of G , so (2.9) entails there exists $l \in \mathbb{Z}$ such that $F(z)/G(z) = C \exp(2\pi lz/a)$. But the function on the lhs equals $F(z)F(-z)$ and hence is even. Therefore, we have $l=0$ and the third assertion follows.

To prove the last assertion, consider the function $H(z) \equiv F(-z)$. It satisfies

$$\frac{H(z+ia/2)}{H(z-ia/2)} = \frac{F(-z-ia/2)}{F(-z+ia/2)} = 1/\Phi(-z) = \Phi(z), \quad (2.12)$$

and so it is a second minimal solution. Thus we must have $F(-z) = C \exp(2\pi lz/a) F(z)$. Putting $z=0$ yields $C=1$ and putting $z=ia/2$ yields $(-)^l \Phi(0)=1$, so that l is even. But then $\exp(2\pi kz/a)F(z)$ with $k \equiv l/2$ is an even minimal solution. \square

Thus far, we have been dealing with meromorphic AΔEs of the multiplicative form (2.1). To study these in more detail and, in particular, to construct minimal solutions, it turns out to be convenient to also consider AΔEs of the additive form

$$f(z+ia/2) - f(z-ia/2) = \phi(z), \quad a > 0. \quad (2.13)$$

Here, $\phi(z)$ is assumed to be meromorphic in a strip $|\operatorname{Im} z| < s$, $s \in (0, \infty)$, and we restrict attention to functions $f(z)$ that are meromorphic in the strip $|\operatorname{Im} z| < s + a/2$ and that satisfy (2.13) for $|\operatorname{Im} z| < s$; the term “solution to (2.13)” will be used only for such functions. The function $\phi(z)$ and the associated AΔE (2.13) will be termed *regular* iff $\phi(z)$ is analytic in $|\operatorname{Im} z| < s$, and a solution $f(z)$ to a regular AΔE will be called *regular* iff $f(z)$ is analytic in $|\operatorname{Im} z| < s + a/2$.

Obviously, taking logarithms of a regular AΔE of the multiplicative form (2.1) leads to a regular AΔE of the additive form (2.13). Since the meromorphic function $\Phi(z)$ may have zeros and/or poles for $|\operatorname{Im} z| \geq s$, its logarithm may have branch points for $|\operatorname{Im} z| \geq s$. Such branch points are irrelevant for studying the AΔE (2.1), and therefore we restrict attention to the strip $|\operatorname{Im} z| < s$ in the additive case.

The above-mentioned notions and results connected to (2.1) have obvious analogs for (2.13). In particular, a regular solution $f(z)$ to a regular AΔE (2.13) will be termed *minimal* iff it is polynomially bounded in $|\operatorname{Im} z| \leq a/2$, and a necessary condition for the existence of minimal solutions is that $\phi(z)$ be polynomially bounded on \mathbb{R} . Of course, in the additive case two minimal solutions can only differ by a constant, cf. the proof of Theorem II.1.

Let us now compare the above to the older literature on first order AΔEs, cf. in particular Refs. 5–8. Here, one usually considers additive AΔEs of the form

$$u(w+1) - u(w) = b(w). \quad (2.14)$$

Of course, these are essentially equal to (2.13), as follows by making the change of variables $z = ia(w + 1/2)$ in (2.13). But these different conventions reflect a different emphasis. Indeed, our main interest is in the behavior of $\phi(z)$ and associated solutions in the strip $|\operatorname{Im} z| \leq a/2$; in particular, we shall obtain representations for minimal solutions that hold true in this strip, cf. the next two subsections.

By contrast, Nörlund⁵ singles out the “principal solution” (Hauptlösung) to (2.14) by imposing conditions on $b(w)$ for $\operatorname{Re} w \rightarrow \infty$; accordingly, his principal solution can be characterized among all other solutions by its having the slowest possible increase for $\operatorname{Re} w \rightarrow \infty$. The principal solution equals the obvious iteration solution to (2.14) whenever $b(w)$ goes to 0 sufficiently fast for $\operatorname{Re} w \rightarrow \infty$, but it can be defined for larger classes of right-hand sides by modifying the iteration, cf. *loc. cit.* Chapters 3 and 4. As we have already seen [cf. $\Phi_3(z)$ in (2.6)], an iteration solution need not be regular, and so, *a fortiori*, it need not be minimal. Moreover, minimality concerns the asymptotics for $\operatorname{Im} w \rightarrow \pm\infty$, and not $\operatorname{Re} w \rightarrow \pm\infty$.

If one writes the hyperbolic and elliptic AΔEs occurring below (for which we construct minimal solutions) in the form (2.14), then Nörlund’s conditions are violated, and no principal solution exists. On the other hand, Nörlund’s conditions allow right-hand side functions $\phi(z)$ in (2.13) that are not polynomially bounded on \mathbb{R} ; in that case, (2.13) does not admit minimal solutions. For the regular trigonometric and rational AΔEs occurring below, both Nörlund’s and our solution methods apply, and the principal solution is then a minimal solution. Our Fourier series representation for the trigonometric case is however very different from the representations for the principal solution occurring in Ref. 5.

B. Fourier transform solutions

In this subsection we obtain minimal solutions to a large class of AΔEs by exploiting Fourier transformation on $L^2(\mathbb{R})$. (This class contains the AΔEs that occur in the hyperbolic context, cf. the Introduction.) Our normalization reads

$$\hat{\Psi}(y) \equiv \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \Psi(x) e^{ixy} \quad (2.15)$$

so that

$$\Psi(x) = \int_{-\infty}^{\infty} dy \hat{\Psi}(y) e^{-ixy}. \quad (2.16)$$

Of course, we may and will use the definition (2.15) for $\Psi \in L^1(\mathbb{R})$, too; in this case, recall $\hat{\Psi}(y) \rightarrow 0$ for $y \rightarrow \pm\infty$ (Riemann-Lebesgue lemma). We also have occasion to use the distributional Fourier transform

$$\int_{-\infty}^{\infty} dy e^{-2iyz} \mathbf{P} \frac{1}{\operatorname{sh} ay} = -\frac{i\pi}{a} \operatorname{th} \left(\frac{\pi z}{a} \right), \quad |\operatorname{Im} z| < a/2, \quad (2.17)$$

where \mathbf{P} denotes the principal value. (This formula can be verified by a straightforward contour integration.)

Theorem II.2: Assume $\phi(z)$ is a function with the following properties:

$$\phi(z) \text{ is analytic in a strip } |\operatorname{Im} z| < s, s \in (0, \infty), \quad (2.18)$$

$$\phi(x) \in L^1(\mathbb{R}), \quad (2.19)$$

$$\hat{\phi}(y) \in L^1(\mathbb{R}), \quad (2.20)$$

$$\hat{\phi}(y) = O(y), y \rightarrow 0. \quad (2.21)$$

Then the AΔE

$$f(z + ia/2) - f(z - ia/2) = \phi(z), \quad a > 0, \quad |\operatorname{Im} z| < s, \quad (2.22)$$

has a unique solution $f(a; z)$ such that

$$f(a; z) \text{ is analytic in the strip } |\operatorname{Im} z| < s + a/2, \quad (2.23)$$

$$f(a; z) \text{ is bounded in the strip } |\operatorname{Im} z| \leq a/2, \quad (2.24)$$

$$\lim_{x \rightarrow \pm \infty} f(a; x + it) = 0, \quad t \in [-a/2, a/2]. \quad (2.25)$$

Explicitly, this solution can be written as

$$f(a; z) = \int_{-\infty}^{\infty} dy \frac{\hat{\phi}(2y)}{\operatorname{sh} ay} e^{-2iyz}, \quad |\operatorname{Im} z| \leq a/2, \quad (2.26)$$

or as

$$f(a; z) = \frac{1}{2ia} \int_{-\infty}^{\infty} du \phi(u) \operatorname{th} \frac{\pi}{a}(z - u), \quad |\operatorname{Im} z| < a/2. \quad (2.27)$$

It satisfies the addition formula

$$f\left(\frac{a}{k}; z\right) = \sum_{j=1}^k f\left(a; z + \frac{ia}{2k}(k+1-2j)\right). \quad (2.28)$$

If $\phi(z)$ is even/odd, then $f(a; z)$ is odd/even. Finally, let $\psi(x)$ be the following primitive of $\phi(x)$, $x \in \mathbb{R}$:

$$\psi(x) = \frac{1}{2} \left(\int_{-\infty}^x du \phi(u) - \int_x^{\infty} du \phi(u) \right). \quad (2.29)$$

Then one has

$$\lim_{a \rightarrow 0} ia f(a; z) = \psi(z) \quad (2.30)$$

uniformly on compact subsets of the strip $|\operatorname{Im} z| < s$.

Proof: First we prove uniqueness. Thus, let $d(z)$ be the difference of two solutions to (2.22) with properties (2.23)–(2.25). Then $d(z)$ is an analytic function in $|\operatorname{Im} z| < s + a/2$, satisfying $d(z + ia/2) = d(z - ia/2)$ for $|\operatorname{Im} z| < s$. Therefore, $d(z)$ extends to an ia -periodic entire function. By virtue of (2.24), $d(z)$ is bounded in the period strip $|\operatorname{Im} z| \leq a/2$, so $d(z)$ is constant in view of Liouville's theorem. On account of (2.25) this constant equals 0, so uniqueness follows.

Next, we use (2.19) and (2.21) to infer that the function $\hat{\phi}(2y)/\operatorname{sh} ay$ is bounded and satisfies

$$|\hat{\phi}(2y)/\operatorname{sh} ay| = o(e^{-a|y|}), \quad y \rightarrow \pm \infty. \quad (2.31)$$

Thus, defining a function $f(z)$ by the rhs of (2.26), it is clear that $f(z)$ is analytic in $|\operatorname{Im} z| < a/2$ and that $f(x + it)$ converges to 0 for $x \rightarrow \pm \infty$ and $|t| < a/2$. Moreover, using also (2.20), we infer that the functions

$$b_{\pm}(x) \equiv \int_{-\infty}^{\infty} dy \frac{\hat{\phi}(2y)}{\operatorname{sh} ay} e^{\pm ay} e^{-2iyx}, \quad x \in \mathbb{R}, \quad (2.32)$$

are continuous and vanish at $\pm \infty$, and that we have

$$\lim_{t \uparrow 1} f(x \pm ita/2) = b_{\pm}(x) \quad (2.33)$$

uniformly on \mathbb{R} .

Now consider the auxiliary function

$$A(z) \equiv f(z - ia/2) + \phi(z). \quad (2.34)$$

Clearly, $A(z)$ is analytic in the strip

$$S_+ \equiv \{z \in \mathbb{C} \mid \operatorname{Im} z \in (0, \gamma)\}, \quad \gamma \equiv \min(s, a), \quad (2.35)$$

and $A(x + i\epsilon)$ converges to $b_-(x) + \phi(x)$ as $\epsilon \downarrow 0$, uniformly for x in compact subsets of \mathbb{R} . But from (2.32) we have

$$b_+(x) - b_-(x) = 2 \int_{-\infty}^{\infty} dy \hat{\phi}(2y) e^{-2iyx} = \phi(x), \quad (2.36)$$

so this boundary value is equal to $b_+(x)$. On the other hand, the function $f(z + ia/2)$ is analytic in the strip $\operatorname{Im} z \in (-a, 0)$ and converges uniformly on \mathbb{R} to $b_+(x)$ as $\operatorname{Im} z \uparrow 0$. Consequently, we may invoke Painlevé's lemma to deduce that $f(z + ia/2)$ extends to an analytic function in $\operatorname{Im} z \in (-a, \gamma)$, which coincides with $A(z)$ when $z \in S_+$. That is, the AΔE (2.22) holds true for $z \in S_+$.

We may now exploit (2.22) for $z \in S_+$ to deduce that $f(z)$ extends to an analytic function in $|\operatorname{Im} z| < s + a/2$. Since the functions $f(x \pm ia/2)$ equal $b_{\pm}(x)$, they converge to 0 for $x \rightarrow \pm\infty$. Moreover, recalling the definition of $f(z)$, we obtain

$$|f(z)| \leq \int_{-\infty}^{\infty} dy \frac{|\hat{\phi}(2y)|}{|\operatorname{sh} ay|} e^{a|y|}, \quad |\operatorname{Im} z| \leq a/2, \quad (2.37)$$

and in view of (2.20) and (2.21) the rhs is finite. Therefore, the rhs of (2.26) defines a solution to (2.22) with the properties (2.23)–(2.25).

Next, we prove (2.27). Replacing the integral in (2.26) by a principal value integral, and the functions ϕ and $\hat{\phi}$ in (2.27) and (2.26) by a Schwartz space function χ and its Fourier transform $\hat{\chi}$, resp., the equality of the resulting integrals is clear from (2.17) and the Plancherel relations. Since $S(\mathbb{R})$ is dense in $L^1(\mathbb{R})$, we deduce (2.27) from (2.26).

The function at the rhs of (2.28) obviously solves (2.22) with a replaced by a/k . Since it also has the properties (2.23)–(2.25) that uniquely determine $f(a/k; z)$, we obtain (2.28). Alternatively, (2.28) follows directly from the representation (2.26) by using the elementary identity

$$\sum_{j=1}^k \exp\left(\frac{ay}{k}(k+1-2j)\right) = \frac{\operatorname{sh}(ay)}{\operatorname{sh}(ay/k)}. \quad (2.38)$$

The parity assertion can be read off from both of the representations (2.26) and (2.27).

It remains to prove the last assertion. To this end we first observe that the representation (2.27) entails that (2.30) holds true pointwise for $z = x \in \mathbb{R}$. Next, we use the bound (2.37) and the assumptions (2.20) and (2.21) to infer that the function $af(a; z)$ remains bounded by an a -independent constant in the strip $|\operatorname{Im} z| \leq a/2$ as $a \rightarrow 0$. By iteration of the AΔE (2.22) we now deduce that $af(a; z)$ remains bounded in compact subsets of the strip $|\operatorname{Im} z| < s$ as $a \rightarrow 0$. Therefore, the last assertion follows from Vitali's theorem. \square

For our purposes the conditions (2.18)–(2.21) on $\phi(z)$ are sufficiently weak. In general, however, the conditions (2.20) and (2.21) may be difficult to check. Requiring solely (2.18) and (2.19), the rhs of (2.27) defines a function $f(z)$ that is clearly analytic in the strip $|\operatorname{Im} z| < a/2$ and that satisfies

$$\lim_{x \rightarrow \pm\infty} f(x \pm it) = \pm \frac{1}{2ia} \int_{-\infty}^{\infty} du \phi(u), \quad t \in (-a/2, a/2). \quad (2.39)$$

We conjecture that this function is in fact a solution to (2.22) satisfying (2.23) and (2.24).

Returning to the assumptions of the theorem, let us note that (2.21) entails that the primitive $\psi(x)$ (2.29) vanishes at $\pm\infty$. Thus, writing $\phi(u) = \psi'(u)$ in the representation (2.27), and integrating by parts, we obtain the formula

$$f(a; z) = \frac{\pi}{2ia^2} \int_{-\infty}^{\infty} du \frac{\psi(u)}{\operatorname{ch}^2 \frac{\pi}{a}(z-u)}, \quad |\operatorname{Im} z| < a/2. \quad (2.40)$$

Comparing this representation to Eq. (14) in Chapter 4 of Nörlund's monograph,⁵ one sees that the solution $f(a; z)$ and Nörlund's principal solution differ only by a constant whenever $\phi(z)$ satisfies not only the assumptions of Theorem II.2, but also the various restrictions that Nörlund needs for his principal solution to exist and admit the representation (14) in *loc. cit.* (As already mentioned, his assumptions on $\phi(z)$ are quite different from ours, cf. the discussion after (2.14).)

It is also of interest to observe that the assumptions (2.19)–(2.21) entail that $\hat{\phi}(y)$ is an $L^2(\mathbb{R})$ -function in the domain of the unbounded self-adjoint multiplication operator $1/\operatorname{sh}(ay/2)$. From this point of view the function $f(a; x)$, $x \in \mathbb{R}$, given by (2.26), is the obvious $L^2(\mathbb{R})$ -solution to (2.22) with $z \in \mathbb{R}$, reinterpreted as a Hilbert space equation. (Indeed, the function $\hat{f}(a; y)$ —being equal to $\hat{\phi}(y)/2\operatorname{sh}(ay/2)$ —is in the domain of multiplication by $\exp(\pm ay/2)$.)

We proceed by generalizing the above key result Theorem II.2. We shall detail this generalization in the multiplicative context (2.1); the additive version will be clear from this.

Theorem II.3: Assume $\Phi(z)$ is a meromorphic function that has no poles and zeros in the strip $|\operatorname{Im} z| < s$ for some $s \in (0, \infty)$. Setting

$$\phi_l(z) \equiv \left(\frac{d}{dz} \right)^l \ln \Phi(z), \quad l \in \mathbb{N}, \quad (2.41)$$

assume there exists $k \in \mathbb{N}^*$ such that $\phi(z) \equiv \phi_k(z)$ satisfies (2.18)–(2.21). Then the $A\Delta E$

$$\frac{F(z + ia/2)}{F(z - ia/2)} = \Phi(z) \quad (2.42)$$

admits minimal solutions. Any minimal solution can be written as

$$F(z) = \exp(e(z) + P(z)), \quad (2.43)$$

where

$$e(z) \equiv \int_{-\infty}^{\infty} dy \frac{\hat{\phi}(2y)}{\operatorname{sh} ay} (-2iy)^{-k} \left(e^{-2iyz} - \sum_{j=0}^{k-1} \frac{(-2iyz)^j}{j!} \right), \quad |\operatorname{Im} z| \leq a/2, \quad (2.44)$$

and

$$P(z) \equiv \sum_{j=0}^k c_j z^j / j!, \quad c_0, \dots, c_k \in \mathbb{C}. \quad (2.45)$$

The coefficients c_2, \dots, c_k are uniquely determined, whereas c_1 is uniquely determined mod $2\pi/a$.

Proof: Consider the AΔEs

$$f_l(z + ia/2) - f_l(z - ia/2) = \phi_l(z), \quad l = 0, \dots, k. \quad (2.46)$$

By virtue of Theorem II.2 the function

$$f_k(z) \equiv \int_{-\infty}^{\infty} dy \frac{\hat{\phi}(2y)}{\text{sh}ay} e^{-2iyz}, \quad |\text{Im}z| \leq a/2, \quad (2.47)$$

admits an analytic continuation to $|\text{Im}z| < s + a/2$ and satisfies (2.46) with $l = k$. Introducing

$$f_{k-1}(z) \equiv c_k z + \int_0^z ds f_k(s), \quad c_k \in \mathbb{C}, \quad (2.48)$$

we infer that the rhs of the resulting equation

$$f_{k-1}(z + ia/2) - f_{k-1}(z - ia/2) = iac_k + \int_{z-ia/2}^{z+ia/2} ds f_k(s) \quad (2.49)$$

equals $\phi_{k-1}(z)$ for a suitable choice of c_k [since its z -derivative equals $\phi_k(z)$]; specifically, we may and will choose c_k such that

$$iac_k + \int_{-ia/2}^{ia/2} ds f_k(s) = \phi_{k-1}(0). \quad (2.50)$$

Proceeding recursively, we obtain functions $f_k(z), f_{k-1}(z), \dots, f_0(z)$ related by

$$f_{l-1}(z) = c_l z + \int_0^z ds f_l(s), \quad l = 1, \dots, k, \quad (2.51)$$

with c_l given by

$$c_l = \frac{1}{ia} \left(\phi_{l-1}(0) - \int_{-ia/2}^{ia/2} ds f_l(s) \right), \quad l = 1, \dots, k. \quad (2.52)$$

Then $f_l(z), l \in \{0, \dots, k\}$, is analytic in $|\text{Im}z| < s + a/2$ and is a minimal solution to (2.46). Moreover, from (2.51) and (2.47) one easily sees that $f_0(z)$ equals the sum of $e(z)$ and a polynomial of degree $\leq k$. The proof can now be completed by invoking Theorem II.1. \square

In Appendix A we show (among other things) how the above results can be used to arrive at the psi and gamma functions, and derive various salient features along the way. Here, we add two applications exemplifying the above, yielding identities we have occasion to use later on. First, consider the function

$$F(z) = \text{cth}z - \frac{\pi}{a} \text{cth} \frac{\pi z}{a}. \quad (2.53)$$

It satisfies the AΔE

$$F(z+ia/2)-F(z-ia/2)=\operatorname{cth}(z+ia/2)-\operatorname{cth}(z-ia/2)\equiv\chi(z). \quad (2.54)$$

Inverting (2.17) yields the distributional Fourier transforms

$$\int_{-\infty}^{\infty} dx \operatorname{cth} \alpha(x \pm i\beta) e^{ixy} = \frac{i\pi}{\alpha} \frac{\exp \pm (-\pi y/2\alpha + \beta y)}{\operatorname{sh}(\pi y/2\alpha)}, \quad \alpha > 0, \quad \beta \in (0, \pi/\alpha), \quad (2.55)$$

so we have

$$\hat{\chi}(y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \chi(x) e^{ixy} = i \frac{\operatorname{sh}(a-\pi)/2}{\operatorname{sh} y \pi/2}, \quad a \in (0, 2\pi). \quad (2.56)$$

Thus, $\chi(z)$ satisfies the assumptions (2.18)–(2.20), but not (2.21). But $\phi(z) \equiv \chi'(z)$ does satisfy (2.18)–(2.21), since

$$\hat{\phi}(y) = y \frac{\operatorname{sh}(a-\pi)/2}{\operatorname{sh} y \pi/2}. \quad (2.57)$$

Therefore, we obtain a solution

$$f(z) = 4 \int_0^{\infty} dy \frac{y \operatorname{sh}(a-\pi)y}{\operatorname{sh} a y \operatorname{sh} \pi y} \cos 2yz \quad (2.58)$$

to the AΔE (2.22). Now since $F'(z)$ satisfies (2.22), too, and obviously has the properties (2.23)–(2.25), we must have $f(z) = F'(z)$, by uniqueness. Integrating the resulting identity w.r.t. z , we obtain

$$\operatorname{cth} z - \frac{\pi}{a} \operatorname{cth} \frac{\pi z}{a} = 2 \int_0^{\infty} dy \frac{\operatorname{sh}(a-\pi)y}{\operatorname{sh} a y \operatorname{sh} \pi y} \sin 2yz. \quad (2.59)$$

Here we have $a \in (0, 2\pi)$ in view of the restriction in (2.56). But for $z \in \mathbb{R}$ the integral converges for any $a > 0$, and so it readily follows that (2.59) holds for any $a > 0$ (taking $|\operatorname{Im} z|$ small enough, of course). Integrating once more now yields

$$\ln(\operatorname{sh} z) - \ln\left(\frac{a}{\pi} \operatorname{sh} \frac{\pi z}{a}\right) = \int_0^{\infty} \frac{dy}{y} \frac{\operatorname{sh}(a-\pi)y}{\operatorname{sh} a y \operatorname{sh} \pi y} (1 - \cos 2yz), \quad a > 0. \quad (2.60)$$

Second, consider the function

$$h(z) \equiv \frac{z}{a} \operatorname{cth} \frac{\pi z}{a}. \quad (2.61)$$

It satisfies the AΔE

$$h(z+ia/2) - h(z-ia/2) = i \operatorname{th} \frac{\pi z}{a}. \quad (2.62)$$

Therefore, $h''(z)$ satisfies the AΔE

$$f(z+ia/2) - f(z-ia/2) = \frac{i\pi}{a} \frac{d}{dz} \left(\frac{1}{\operatorname{ch}^2} \frac{\pi z}{a} \right) \equiv \phi(z). \quad (2.63)$$

Now one readily verifies

$$\hat{\phi}(y) = \frac{ay^2}{2\pi \operatorname{sh}(ay/2)}, \quad a > 0, \quad (2.64)$$

so $\phi(z)$ satisfies the assumptions of Theorem II.2. The resulting solution

$$f(z) = \frac{4a}{\pi} \int_0^\infty dy \frac{y^2}{\operatorname{sh}^2 ay} \cos 2yz \quad (2.65)$$

must then be equal to $h''(z)$, since $h''(z)$ clearly has the properties (2.23)–(2.25). Integrating twice w.r.t. z we now obtain

$$\pi z \operatorname{cth} \frac{\pi z}{a} = a + a^2 \int_0^\infty dy \frac{(1 - \cos 2yz)}{\operatorname{sh}^2 ay}, \quad a > 0. \quad (2.66)$$

The identities (2.66) and (2.60) can be combined to evaluate integrals occurring below. First, they entail that for $a \in (0, \pi)$ one has

$$\frac{a\pi}{a-\pi} \ln \left(\frac{\pi \operatorname{sh} z}{a \operatorname{sh} \frac{\pi z}{a}} \right) - \pi z \operatorname{cth} \frac{\pi z}{a} + a = \int_0^\infty dy \left(\frac{a\pi \operatorname{sh}(a-\pi)y}{(a-\pi)y \operatorname{sh} ay \operatorname{sh} \pi y} - \frac{a^2}{\operatorname{sh}^2 ay} \right) (1 - \cos 2yz). \quad (2.67)$$

Taking $z \rightarrow \infty$ and using the Riemann-Lebesgue lemma we obtain the integral

$$\frac{a\pi}{a-\pi} \ln \frac{\pi}{a} + a = \int_0^\infty dy \left(\frac{a\pi \operatorname{sh}(a-\pi)y}{(a-\pi)y \operatorname{sh} ay \operatorname{sh} \pi y} - \frac{a^2}{\operatorname{sh}^2 ay} \right). \quad (2.68)$$

Adding the elementary integral

$$\int_0^\infty dy \left(\frac{a^2}{\operatorname{sh}^2 ay} - \frac{1}{y^2} \right) = -a \quad (2.69)$$

yields

$$\ln \frac{\pi}{a} = \int_0^\infty \frac{dy}{y} \left(\frac{\operatorname{sh}(a-\pi)y}{\operatorname{sh} ay \operatorname{sh} \pi y} - \frac{(a-\pi)}{a\pi y} \right), \quad (2.70)$$

and combining this with (2.60) we get

$$\ln \left(\operatorname{sh} \frac{\pi z}{a} \right) - \ln(\operatorname{sh} z) = \int_0^\infty \frac{dy}{y} \left(\frac{\operatorname{sh}(a-\pi)y}{\operatorname{sh} ay \operatorname{sh} \pi y} \cos 2yz - \frac{(a-\pi)}{a\pi y} \right), \quad a > 0. \quad (2.71)$$

Just as in the above examples, AΔEs with a -dependent right-hand side functions will be encountered later on. The last theorem of this subsection concerns the limit $a \rightarrow 0$ in this setting. It is convenient to use the assumptions of Theorem II.2 as a starting point; corresponding results in the slightly more general context of Theorem II.3 can then be obtained by k -fold integration.

Specifically, we consider an AΔE of the form

$$f(z + ia/2) - f(z - ia/2) = \phi_a(z), \quad a > 0, \quad (2.72)$$

where $\phi_a(z)$ satisfies the assumptions (2.18)–(2.21) for any $a \in (0, a_0]$. (Of course, the choice of a_0 is irrelevant for the limit $a \rightarrow 0$.) We allow dependence of the maximal number $s_m \in (0, \infty]$ in (2.18) on a ; in particular, one may have $s_m \rightarrow 0$ as $a \rightarrow 0$. However, we do assume that for any $a \in (0, a_0]$ the function $\phi_a(z)$ is analytic in the open right half plane

$$\mathcal{R}_0 \equiv \{z \in \mathbb{C} \mid \operatorname{Re} z > 0\}. \quad (2.73)$$

Moreover, we assume that for any compact $K \subset \mathcal{R}_0$ there exists $C_K > 0$ with

$$|\phi_a(z) - a\chi(z)| \leq C_K a^2, \quad \forall (a, z) \in (0, a_0] \times K, \quad (2.74)$$

where $\chi(z)$ is analytic in \mathcal{R}_0 .

Now let $f_a(z)$ be the unique solution to (2.72) given by Theorem II.2 (with $\phi(z) \rightarrow \phi_a(z)$, of course). Thus, $f_a(z)$ is analytic in the strip $|\operatorname{Im} z| < a/2 + s_m(a)$ and in \mathcal{R}_0 . We are now in the position to state the next result.

Theorem II.4: *In addition to the above assumptions, let*

$$|f_a(z)| \leq C_{\delta, M}, \quad \forall (a, z) \in (0, a_0] \times \{z \in \mathbb{C} \mid \operatorname{Re} z \in [\delta, M], |\operatorname{Im} z| \leq a/2\}, \quad (2.75)$$

for any $\delta > 0$ and $M > \delta$, and let the pointwise limit

$$\lim_{a \downarrow 0} f_a(z) \equiv f(z) \quad (2.76)$$

exist for any $z \in (0, \infty)$. Then $f_a(z)$ converges uniformly on compact subsets of \mathcal{R}_0 to a function $f(z)$ that is analytic in \mathcal{R}_0 . Moreover, one has

$$f'(z) = -i\chi(z), \quad z \in \mathcal{R}_0 \quad (2.77)$$

with $\chi(z)$ defined by (2.74).

Proof: Upward iteration of the AΔE (2.72) yields

$$f_a(z + iLa) = f_a(z) + \sum_{j=1}^L \phi_a(z + (j-1/2)ia), \quad |\operatorname{Im} z| \leq a/2. \quad (2.78)$$

Choosing

$$L = N[a^{-1}], \quad \operatorname{Re} z \in [\delta, M], \quad 0 < \delta < M, \quad (2.79)$$

in this equation, the arguments of ϕ_a occurring on the rhs stay in a closed rectangle $K(N, \delta, M) \subset \mathcal{R}_0$ as $a \downarrow 0$. Thus we may invoke the bounds (2.74) and (2.75) to conclude that $f_a(z)$ remains bounded for $\operatorname{Re} z \in [\delta, M], \operatorname{Im} z \in [0, N]$, as $a \downarrow 0$. Similarly, iterating downwards L times and requiring (2.79), we deduce that $f_a(z)$ remains bounded for $\operatorname{Re} z \in [\delta, M], \operatorname{Im} z \in [-N, 0]$.

Combining uniform boundedness of $f_a(z)$ on compacts of \mathcal{R}_0 with the pointwise convergence assumption (2.76), it follows from Vitali's theorem that $f_a(z)$ converges uniformly on compacts of \mathcal{R}_0 to a function $f(z)$ that is analytic in \mathcal{R}_0 . Therefore, it remains to prove (2.77).

To this end, we use (2.72) to write

$$\frac{\phi_a(z)}{ia} = f'_a(z) + \frac{1}{ia} \int_{z-ia/2}^{z+ia/2} dw (f'_a(w) - f'_a(z)), \quad z \in \mathcal{R}_0. \quad (2.80)$$

Clearly, the second term on the rhs can be majorized by

$$\sup_{w \in \{z+ib \mid b \in [-a/2, a/2]\}} |f'_a(w) - f'_a(z)|. \quad (2.81)$$

Now $f'_a(z)$ converges to $f'(z)$ uniformly on compacts $K \subset \mathcal{R}_0$, and the lhs of (2.80) converges to $-i\chi(z)$ uniformly on K [due to (2.74)], so one easily deduces (2.77). \square

We conclude this subsection with some comments on the assumptions of the theorem just obtained. In later applications, the assumptions on $\phi_a(z)$ are easily verified. Moreover, fixing $z \in \mathcal{R}_0$, the function $\phi_a(z)$ is actually real-analytic in a for $a \in \mathbb{R}$. (Note this property is stronger than (2.74).) Possibly, these properties already entail the hypotheses (2.75) and (2.76), but we believe this is not true in general. (Observe that the function $f_a(z)$ is not likely to be analytic at $a=0$ for $z \in \mathcal{R}_0$.)

The above convergence result should also be compared to the last assertion of Theorem II.2. Taking $\phi_a(z) \equiv a\phi(z)$, one sees that this assertion amounts to a simple special case of Theorem II.4—except that the analyticity region is different, and that the constant left undetermined in $f(z) = -i\psi(z)$ by (2.77) is fixed in terms of $\chi(z) = \phi(z)$. In this connection we point out that the choice of the region \mathcal{R}_0 (2.73) in which $\phi_a(z)$ is assumed to remain analytic as $a \rightarrow 0$ is determined more by convenience of exposition than by necessity. Indeed, as will be exemplified by Prop. III.7 below, the maximal region with this property can be larger, and correspondingly one can obtain convergence in this larger region.

C. Fourier series solutions

We proceed by obtaining results that will enable us to solve AΔEs occurring in the trigonometric and elliptic contexts. Correspondingly, we will be dealing with meromorphic functions that are periodic in the real direction. It is convenient to parametrize this period by $\pi/r, r \in (0, \infty)$. For $\Psi(x) \in L^2([-\pi/2r, \pi/2r], dx)$ we employ Fourier coefficients

$$\hat{\Psi}_n \equiv \frac{r}{\pi} \int_{-\pi/2r}^{\pi/2r} dx \Psi(x) e^{2inrx}, \quad n \in \mathbb{Z}, \quad (2.82)$$

so that

$$\Psi(x) = \sum_{n \in \mathbb{Z}} \hat{\Psi}_n e^{-2inrx} \quad (2.83)$$

with the series converging in the L^2 -topology.

As we have seen in the previous subsection, the AΔE (2.22) naturally leads to hyperbolic functions when ϕ satisfies (2.18)–(2.21), cf. (2.26) and (2.27). In much the same way, periodicity of $\phi(z)$ leads to the emergence of elliptic functions. It is convenient to collect some features of the functions that arise before stating the analog of Theorem II.2. First, we recall the product representations of the Weierstrass σ -function (cf., e.g., Ref. 16): We have, taking $r, a > 0$,

$$\sigma\left(z; \frac{\pi}{2r}, \frac{ia}{2}\right) = \exp(\eta z^2 r / \pi) \frac{\sin rz}{r} \prod_{k=1}^{\infty} \frac{(1 - p^k \exp(2irz))(z \rightarrow -z)}{(1 - p^k)^2} \quad (2.84)$$

with

$$p \equiv \exp(-2ar) \quad (2.85)$$

or, alternatively,

$$\sigma\left(z; \frac{\pi}{2r}, \frac{ia}{2}\right) = \exp(\eta' z^2 / ia) \frac{\operatorname{sh} \pi z / a}{\pi / a} \prod_{k=1}^{\infty} \frac{(1 - \bar{p}^k \exp(2\pi z / a))(z \rightarrow -z)}{(1 - \bar{p}^k)^2} \quad (2.86)$$

with

$$\tilde{p} \equiv \exp(-2\pi^2/ar). \quad (2.87)$$

Here, η and η' are connected by Legendre's relation

$$\eta' = i\eta ar/\pi - ir. \quad (2.88)$$

The function

$$s(r, a; z) \equiv \sigma\left(z; \frac{\pi}{2r}, \frac{ia}{2}\right) \exp(-\eta z^2 r/\pi) \quad (2.89)$$

plays a key role in the sequel. In view of (2.84) s is odd and π/r -antiperiodic, and satisfies

$$\lim_{a \rightarrow \infty} s(r, a; z) = \frac{\sin rz}{r} \quad (\text{uniformly on compacts}). \quad (2.90)$$

Moreover, using (2.86) and (2.88) one sees that s solves the AΔE

$$\frac{s(z + ia/2)}{s(z - ia/2)} = -\exp(-2irz) \quad (2.91)$$

and obeys

$$\lim_{r \rightarrow 0} s(r, a; z) = \frac{\operatorname{sh} \pi z/a}{\pi/a} \quad (\text{uniformly on compacts}). \quad (2.92)$$

Note that $s(r, a; z)$ is not a regular solution to the regular AΔE (2.91): It has zeros for $\operatorname{Im} z = 0$.

Next, using the power series for $\ln(1-x)$, $|x| < 1$, one easily verifies the identity

$$\prod_{k=1}^{\infty} (1 - p^k \exp(2irz))(z \rightarrow -z) = \exp\left(-\sum_{n=1}^{\infty} \frac{e^{-nra}}{n \operatorname{sh} nra} \cos 2nrz\right), \quad |\operatorname{Im} z| < a. \quad (2.93)$$

Combining this with (2.84) and (2.89) one obtains

$$s(r, a; z) = \frac{\sin rz}{r} \exp\left(\sum_{n=1}^{\infty} \frac{e^{-nra}}{n \operatorname{sh} nra} (1 - \cos 2nrz)\right), \quad |\operatorname{Im} z| < a. \quad (2.94)$$

From this representation we deduce

$$\frac{s'(r, a; z)}{s(r, a; z)} = r \cot rz + 2r \sum_{n=1}^{\infty} \frac{e^{-nra}}{\operatorname{sh} nra} \sin 2nrz, \quad |\operatorname{Im} z| < a. \quad (2.95)$$

Using the elementary Fourier series

$$\cot r(z + ia/2) = -i - 2i \sum_{n=1}^{\infty} e^{-nra} e^{2inrz}, \quad \operatorname{Im} z > -a/2, \quad (2.96)$$

we finally obtain

$$K(r, a; z) = ir + ir \sum_{n \in \mathbb{Z}^*} \frac{e^{-2inrz}}{\operatorname{sh} nra}, \quad |\operatorname{Im} z| < a/2, \quad (2.97)$$

where we have introduced

$$K(r, a; z) \equiv \frac{d}{dz} \ln s(r, a; z + ia/2). \quad (2.98)$$

Note that (2.92) entails

$$\lim_{r \rightarrow 0} K(r, a; z) = \frac{\pi}{a} \operatorname{th} \frac{\pi z}{a} \quad (2.99)$$

uniformly on compact subsets of $|\operatorname{Im} z| < a/2$.

Theorem II.5: Assume $\phi(z)$ is a function with the following properties:

$$\phi(z) \text{ is analytic in a strip } |\operatorname{Im} z| < s, s \in (0, \infty), \quad (2.100)$$

$$\phi(z) \text{ has period } \pi/r, \quad (2.101)$$

$$\hat{\phi}_0 = 0. \quad (2.102)$$

Then the AΔE (2.22) has a unique solution $f(a; z)$ such that

$$f(a; z) \text{ is analytic in the strip } |\operatorname{Im} z| < s + a/2, \quad (2.103)$$

$$f(a; z) \text{ has period } \pi/r, \quad (2.104)$$

$$\hat{f}_0 = 0. \quad (2.105)$$

Explicitly, this solution can be written as

$$f(a; z) = \frac{1}{2} \sum_{n \in \mathbb{Z}^*} \frac{\hat{\phi}_n e^{-2inrz}}{\operatorname{sh} nra}, \quad |\operatorname{Im} z| \leq a/2, \quad (2.106)$$

or as

$$f(a; z) = \frac{1}{2i\pi} \int_{-\pi/2r}^{\pi/2r} du \phi(u) K(r, a; z - u), \quad |\operatorname{Im} z| < a/2. \quad (2.107)$$

It obeys the addition formula (2.28). If $\phi(z)$ is even/odd, then $f(a; z)$ is odd/even. Finally, the limit relation (2.30) holds true uniformly on compact subsets of the strip $|\operatorname{Im} z| < s$, with $\psi(x)$ the primitive of $\phi(x)$ that satisfies $\hat{\psi}_0 = 0$.

Proof: In order to prove uniqueness, we argue as in the proof of Theorem II.2 to conclude that the difference $d(z)$ of two solutions satisfying (2.103)–(2.105) extends to an ia -periodic entire function. Since $d(z)$ has period π/r , too, we deduce that $d(z)$ equals a constant d . Now we have $0 = \hat{d}_0 = \pi d/r$ by (2.105), and so uniqueness follows.

Next, we define a function $f(z)$ by the rhs of (2.106). Clearly, $f(z)$ is analytic in $|\operatorname{Im} z| < a/2$ and has properties (2.104) and (2.105). Moreover, the functions

$$b_{\pm}(x) \equiv \frac{1}{2} \sum_{n \in \mathbb{Z}^*} \frac{\hat{\phi}_n e^{\pm nra}}{\operatorname{sh} nra} e^{-2inrx}, \quad x \in \mathbb{R}, \quad (2.108)$$

are smooth and π/r -periodic, and (2.33) holds true uniformly on \mathbb{R} . (Note that the Fourier coefficients $\hat{\phi}_n$ form a fast decreasing sequence, since $\phi(x)$ is real-analytic and π/r -periodic.) Since we also have

$$b_+(x) - b_-(x) = \sum_{n \in \mathbb{Z}^*} \hat{\phi}_n e^{-2inrx} = \phi(x), \quad (2.109)$$

the reasoning in the proof of Theorem II.2 can be repeated, showing that $f(z)$ solves (2.22) and has property (2.103).

The representation (2.107) follows from (2.106) and the Fourier series (2.97) by using the Plancherel relations and (2.102). The addition formula (2.28) follows in the same way as in the proof of Theorem II.2. The parity claim is obvious from either (2.106) or (2.107). Using (2.106) with $z \in \mathbb{R}$, it follows from routine arguments that

$$\lim_{a \rightarrow 0} iaf(a; x) = \sum_{n \in \mathbb{Z}^*} \frac{\hat{\phi}_n e^{-2inrx}}{-2inr} \equiv \psi(x), \quad x \in \mathbb{R}, \quad (2.110)$$

and that $\psi(x)$ is a primitive of $\phi(x)$ with $\hat{\psi}_0 = 0$. The uniform convergence assertion then follows in the same way as before from Vitali's theorem. \square

Recalling the limit (2.99), one sees that the representation (2.107) turns into (2.27) for $r \rightarrow 0$. More precisely, this holds true for functions $\phi(r; u)$ with a suitable dependence on r . Clearly, one needs some restrictions on this dependence to ensure uniform convergence for z in compacts of the strip $|\operatorname{Im} z| < a/2$ (say), but we shall not pursue this. (For an explicit example, see Prop. III.12 in Subsection III B.)

We continue with an analog of Theorem II.3.

Theorem II.6: *With (2.18)–(2.21) replaced by (2.100)–(2.102) and (2.44) replaced by*

$$e(z) \equiv \frac{1}{2} \sum_{n \in \mathbb{Z}^*} \frac{\hat{\phi}_n}{\operatorname{sh} nra} (-2inr)^{-k} \left(e^{-2inrz} - \sum_{j=0}^{k-1} \frac{(-2inrz)^j}{j!} \right), \quad |\operatorname{Im} z| \leq a/2, \quad (2.111)$$

the assertions of Theorem II.3 hold true.

Proof: With Theorem II.2 replaced by Theorem II.5, and (2.47) by

$$f_k(z) \equiv \frac{1}{2} \sum_{n \in \mathbb{Z}^*} \frac{\hat{\phi}_n}{\operatorname{sh} nra} e^{-2inrz}, \quad |\operatorname{Im} z| \leq a/2, \quad (2.112)$$

the reasoning in the proof of Theorem II.3 applies verbatim; note that boundedness of $f_k(z)$ in the strip $|\operatorname{Im} z| \leq a/2$ entails polynomial boundedness of $f_l(z)$ in this strip. \square

We conclude this subsection with a result pertaining to AΔEs (2.72), adapting the assumptions of the previous subsection to the periodic context. Thus, for any $a \in (0, a_0]$ the right-hand side $\phi_a(z)$ is assumed to satisfy (2.100)–(2.102) and to be analytic in the open period strip

$$\mathcal{R}_r \equiv \{z \in \mathbb{C} \mid \operatorname{Re} z \in (0, \pi/r)\}. \quad (2.113)$$

Furthermore, the bound (2.74) is assumed to be valid for any compact $K \subset \mathcal{R}_r$, with $\chi(z)$ analytic in \mathcal{R}_r .

Denoting by $f_a(z)$ the unique solution to (2.72) given by Theorem II.5, we are prepared to state the analog of Theorem II.4.

Theorem II.7: Assume in addition to the above that (2.75) holds true for any $\delta \in (0, \pi/r)$ and $M \in (\delta, \pi/r)$, and that the pointwise limit (2.76) exists for any $z \in (0, \pi/r)$. Then the assertions of Theorem II.4 hold true, with \mathcal{R}_0 replaced by \mathcal{R}_r .

Proof: Taking $M < \pi/r$ in (2.79) and replacing \mathcal{R}_0 by \mathcal{R}_r , the proof of Theorem II.4 applies verbatim. \square

The comments after Theorem II.4 apply with obvious changes to Theorem II.7, so we shall not spell them out again.

III. GENERALIZED GAMMA FUNCTIONS

A. The hyperbolic case

Consider the integral

$$\int_0^\infty \frac{dy}{y} \left(\frac{\sin 2yz}{2 \operatorname{sha}_+ y \operatorname{sha}_- y} - \frac{z}{a_+ a_- y} \right) \equiv g(a_+, a_-; z), \quad (3.1)$$

where we take $a_\delta \in (0, \infty)$, $\delta = +, -$, until further notice. Obviously, this integral converges absolutely provided z belongs to the strip

$$S \equiv \{z \in \mathbb{C} \mid |\operatorname{Im} z| < (a_+ + a_-)/2\}, \quad (3.2)$$

and it defines a function g that is analytic in S . In this subsection we study the function

$$G(z) \equiv \exp(ig(z)) \quad (3.3)$$

in considerable detail. (Here and in the sequel, we suppress the dependence on a_+, a_- whenever this causes no confusion.) We shall collect our results in propositions that concern various features of $G(z)$.

Proposition III.1 (defining AΔEs): The function $G(z)$ is analytic and has no zeros in the strip S . It extends to a meromorphic function that is a minimal solution to the three AΔEs

$$\frac{G(z + ia_\delta/2)}{G(z - ia_\delta/2)} = 2 \operatorname{ch}(\pi z/a_\delta), \quad \delta = +, -, \quad (3.4)$$

and

$$\frac{G(z + i(a_+ - a_-)/2)}{G(z - i(a_+ - a_-)/2)} = \frac{\operatorname{sh}(\pi z/a_-)}{\operatorname{sh}(\pi z/a_+)}. \quad (3.5)$$

It is the unique minimal solution satisfying

$$G(0) = 1, \quad |G(x)| = 1, \quad x \in \mathbb{R}. \quad (3.6)$$

Proof: The first assertion is clear from (3.1)–(3.3). Taking $\delta = +$ in (3.4) and denoting the rhs by $\Phi(z)$, the assumptions of Theorem II.3 are satisfied, with $a = a_+$, $s = a_-/2$ and $k = 3$. Indeed, we have

$$\phi(z) \equiv \left(\frac{d}{dz} \right)^3 \ln \Phi(z) = \frac{\pi}{a_-} \left(\frac{d}{dz} \right)^2 \operatorname{th}(\pi z/a_-) \quad (3.7)$$

so that (cf. (217))

$$\hat{\phi}(y) = \frac{-iy^2}{2\operatorname{sh}(a_-y/2)}. \quad (3.8)$$

From this the properties (2.18)–(2.21) are evident.

As a consequence the AΔE at hand admits minimal solutions; these can be written as (2.43)–(2.44) with $k=3$ and

$$\begin{aligned} e(z) &= -\frac{1}{4} \int_{-\infty}^{\infty} \frac{dy}{\operatorname{sha}_+ y \operatorname{sha}_- y} (e^{-2iyz} - (1 - 2iyz - 2y^2 z^2)) \\ &= i \int_0^{\infty} \frac{dy}{\operatorname{sha}_+ y \operatorname{sha}_- y} \left(\frac{\sin 2yz}{2y} - z \right). \end{aligned} \quad (3.9)$$

To determine c_1, c_2, c_3 we follow the proof of Theorem II.3. Thus, we start from

$$f_3(z) = -4i \int_0^{\infty} dy y^2 \cos(2yz) / \operatorname{sha}_+ y \operatorname{sha}_- y, \quad (3.10)$$

cf. (2.47). Then we get

$$\int_0^z ds f_3(s) = -2i \int_0^{\infty} dy y \sin(2yz) / \operatorname{sha}_+ y \operatorname{sha}_- y \quad (3.11)$$

so that

$$\int_{-ia_+/2}^{ia_+/2} ds f_3(s) = 4 \int_0^{\infty} dy y / \operatorname{sha}_- y = \left(\frac{\pi}{a_-} \right)^2. \quad (3.12)$$

From (2.50) we then have $c_3=0$, and so

$$f_2(z) = -2i \int_0^{\infty} dy y \sin(2yz) / \operatorname{sha}_+ y \operatorname{sha}_- y. \quad (3.13)$$

Now $f_2(z)$ is odd, so (2.52) yields $c_2=0$. Hence,

$$f_1(z) = i \int_0^{\infty} dy (\cos(2yz) - 1) / \operatorname{sha}_+ y \operatorname{sha}_- y, \quad (3.14)$$

cf. (2.51), so that

$$\int_0^z ds f_1(s) = i \int_0^{\infty} dy \left(\frac{\sin 2yz}{2y} - z \right) / \operatorname{sha}_+ y \operatorname{sha}_- y = e(z), \quad (3.15)$$

cf. (3.9). Now we have

$$\pm e(\pm ia_+/2) = \frac{1}{2} \int_0^{\infty} dy \left(\frac{a_+}{\operatorname{sha}_+ y \operatorname{sha}_- y} - \frac{1}{y \operatorname{sha}_- y} \right). \quad (3.16)$$

Also, recalling (A33) and (A34), we may write

$$\ln 2 = \int_0^\infty dy \left(\frac{1}{a-y^2} - \frac{1}{y \operatorname{sha} y} \right). \quad (3.17)$$

Using (2.52) once more, we obtain

$$c_1 = (ia_+)^{-1} (\ln 2 - e(ia_+/2) + e(-ia_+/2)) = i \int_0^\infty dy \left(\frac{1}{\operatorname{sha}_+ y \operatorname{sha}_- y} - \frac{1}{a_+ a_- y^2} \right). \quad (3.18)$$

Combining (2.51) with (3.15) now yields

$$f_0(z) = c_1 z + e(z) = ig(a_+, a_-; z), \quad (3.19)$$

cf. (3.1). In view of (3.3), this entails that $G(z)$ solves (3.4) with $\delta = +$. Since the function G is manifestly symmetric in a_+, a_- , it solves (3.4) with $\delta = -$, too.

To prove that G also satisfies the AΔE (3.5), we observe that we may write

$$\frac{G(z + i(a_+ - a_-)/2)}{G(z - i(a_+ - a_-)/2)} = \frac{G(z - ia_-/2 + ia_+/2)}{G(z - ia_-/2 - ia_+/2)} \cdot \frac{G(z - ia_+/2 - ia_-/2)}{G(z - ia_+/2 + ia_-/2)}. \quad (3.20)$$

From (3.4) we now deduce that (3.5) holds true. Finally, the uniqueness assertion is clear from Theorem II.1. \square

We point out that the identity (2.71) can also be obtained from the AΔE (3.5). Similarly, the proposition entails the identity

$$\int_0^\infty \frac{dy}{y} \left(\frac{1}{ay} - \frac{\cos 2yz}{\operatorname{sha} y} \right) = \ln \left(2 \operatorname{ch} \frac{\pi z}{a} \right), \quad a > 0, \quad |\operatorname{Im} z| < a/2. \quad (3.21)$$

Indeed, this identity amounts to the function ig [as given by (3.1)] satisfying the additive versions of the AΔEs (3.4). The integral (3.21) can also be derived directly from (A33), (A34) and (2.17). In this way one can obtain a shorter proof of (3.4). The above proof, however, shows how the function $G(z)$ emerges from the general theory presented in Subsection II B, when one takes one of the AΔEs (3.4) as a starting point.

Proposition III.2 (automorphy properties): *One has*

$$G(-z) = 1/G(z), \quad (3.22)$$

$$G(a_-, a_+; z) = G(a_+, a_-; z), \quad (3.23)$$

$$G(\lambda a_+, \lambda a_-; \lambda z) = G(a_+, a_-; z), \quad \lambda \in (0, \infty). \quad (3.24)$$

For any $M, N \in \mathbb{N}^*$ one has the multiplication formula

$$G\left(\frac{a_+}{M}, \frac{a_-}{N}; z\right) = \prod_{j=1}^M \prod_{k=1}^N G\left(a_+, a_-; z + \frac{ia_+}{2M}(M+1-2j) + \frac{ia_-}{2N}(N+1-2k)\right). \quad (3.25)$$

Proof: All of these properties readily follow from the integral representation (3.1)–(3.3) and meromorphy of G . Indeed, the first three are immediate from (3.1). Taking first $N=1$ in (3.25), and using (3.1) and the identity (2.38) to rewrite the rhs, one obtains the desired result for $G(a_+/M, a_-; z)$; the general case then follows by using (3.23). \square

Note that when one takes $M=N$ in the formula (3.25), one can use (3.24) to write its lhs as $G(a_+, a_-; Nz)$.

Proposition III.3 (zeros, poles, residues): *The zeros and poles of $G(z)$ are given by*

$$z_{kl}^+ \equiv i(a_+(k+1/2) + a_-(l+1/2)), \quad k, l \in \mathbb{N} \quad (\text{zeros}), \quad (3.26)$$

$$z_{kl}^- \equiv -z_{kl}^+, \quad k, l \in \mathbb{N} \quad (\text{poles}). \quad (3.27)$$

For a given $(k_0, l_0) \in \mathbb{N}^2$, the multiplicities of the pole $z_{k_0 l_0}^-$ and zero $z_{k_0 l_0}^+$ are equal to the number of distinct pairs $(k, l) \in \mathbb{N}^2$ such that $z_{kl}^+ = z_{k_0 l_0}^+$; in particular, for $a_+ / a_- \notin \mathbb{Q}$ all poles and zeros are simple. The pole at z_{00}^- is simple and has residue

$$r_{00} = \frac{i}{2\pi} (a_+ a_-)^{1/2}. \quad (3.28)$$

More generally, if the quantity

$$t_{kl} \equiv \prod_{m=1}^k \sin(\pi m a_+ / a_-) \prod_{n=1}^l \sin(\pi n a_- / a_+) \quad (3.29)$$

is non-zero, then the pole at z_{kl}^- is simple and has residue

$$r_{kl} = (-)^{kl} (-1/2)^{k+l} r_{00} / t_{kl}. \quad (3.30)$$

Conversely, if z_{kl}^- is a simple pole, then one has $t_{kl} \neq 0$.

Proof: In view of (3.23), we may assume $a_+ \leq a_-$. Iterating the AΔE (3.4) with $\delta = +$ we obtain

$$G(z - iM a_+) = P_M(z) G(z), \quad M \in \mathbb{N}^*, \quad (3.31)$$

where

$$P_M(z) \equiv \left(\prod_{m=1}^M 2 \operatorname{ch} \frac{\pi}{a_-} (z - i a_+ (m - 1/2)) \right)^{-1}. \quad (3.32)$$

Now the poles of $P_M(z)$ occur at (and only at)

$$z_{ml} \equiv i a_+ (m - 1/2) - i a_- (l + 1/2), \quad m = 1, \dots, M, \quad l \in \mathbb{Z}. \quad (3.33)$$

Introducing the strip

$$S_- \equiv \{z \in \mathbb{C} \mid \operatorname{Im} z \in a_- [-1/2, 1/2]\}, \quad (3.34)$$

and fixing $m \in \{1, \dots, M\}$, there exists a unique $l \geq 0$ such that $z_{ml} \in S_-$. Since $G(z)$ is analytic and non-zero in S_- , it now follows from (3.31) that G has M and only M poles (counting multiplicity) in the shifted strip $S_- - iM a_+$; these occur at $z_{kl}^-, k = 0, \dots, M-1$, with $l \in \mathbb{N}$ uniquely determined by k and M .

Now for a given pair $(k_0, l_0) \in \mathbb{N}^2$ one can find some $M_0 > k_0$ such that $z_{k_0 l_0}^- \in S_- - iM_0 a_+$ (since the shifted strips cover the lower half plane). Also, for any pair $(k, l) \in \mathbb{N}^2$ such that $z_{kl}^- = z_{k_0 l_0}^-$, one must have $k < M_0$ (since $z_{kl}^- \in S_- - iM_0 a_+$ entails $a_+(k+1/2) + a_-(l+1/2) \leq a_+ M_0$). Consequently, the multiplicity of the pole of $P_{M_0}(z)$ at $z = z_{k_0 l_0}^- + iM_0 a_+$ equals the number of pairs satisfying $z_{kl}^- = z_{k_0 l_0}^-$.

The upshot is that the poles of $G(a_+, a_-; z)$ in the lower half plane are given by (3.27) and have the asserted multiplicity. Since G is non-zero in S_- and P_M has no zeros at all, it follows from (3.31) that G is non-zero in the lower half plane. Recalling (3.22), the first two assertions easily follow.

To prove the third one, we use (3.4) with $\delta = +$ to get

$$G(z - i(a_+ + a_-)/2) = \left(-2i \operatorname{sh} \frac{\pi z}{a_-} \right)^{-1} G(z + i(a_+ - a_-)/2). \quad (3.35)$$

From this we read off

$$r_{00} = \frac{ia_-}{2\pi} G(i(a_+ - a_-)/2). \quad (3.36)$$

Similarly, using (3.4) with $\delta = -$ we obtain

$$r_{00} = \frac{ia_+}{2\pi} G(i(a_- - a_+)/2). \quad (3.37)$$

Combining these two expressions for r_{00} with (3.22), we deduce

$$G(i(a_+ - a_-)/2) = (a_+/a_-)^{1/2}, \quad (3.38)$$

and so (3.28) follows. (Note that (3.1) and (3.3) entail that G is positive for $z \in i(a_+ + a_-) \times (-1/2, 1/2)$. Note also that (3.38) can be derived from (3.5).)

Finally, we exploit both AΔEs (3.4) to write

$$G(z + z_{kl}^-) = (-)^{kl+k+l} \left(\prod_{m=1}^k 2i \operatorname{sh} \frac{\pi}{a_-} (z - ima_+) \prod_{n=1}^l 2i \operatorname{sh} \frac{\pi}{a_+} (z - ina_-) \right)^{-1} G(z + z_{00}^-). \quad (3.39)$$

Taking $z \rightarrow 0$ in this identity, the remaining assertions follow. \square

In principle, the residue at $z_{k_0 l_0}^-$ can still be determined by using (3.30) even when $z_{k_0 l_0}^-$ is not a simple pole. Indeed, in that case one must have $a_+/a_- \in \mathbb{Q}$; choosing sequences $a_{\delta,n} \rightarrow a_\delta$, $\delta = +, -$, for $n \rightarrow \infty$ such that $a_{+,n}/a_{-,n} \notin \mathbb{Q}$, the residue equals the limit of the sum of the residues at the simple poles that coalesce at $z_{k_0 l_0}^-$. There is presumably an explicit formula for the limit, but we have not pursued this.

It is evident from (3.3) and the above that $g(z)$ extends from an analytic function in S to a multi-valued function with logarithmic branch points at (3.26) and (3.27). It is convenient to specialize to the branch obtained by restricting z to the cut plane $\mathbb{C}(a_+ + a_-)$, where

$$\mathbb{C}(d) \equiv \mathbb{C} \setminus \{\pm i[d/2, \infty)\}, \quad d > 0. \quad (3.40)$$

This branch will be again denoted $g(z)$. Asymptotic properties for $\operatorname{Re} z \rightarrow \pm \infty$ are most easily obtained for the special case $a_+ = a_- \equiv a$; the general case can then be handled by a comparison argument, cf. Prop. III.4 below.

We start from the identity

$$g(a, a; z) = -\frac{1}{\pi} b(\pi z/a), \quad (3.41)$$

where we have introduced

$$b(w) \equiv \int_0^w dt t \operatorname{ctht}, \quad w \in \mathbb{C}(2\pi). \quad (3.42)$$

(To see that this holds true, use (3.1) on the lhs and take z -derivatives; this yields a linear combination of the identities (2.66) and (2.69).) Next, we write $\text{cht} = \text{sht} + e^{-t}$ to obtain

$$b(w) = w^2/2 + c_+ - b_+(w), \quad \text{Re } w > 0, \quad (3.43)$$

where

$$b_+(w) \equiv \int_w^\infty dt \frac{te^{-t}}{\text{sht}}, \quad \text{Re } w > 0, \quad (3.44)$$

$$c_+ \equiv \int_0^\infty dt \frac{te^{-t}}{\text{sht}} = \frac{F_1'(0)}{2i} = \sum_{m=1}^\infty \frac{1}{2m^2} = \frac{\pi^2}{12}, \quad (3.45)$$

cf. (A8) and (A10). From this representation we read off the bounds

$$b(w) = \frac{w^2}{2} + \frac{\pi^2}{12} + O(\exp((\epsilon - 2)w)), \quad \text{Re } w \rightarrow \infty, \quad (3.46)$$

$$b'(w) = w + O(\exp((\epsilon - 2)w)), \quad \text{Re } w \rightarrow \infty. \quad (3.47)$$

Here, ϵ is a fixed positive number and the bounds hold true uniformly for $\text{Im } w$ varying over compact subsets of \mathbb{R} .

Of course, these bounds entail bounds on $g(a, a; z)$ via (3.41). More generally, they can be exploited to derive bounds on $g(a_+, a_-; z)$, as will now be detailed.

Proposition III.4 (asymptotics): Fixing $\epsilon > 0$ and setting

$$a_m \equiv \max(a_+, a_-) \quad (3.48)$$

one has

$$\pm g(a_+, a_-; z) = -\frac{\pi z^2}{2a_+a_-} - \frac{\pi}{24} \left(\frac{a_+}{a_-} + \frac{a_-}{a_+} \right) + O(\exp(\pm(\epsilon - 2\pi/a_m)z)), \quad \text{Re } z \rightarrow \pm\infty, \quad (3.49)$$

$$\pm g'(a_+, a_-; z) = -\frac{\pi z}{a_+a_-} + O(\exp(\pm(\epsilon - 2\pi/a_m)z)), \quad \text{Re } z \rightarrow \pm\infty, \quad (3.50)$$

where the bounds are uniform for $\text{Im } z$ in \mathbb{R} -compacts.

Proof: Since g is odd in z , it suffices to verify the $\text{Re } z \rightarrow \infty$ asymptotics. Now when $a_+ = a_-$, the formulas (3.49) and (3.50) are immediate from (3.41), and (3.46) and (3.47), resp. Since g is symmetric in a_+, a_- , it remains to consider the case $a_+ < a_-$.

To this end we rewrite (3.1) as

$$a_+a_-g(a_+, a_-; z) = a^2g(a, a; z) + d(z), \quad (3.51)$$

where we have introduced

$$a \equiv \left(\frac{a_+^2 + a_-^2}{2} \right)^{1/2}, \quad (3.52)$$

$$d(z) \equiv \int_0^\infty dy I(y) \sin 2yz, \quad (3.53)$$

with

$$I(y) \equiv \frac{1}{2y} \left(\frac{a_+ a_-}{\operatorname{sha}_+ y \operatorname{sha}_- y} - \frac{a^2}{\operatorname{sh}^2 a y} \right). \quad (3.54)$$

Here, we take z in the strip S (3.2), so that the integral converges (note $a_+ + a_- \leq 2a$). Now we have

$$I(y) = c(a_+, a_-)y + O(y^3), \quad y \rightarrow 0, \quad (3.55)$$

so $I(y)$ is analytic in the strip $|\operatorname{Im} y| < \pi/a_-$. Hence, fixing $z \in S$ and $r \in (0, \pi/a_-)$, we may shift contours to obtain

$$2id(z) = e^{-2rz} \int_{-\infty}^{\infty} du I(u + ir) e^{2iuz}. \quad (3.56)$$

From this we deduce that $d(z)$ and $d'(z)$ are $O(e^{-2rz})$ for $\operatorname{Re} z \rightarrow \infty$, uniformly for z in a closed substrip of S .

Combining these bounds with (3.51) and the $\operatorname{Re} z \rightarrow \infty$ asymptotics of $g(a, a; z)$, we deduce that (3.49) and (3.50) hold true uniformly for z in the strip $|\operatorname{Im} z| \leq a_+$. Finally, we exploit the AΔEs

$$g(z \pm ia_+) = g(z) \mp i \ln \left(2 \operatorname{ch} \frac{\pi}{a_-} (z \pm ia_+/2) \right) \quad (3.57)$$

to infer that the bounds hold uniformly for $|\operatorname{Im} z| \leq 2a_+$; by iteration, the proposition now follows. \square

Thus far, we have taken a_+ and a_- positive. However, fixing $z \in \mathbb{R}$, it is already obvious from (3.1) that $G(a_+, a_-; z)$ extends to a function that is analytic and non-zero for a_+, a_- in the (open) right half plane. Note this is consistent with the analytic continuation of (3.26) and (3.27): The imaginary part of the rhs is non-zero for a_+, a_- in the right half plane.

More generally, we shall now prove that G can be continued to a function that is meromorphic in a_+, a_- and z , provided the ratio variable

$$\rho \equiv a_- / a_+ \quad (3.58)$$

stays away from the negative real axis. To this end we consider the auxiliary function

$$A(\rho, \lambda) \equiv \prod_{j=0}^{\infty} F((j+1/2)\rho, \lambda), \quad \rho \in \mathbb{C}^-, \quad \lambda \in \mathbb{C}, \quad (3.59)$$

where \mathbb{C}^- denotes the cut plane (A15). In view of (B22) and (B19) this is a well-defined meromorphic function in $\mathbb{C}^- \times \mathbb{C}$. Moreover, from (A40) we readily deduce

$$A(\rho, \lambda) = \exp \left(\int_0^{\infty} \frac{dt}{t \operatorname{sh}(\rho t/2)} (2\lambda - \operatorname{sh}(\lambda t) \operatorname{cth}(t/2)) \right), \quad \rho > 0, \quad |\operatorname{Re} \lambda| < \rho. \quad (3.60)$$

Now from (3.1) and (3.3) we have

$$\begin{aligned}
G(z+ia_+/2)G(z-ia_+/2) &= \exp\left(\int_0^\infty \frac{dt}{t \operatorname{sh}(\rho t/2)} \left(\operatorname{sh}(itz/a_+) \operatorname{cth}(t/2) - \frac{2iz}{a_+} \right)\right) \\
&\quad \times \exp\left(i \int_0^\infty \frac{dt}{t} \left(\frac{2z}{a_+ \operatorname{sh}(a_-t/2a_+)} - \frac{4z}{a_-t} \right)\right) \\
&= A(\rho, -iz/a_+) \exp\left(-\frac{2iz}{a_+} \ln 2\right), \tag{3.61}
\end{aligned}$$

where we used (A33) and (A34). Next, we introduce the new variable

$$\lambda \equiv -iz/a_+ \tag{3.62}$$

and combine (3.61) and the AΔE (3.4) to deduce

$$G(a_+, a_-; z+ia_+/2)^2 = A(\rho, \lambda) \exp(2\lambda \ln 2) \cdot 2 \cos(\pi \lambda / \rho). \tag{3.63}$$

We are now prepared for the following proposition.

Proposition III.5 (meromorphic continuation): *The function $G(a_+, a_-; z)$ admits analytic continuation to a function that is meromorphic in a_+, a_- and z , provided $\rho \equiv a_-/a_+$ stays in \mathbb{C}^- . Fixing a_+, a_- with $\operatorname{Im} \rho \neq 0$, one obtains a meromorphic function whose zeros and poles are simple and located at (3.26) and (3.27), resp.*

Proof: The function

$$B(\rho, \lambda) \equiv A(\rho, \lambda) \cos(\pi \lambda / \rho) \tag{3.64}$$

is meromorphic in $\mathbb{C}^- \times \mathbb{C}$, so in view of (3.63) we need only show that for $\rho \notin \mathbb{R}$ all of its zeros and poles are double and located at

$$\lambda = k + (l+1/2)\rho, \quad k, l \in \mathbb{N} \quad (\text{zeros}), \tag{3.65}$$

$$\lambda = -k - 1 - (l+1/2)\rho, \quad k, l \in \mathbb{N} \quad (\text{poles}). \tag{3.66}$$

Recalling the definitions (2.59) and (A39), we obtain the representation

$$B(\rho, \lambda) = \cos(\pi \lambda / \rho) \prod_{j=0}^{\infty} \frac{\Gamma((j+1/2)\rho + \lambda)}{\Gamma((j+1/2)\rho - \lambda)} \frac{\Gamma(1 + (j+1/2)\rho + \lambda)}{\Gamma(1 + (j+1/2)\rho - \lambda)} \exp(-4\lambda \ln(j+1/2)\rho) \tag{3.67}$$

from which these features can be read off. \square

Of course, the proposition just proved entails that various formulas involving G can be analytically continued. We mention specifically (3.4), (3.5), (3.22)–(3.25) [note one can take $\lambda \in \mathbb{C}^*$ in (3.24)], (3.28)–(3.30), and the special values

$$G(ia_\delta - a_-_\delta/2) = (a_\delta/a_-_\delta)^{1/2}, \quad G(\pm ia_\delta/2) = 2^{\pm 1/2}, \quad \delta = +, -. \tag{3.68}$$

(These values easily follow from (3.1)–(3.5).)

We proceed by detailing the relation to the gamma function. To this end we introduce

$$H(\rho; z) \equiv G(1, \rho; \rho z + i/2) \exp(iz \ln(2\pi\rho) - 2^{-1} \ln(2\pi)), \quad \rho \in \mathbb{C}^-, \quad z \in \mathbb{C}. \tag{3.69}$$

This renormalized version of $G(a_+, a_-; z)$ is such that the two AΔEs (3.4) translate into the AΔE

$$\frac{H(\rho; z + i/2)}{H(\rho; z - i/2)} = \frac{i \operatorname{sh} \pi \rho z}{\pi \rho} \quad (3.70)$$

and functional equation

$$H(\rho; z)H(\rho; -z) = \frac{\operatorname{ch} \pi z}{\pi}. \quad (3.71)$$

(Use (3.22) to check (3.71).) We shall now show that the $\rho \rightarrow 0$ limit of $H(\rho; z)$ exists and equals $1/\Gamma(iz + 1/2)$. Accordingly, (3.70) and (3.71) turn into the AΔE and functional equation satisfied by the gamma function.

Proposition III.6 (relation to gamma function): *Taking $\rho \in (0, \infty)$, one has*

$$\lim_{\rho \downarrow 0} H(\rho; z) = 1/\Gamma(iz + 1/2) \quad (3.72)$$

uniformly for z in \mathbb{C} -compacts. More generally, fix $\epsilon \in (0, \infty)$, $\phi \in (0, \pi)$, and an arbitrary compact $K \subset \mathbb{C}$. Then there exists $\delta = \delta(\epsilon, \phi, K) \in (0, \infty)$ such that

$$|H(\rho; z)\Gamma(iz + 1/2) - 1| < \epsilon, \quad z \in K, \quad |\operatorname{Arg} \rho| \leq \pi - \phi, |\rho| \in (0, \delta]. \quad (3.73)$$

Proof: We begin by proving (3.72). Since the function $1/\Gamma(iz + 1/2)$ is entire, we need only show

$$\lim_{\rho \downarrow 0} P(\rho; z) = 1 \quad (\text{uniformly on compacts}), \quad (3.74)$$

$$P(\rho; z) \equiv H(\rho; z)\Gamma(iz + 1/2). \quad (3.75)$$

Now from Prop. III.3 we see that the poles of $\Gamma(iz + 1/2)$ are matched by zeros of $H(\rho; z)$, so that $P(\rho; z)$ has no poles and zeros in the strip

$$S_\rho \equiv \{z \in \mathbb{C} \mid |\operatorname{Im} z| < 1/2 + 1/\rho\}. \quad (3.76)$$

We continue by deriving an integral representation for $P(\rho; z)$ that holds true in S_ρ . To this end we first take $|\operatorname{Im} z| < 1/2$. Then we may use (3.3) and (3.1) to write

$$G(1, \rho; \rho z + i/2) = \exp \left(\int_0^\infty \frac{dy}{y} \left(\frac{e^{2i\rho y z} e^{-y} - e^{-2i\rho y z} e^y}{4 \operatorname{sh} y \operatorname{sh} \rho y} - \frac{iz}{y} + \frac{1}{2\rho y} \right) \right). \quad (3.77)$$

Also, from (A37) we obtain

$$\frac{\Gamma(iz + 1/2)}{(2\pi)^{1/2}} = \exp \left(\int_0^\infty \frac{dy}{y} \left(iz e^{-2\rho y} - \frac{1}{2\rho y} + \frac{e^{-2i\rho y z} (e^y - e^{-y})}{4 \operatorname{sh} y \operatorname{sh} \rho y} \right) \right). \quad (3.78)$$

Finally, combining (A37) (with $z = 1/2$) and the integral (A29), we write the remaining factor in (3.69) as

$$\exp(iz \ln(2\pi\rho)) = \exp \left(\int_0^\infty \frac{dy}{y} \left(\frac{iz}{y} - \frac{iz e^{-y}}{\operatorname{sh} y} - iz e^{-2\rho y} \right) \right). \quad (3.79)$$

Putting the pieces together, we obtain

$$P(\rho; z) = \exp\left(\frac{i}{2} \int_0^\infty \frac{dy}{y} \frac{e^{-y}}{\operatorname{sh} y \operatorname{sh} \rho y} (\sin(2\rho y z) - 2z \operatorname{sh} \rho y)\right). \quad (3.80)$$

Clearly, this representation can be analytically continued to the strip S_ρ , as announced above. Now we fix a compact $K \subset \mathbb{C}$ and note $K \subset S_\rho$ for ρ small enough. Rewriting the integral in (3.80) as

$$\frac{1}{c} \int_0^\infty dy \frac{e^{-cy} (\sin(2yz) - 2z \operatorname{sh} y)}{y^2 \operatorname{sh} y} \left(\frac{cy}{\operatorname{sh} cy} \right), \quad c \equiv 1/\rho, \quad (3.81)$$

it becomes evident that it converges to 0 for $c \rightarrow \infty$ uniformly on K . Consequently, we have now proved that (3.72) holds true uniformly on compacts.

To prove the stronger assertion (3.73), we observe that for $z \in K$ and $c > 0$ large enough, the contour in (3.81) may be rotated to $e^{i\chi}y$, $y \in [0, \infty)$, with $|\chi| \leq (\pi - \phi)/2$, cf. the proof of Theorem B.1. The resulting integral can now be estimated in an obvious way for $c \in \mathbb{C}$ with $|c|$ large enough and $|\operatorname{Arg}(e^{i\chi}c)| \leq (\pi - \phi)/2$, and then (3.73) easily follows. \square

The function $P(\rho; z)$ (3.75) is of some interest in itself: It is the unique minimal solution to the AΔE

$$\frac{F(z + i/2)}{F(z - i/2)} = \frac{\operatorname{sh} \pi \rho z}{\pi \rho z} \quad (3.82)$$

[cf. (3.70)] that satisfies $F(0) = 1, |F(x)| = 1, x \in \mathbb{R}$. Note that the representation (3.80) can be understood from Theorem II.3.

We conclude this subsection by deriving two more zero step size limits, now involving the function $G(\pi, a; \cdot)$ for $a \rightarrow 0$. (The choice $a_+ = \pi$ is notationally convenient; the scaling relation (3.24) can be used for other a_+ -values.) In fact, we shall phrase the limits in terms of the branch $g(z) = -i \ln G(z)$ defined in the cut plane $\mathbb{C}(\pi + a)$, cf. the paragraph containing (3.40). Introducing the functions

$$d_a(\lambda, \mu; z) \equiv g(\pi, a; z + i\lambda a) - g(\pi, a; z + i\mu a), \quad z \in \mathbb{C}(\pi + a), \quad \lambda, \mu \in \mathbb{R}, \quad (3.83)$$

$$D_a(z) \equiv a g(\pi, a; z), \quad z \in \mathbb{C}(\pi + a), \quad (3.84)$$

we are prepared for the following proposition.

Proposition III.7 (zero step size limits): *One has*

$$\lim_{a \downarrow 0} d_a(\lambda, \mu; z) = -i(\lambda - \mu) \ln(2 \operatorname{ch} z), \quad \lambda, \mu \in \mathbb{R}, \quad (3.85)$$

$$\lim_{a \downarrow 0} D_a(z) = - \int_0^z dw \ln(2 \operatorname{ch} w), \quad (3.86)$$

uniformly on compact subsets of the cut plane $\mathbb{C}(\pi)$ (3.40). Here, \ln is real-valued for z and w real, resp., and the integration path in (3.86) belongs to $\mathbb{C}(\pi)$.

Proof: From the AΔE (3.4) with $a_\delta = a, a_{-\delta} = \pi$, we deduce that (3.85) need only be proved for $\lambda, \mu \in [-1/2, 1/2]$. Taking from now on $a \in (0, \pi/4]$ (say), we fix λ and μ in this interval and z in the strip $|\operatorname{Im} z| < \pi/2$. Then we may use (3.1) to write

$$d_a(\lambda, \mu; z) = -i \int_0^\infty \frac{dy}{y} \left(\frac{(\lambda - \mu)}{\pi y} - \frac{\operatorname{sha}(\lambda - \mu)y}{\operatorname{sh} ay} \frac{\cos(2yz + ia(\lambda + \mu))}{\operatorname{sh} \pi y} \right), \quad (3.87)$$

$$D_a(z) = \int_0^\infty \frac{dy}{y^2} \left(\frac{ay}{\operatorname{sh} ay} \frac{\sin 2yz}{2 \operatorname{sh} \pi y} - \frac{z}{\pi} \right). \quad (3.88)$$

From straightforward estimates one sees that these representations entail the limits

$$\lim_{a \downarrow 0} d_a(\lambda, \mu; z) = -i(\lambda - \mu) \int_0^\infty \frac{dy}{y} \left(\frac{1}{\pi y} - \frac{\cos 2yz}{\operatorname{sh} \pi y} \right), \quad (3.89)$$

$$\lim_{a \downarrow 0} D_a(z) = \int_0^\infty \frac{dy}{y^2} \left(\frac{\sin 2yz}{2 \operatorname{sh} \pi y} - \frac{z}{\pi} \right), \quad (3.90)$$

and boundedness for $(a, z) \in (0, \pi/4] \times K$, with K a compact subset of $|\operatorname{Im} z| < \pi/2$.

Invoking now Vitali's theorem and recalling the identity (3.21), it follows that (3.85) and (3.86) hold true uniformly on compacts in $|\operatorname{Im} z| < \pi/2$. Next, we exploit Theorem II.4 to obtain uniform convergence on compacts in the right half plane (2.73). To this end we need only observe that the AΔEs with step size a obeyed by $\partial_z^2 d_a$ and $\partial_z^3 D_a$ satisfy all of the assumptions of Theorem II.4, cf. the proof of Prop. III.1. Similarly, we infer uniform convergence on compacts of the left half plane. Since any compact in $\mathbb{C}(\pi)$ can be written as a union of three compacts in the strip $|\operatorname{Im} z| < \pi/2$ and in the left and right half planes, the proposition now follows. \square

We point out that (3.85) amounts to

$$\lim_{a \downarrow 0} \frac{G(\pi, a; z + i\lambda a)}{G(\pi, a; z + i\mu a)} = \exp((\lambda - \mu) \ln(2 \operatorname{ch} z)), \quad \lambda, \mu \in \mathbb{R}, \quad (3.91)$$

uniformly on compacts in $\mathbb{C}(\pi)$. Observe that the rhs is not meromorphic, unless $\lambda - \mu \in \mathbb{Z}$. The emergence of branch cuts can be understood from the coalescence of zeros and poles taking place for $a \rightarrow 0$, cf. Prop. III.3.

B. The elliptic case

In this subsection we are concerned with a function that is a minimal solution to three AΔEs generalizing the hyperbolic AΔEs (3.4) and (3.5). We study this function along the same lines as in Subsection III A. Our starting point is the infinite series

$$\sum_{n=1}^{\infty} \frac{\sin 2nrz}{2n \operatorname{sh} nra_+ \operatorname{sh} nra_-} \equiv g(r, a_+, a_-; z), \quad (3.92)$$

where we take at first $r, a_\delta \in (0, \infty)$, $\delta = +, -$. Clearly, this series converges absolutely and uniformly for z in an arbitrary compact of the strip S (3.2), so it defines a function g that is analytic in S . As before, it is convenient to suppress the dependence on the parameters whenever this causes no confusion. With this convention, our goal is to study the function $G(z)$ (3.3).

To this end we introduce the ‘‘right-hand side function’’

$$R(r, a; z) \equiv -2ire^{-ar/2} \prod_{k=1}^{\infty} (1 - e^{-2kar})^2 \cdot e^{irz} s(r, a; z + ia/2). \quad (3.93)$$

Using the definition (2.89) of s and the product representation (2.84) of the σ -function, one easily verifies that R can be rewritten

$$R(r, a; z) = \prod_{k=1}^{\infty} (1 - \exp(2irz - (2k-1)ar))(z \rightarrow -z), \quad (3.94)$$

where the infinite product converges absolutely and uniformly on compacts. From this one readily obtains the representation

$$R(r, a; z) = \exp\left(-\sum_{n=1}^{\infty} \frac{\cos 2nrz}{n \operatorname{sh} nra}\right), \quad |\operatorname{Im} z| < a/2. \quad (3.95)$$

(Use the power series for $\ln(1-x)$ to verify this; cf. also (2.93).)

In the sequel it is convenient to employ the abbreviations

$$q_{\delta} \equiv \exp(-a_{\delta} r), \quad (3.96)$$

$$c_{\delta} \equiv -2ir q_{\delta}^{1/2} \prod_{k=1}^{\infty} (1 - q_{\delta}^{2k})^2, \quad (3.97)$$

$$s_{\delta}(z) \equiv s(r, a_{\delta}; z), \quad (3.98)$$

$$R_{\delta}(z) \equiv R(r, a_{\delta}; z) = c_{\delta} e^{irz} s_{\delta}(z + ia_{\delta}/2), \quad (3.99)$$

where $\delta = +, -$. We are now prepared for the following proposition.

Proposition III.8 (defining AΔEs): *With (3.4) replaced by*

$$\frac{G(z + ia_{\delta}/2)}{G(z - ia_{\delta}/2)} = R_{-\delta}(z), \quad \delta = +, -, \quad (3.100)$$

and (3.5) by

$$\frac{G(z + i(a_{+} - a_{-})/2)}{G(z - i(a_{+} - a_{-})/2)} = \prod_{k=1}^{\infty} \left(\frac{1 - q_{-}^{2k}}{1 - q_{+}^{2k}} \right)^2 \cdot \frac{s_{-}(z)}{s_{+}(z)}, \quad (3.101)$$

the assertions of Prop. III.1 hold true.

Proof: In view of (3.99) and (3.95), Theorem II.5 may be invoked to solve the additive form of (3.100). Specifically, we may take

$$\phi(z) \equiv -\sum_{n \in \mathbb{Z}^*} \frac{e^{2inrz}}{2n \operatorname{sh} nra_{-\delta}}, \quad (3.102)$$

$s = a_{-}/2$ and $a = a_{\delta}$. The solution given by (2.106) is then equal to $ig(r, a_{+}, a_{-}; z)$ [cf. (3.92)], and so (3.100) follows.

Next, we use (3.20) and the AΔEs (3.100) to conclude that (3.101) amounts to the identity

$$\frac{s_{-}(z)}{s_{+}(z)} = \prod_{k=1}^{\infty} \left(\frac{1 - q_{+}^{2k}}{1 - q_{-}^{2k}} \right)^2 \cdot \frac{R_{-}(z - ia_{-}/2)}{R_{+}(z - ia_{+}/2)}. \quad (3.103)$$

This identity can be deduced from (3.96)–(3.99), so the proposition follows. \square

Proposition III.9 (automorphy properties): *The function G is periodic with primitive period π/r . It obeys the multiplication formula (3.25) and the period doubling formula*

$$G(2r, a_{+}, a_{-}; z) = G(r, a_{+}, a_{-}; z) G(r, a_{+}, a_{-}; z - \pi/2r). \quad (3.104)$$

Moreover, it satisfies (3.22), (3.23), the scaling relation

$$G(\lambda^{-1}r, \lambda a_{+}, \lambda a_{-}; \lambda z) = G(r, a_{+}, a_{-}; z), \quad \lambda \in (0, \infty), \quad (3.105)$$

and the duplication formula

$$G(r, a_+, a_-; 2z) = \prod_{l, m = +, -} G(r, a_+, a_-; z - i(la_+ + ma_-)/4) \\ \times G(r, a_+, a_-; z - i(la_+ + ma_-)/4 - \pi/2r). \quad (3.106)$$

Proof: These features follow from the series representation (3.92) in the same way as in the hyperbolic case. (Combine (3.25), (3.104) and (3.105) to check (3.106).) \square

Proposition III.10 (zeros, poles, residues): *The zeros and poles of $G(z)$ are given by*

$$z_{jkl}^+ \equiv j\pi/r + z_{kl}^+, \quad j \in \mathbb{Z}, \quad k, l \in \mathbb{N} \quad (\text{zeros}), \quad (3.107)$$

$$z_{jkl}^- \equiv -z_{jkl}^+, \quad j \in \mathbb{Z}, \quad k, l \in \mathbb{N} \quad (\text{poles}), \quad (3.108)$$

with z_{kl}^+ defined by (3.26). The multiplicities of the poles $z_{jk_0l_0}^-$ and zeros $z_{jk_0l_0}^+$, $j \in \mathbb{Z}$, are equal to the number of distinct pairs $(k, l) \in \mathbb{N}^2$ such that $z_{kl}^+ = z_{k_0l_0}^+$. The poles at z_{j00}^- , $j \in \mathbb{Z}$, are simple and have residue

$$r_{00} = i \left(2r \prod_{n=1}^{\infty} (1 - q_-^{2n})(1 - q_+^{2n}) \right)^{-1}. \quad (3.109)$$

Whenever

$$e_{kl} \equiv \prod_{m=1}^k i s_-(i m a_+) \prod_{n=1}^l i s_+(i n a_-) \quad (3.110)$$

is non-zero, the poles at z_{jkl}^- , $j \in \mathbb{Z}$, are simple and have residue

$$r_{kl} = (-)^{kl} \left(\frac{1}{2r} \right)^{k+l} q_-^{(l^2+l)(k+1/2)} q_+^{(k^2+k)(l+1/2)} \prod_{n=1}^{\infty} (1 - q_-^{2n})^{-2k} (1 - q_+^{2n})^{-2l} \cdot r_{00}/e_{kl}. \quad (3.111)$$

Conversely, if z_{jkl}^- is a simple pole, then $e_{kl} \neq 0$.

Proof: We proceed along the same lines as in the proof of Prop. III.3. Here, (3.31) holds true with (3.32) replaced by

$$P_M(z) \equiv \left(\prod_{m=1}^M R_-(z - i a_+(m - 1/2)) \right)^{-1} \quad (3.112)$$

and then the poles of $P_M(z)$ are located at $j\pi/r + z_{ml}$, with $j \in \mathbb{Z}$ and z_{ml} given by (3.33). By periodicity we may restrict attention to poles and zeros on the imaginary axis. In view of (3.22) the first two assertions then follow just as in the hyperbolic case.

Turning to the third one, we now get

$$G(z - i(a_+ + a_-)/2) = (c_- \exp[ir(z - i a_-/2)] s_-(z))^{-1} G(z + i(a_+ - a_-)/2) \quad (3.113)$$

so that [cf. (3.96) and (3.97)]

$$r_{00} = \frac{i}{2r} \prod_{n=1}^{\infty} (1 - q_-^{2n})^{-2} G(i(a_+ - a_-)/2). \quad (3.114)$$

Using symmetry in a_+, a_- , we deduce

$$G(i(a_+ - a_-)/2) = \prod_{n=1}^{\infty} \frac{(1 - q_-^{2n})}{(1 - q_+^{2n})} \quad (3.115)$$

and so (3.109) follows. (Note that (3.115) can also be derived from (3.101).)

Finally, from the AΔEs (3.100) we calculate

$$\begin{aligned} G(z + z_{kl}^-) = & (-)^{kl} \left(c_-^k c_+^l \exp \left(\frac{ra_-}{2} [(l^2 + l)(2k + 1) + k] + \frac{ra_+}{2} [(k^2 + k)(2l + 1) + l] \right) \right. \\ & \cdot \exp(irz[k + l + 2kl]) \prod_{m=1}^k s_-(z - ima_+) \prod_{n=1}^l s_+(z - ina_-) \left. \right)^{-1} \cdot G(z + z_{00}^-). \end{aligned} \quad (3.116)$$

Using (3.96) and (3.97), the remaining assertions readily follow from this. \square

At the elliptic level the choice $a_+ = a_-$ does not appear to yield extra information, as compared to the general case. But since G is π/r -periodic, there is no analog of Prop. III.4, and so we do not need additional information on this special case.

Next, we turn to an analog of Prop. III.5.

Proposition III.11 (meromorphic continuation): *The function G admits the representation*

$$G(r, a_+, a_-; z) = \prod_{m,n=1}^{\infty} \frac{1 - q_+^{2m-1} q_-^{2n-1} e^{-2irz}}{1 - q_+^{2m-1} q_-^{2n-1} e^{2irz}}, \quad q_{\delta} \equiv \exp(-a_{\delta} r). \quad (3.117)$$

It can be analytically continued to a function that is meromorphic in r, a_+, a_- and z , provided $a_+ r$ and $a_- r$ stay in the right half plane. Fixing r, a_+, a_- with $\operatorname{Re}(a_+ r)$ and $\operatorname{Re}(a_- r)$ positive, one obtains a meromorphic function whose zeros and poles are located at (3.107) and (3.108), resp.

Proof: It suffices to prove (3.117), since the remaining assertions are clear from this formula. To this end we observe that the numerator infinite product is the downward iteration solution to both of the AΔEs

$$\frac{F(z + ia_{\delta}/2)}{F(z - ia_{\delta}/2)} = R^{(-)}(a_{-\delta}; z), \quad \delta = +, -, \quad (3.118)$$

with

$$R^{(-)}(a; z) \equiv \prod_{k=1}^{\infty} (1 - e^{-(2k-1)ar} e^{-2irz}). \quad (3.119)$$

Similarly, the denominator infinite product is the upward iteration solution to

$$\frac{F(z + ia_{\delta}/2)}{F(z - ia_{\delta}/2)} = R^{(+)}(a_{-\delta}; z), \quad \delta = +, -, \quad (3.120)$$

with

$$R^{(+)}(a; z) \equiv R^{(-)}(a; -z); \quad (3.121)$$

cf. (2.1)–(2.5). But we have

$$R^{(+)}(a_{\delta}; z)R^{(-)}(a_{\delta}; z) = R_{\delta}(z), \quad (3.122)$$

cf. (3.94), so the rhs \tilde{G} of (3.117) solves the AΔE (3.100). Since both solutions G and \tilde{G} are π/r -periodic, have no zeros and poles in the strip $|\operatorname{Im} z| \leq a_{\delta}/2$, and satisfy $G(0) = \tilde{G}(0) = 1$, we deduce $G = \tilde{G}$. \square

We continue by detailing the relation of the elliptic G -function to the hyperbolic G -function. This relation is the first instance of a general type of limiting transition between meromorphic functions that will reappear several times. Therefore, it is convenient to introduce a term referring to the type of limit involved.

To this end, assume $f_p(z)$ is a family of meromorphic functions parametrized by $p \in \mathbb{C}^N$. We shall say that $f_p(z)$ converges mero-uniformly to a meromorphic function $f(z)$ as $p \rightarrow p_0$ whenever one has $f_p(z) \rightarrow f(z)$ uniformly on compacts not containing poles of $f(z)$, and $1/f_p(z) \rightarrow 1/f(z)$ uniformly on compacts not containing zeros of $f(z)$. (Equivalently, viewing meromorphic functions as holomorphic functions from \mathbb{C} to the Riemann sphere \mathbb{P}^1 , one has $f_p \rightarrow f$ mero-uniformly as $p \rightarrow p_0$ iff the convergence is \mathbb{P}^1 -uniform on arbitrary \mathbb{C} -compacts.)

Defining the renormalized function

$$G_{\text{ren}}(r, a_+, a_-; z) = G(r, a_+, a_-; z) \exp\left(\frac{\pi^2 z}{6ira_+ a_-}\right) \quad (3.123)$$

we are now prepared for the next proposition.

Proposition III.12 (relation to hyperbolic G -function): Fixing $a_+, a_- > 0$, one has

$$\lim_{r \downarrow 0} G_{\text{ren}}(r, a_+, a_-; z) = G(a_+, a_-; z), \quad (3.124)$$

where the limit is mero-uniform.

Proof: Writing $G_{\text{ren}} = \exp(ig_{\text{ren}})$, we obtain

$$g_{\text{ren}}(r, a_+, a_-; z) = r \sum_{n=1}^{\infty} \frac{1}{nr} \left(\frac{\sin 2nrz}{2 \operatorname{sh} nra_+ \operatorname{sh} nra_-} - \frac{z}{nra_+ a_-} \right), \quad z \in S; \quad (3.125)$$

cf. (3.92). Comparing to (3.1), a routine dominated convergence argument now yields

$$\lim_{r \downarrow 0} g_{\text{ren}}(r, a_+, a_-; z) = g(a_+, a_-; z), \quad z \in S, \quad (3.126)$$

uniformly on S -compacts.

Next, we note that G_{ren} satisfies the AΔE

$$\frac{G(z + ia_+/2)}{G(z - ia_+/2)} = R_{-, \text{ren}}(z) \quad (3.127)$$

with

$$R_{-, \text{ren}}(z) \equiv \exp\left(\frac{\pi^2}{6ra_-}\right) R_-(z). \quad (3.128)$$

In view of (3.126) this entails that for $|\operatorname{Im} z| \leq a_-/2$ we have

$$\lim_{r \downarrow 0} R_{-, \text{ren}}(z) = \frac{G(a_+, a_-; z + ia_+/2)}{G(a_+, a_-; z - ia_+/2)} = 2 \operatorname{ch} \frac{\pi z}{a_-}, \quad (3.129)$$

where we used (3.4). Recalling (3.99) and the limit (2.92), we deduce

$$\lim_{r \downarrow 0} \exp\left(\frac{\pi^2}{6ra_-}\right)(ic_-) = \frac{2\pi}{a_-}. \quad (3.130)$$

Using then (2.92) once more, one sees that (3.129) holds uniformly on \mathbb{C} -compacts. Therefore, one may exploit the AΔE (3.127) and uniform convergence of G_{ren} to G on S -compacts to obtain uniform convergence on \mathbb{C} -compacts that do not contain the poles $z_{jk}^-, j, k \in \mathbb{N}$, of G . Moreover, (3.126) entails uniform convergence of $1/G_{\text{ren}}$ to $1/G$ on S -compacts, so one can also use (3.127) and (3.129) to infer $1/G_{\text{ren}} \rightarrow 1/G$ uniformly on compacts not containing the zeros z_{jk}^+ . \square

As a corollary of the proof we obtain the limit

$$\lim_{r \downarrow 0} r \exp\left(\frac{\pi^2}{6ra}\right) \prod_{n=1}^{\infty} (1 - e^{-2nar})^2 = \frac{\pi}{a}, \quad a > 0; \quad (3.131)$$

cf. (3.130) and (3.97). Equivalently, this can be written

$$\lim_{r \downarrow 0} \left(\sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{e^{-nra}}{\text{sh} nra} - \frac{1}{nra} \right) - \ln r \right) = \ln \frac{a}{\pi}, \quad a > 0. \quad (3.132)$$

The last proposition of this subsection is the analog of Prop. III.7 in the previous one. To state it, we introduce the cut plane

$$\mathbb{C}(r, d) \equiv \mathbb{C} \setminus \{\pm i[d/2, \infty) + k\pi/r \mid k \in \mathbb{Z}\}, \quad r, d > 0, \quad (3.133)$$

and define a branch $g(r, A, a; z)$ of $-i \ln G$ in $\mathbb{C}(r, A+a)$ via (3.93) for $|\text{Im} z| < (A+a)/2$. Then we set

$$d_a(r, A, \lambda, \mu; z) \equiv g(r, A, a; z + i\lambda a) - g(r, A, a; z + i\mu a), \quad z \in \mathbb{C}(r, A+a), \quad \lambda, \mu \in \mathbb{R}, \quad (3.134)$$

$$D_a(r, A; z) \equiv a g(r, A, a; z), \quad z \in \mathbb{C}(r, A+a) \quad (3.135)$$

(This should be compared to (3.83) and (3.84).)

Proposition III.13 (zero step size limits): *One has*

$$\lim_{a \downarrow 0} d_a(r, A, \lambda, \mu; z) = -i(\lambda - \mu) \ln R(r, A; z), \quad \lambda, \mu \in \mathbb{R}, \quad (3.136)$$

$$\lim_{a \downarrow 0} D_a(r, A; z) = - \int_0^z dw \ln R(r, A; w), \quad (3.137)$$

uniformly on compact subsets of the cut plane $\mathbb{C}(r, A)$ (3.133). Here, \ln is real-valued for z and w real, resp., and the integration path in (3.137) belongs to $\mathbb{C}(r, A)$.

Proof: This follows in the same way as Prop. III.7, with (3.93), (the logarithm of) (3.95) and Theorem II.7 playing the role of (3.1), (3.21) and Theorem II.4, resp. (Since the limits are π/r -periodic in the strip $|\text{Im} z| < A/2$, one need only handle compacts in \mathcal{R}_r (2.113).) \square

In terms of G , (3.136) reads

$$\lim_{a \downarrow 0} \frac{G(r, A, a; z + i\lambda a)}{G(r, A, a; z + i\mu a)} = \exp((\lambda - \mu) \ln R(r, A; z)), \quad \lambda, \mu \in \mathbb{R}, \quad (3.138)$$

uniformly on compacts in $\mathbb{C}(r, A)$. Once more, the branch cuts arise from coalescence of zeros and poles, cf. Prop. III.10.

C. The trigonometric case

The trigonometric case is most easily understood by viewing it as a limiting case of the elliptic case. In view of (2.90), this should involve sending one of a_+, a_- to ∞ . We shall fix $a_+ \equiv a \in (0, \infty)$ and let $a_- \equiv A$ go to ∞ . To get finite limits, we clearly should shift z in an A -dependent way. We take $z \rightarrow z - iA/2$, and thus wind up with

$$G(r, a; z) \equiv \lim_{A \rightarrow \infty} G(r, a, A; z - iA/2). \quad (3.139)$$

From the product representation (3.117) it is immediate that this limit exists mero-uniformly, yielding

$$G(r, a; z) = \prod_{m=1}^{\infty} (1 - q^{2m-1} e^{2irz})^{-1}, \quad q \equiv e^{-ar}. \quad (3.140)$$

For $\text{Im} z > -a/2$ we can also evaluate the limit (3.139) by using (3.92); this yields the series representation

$$G(r, a; z) = \exp \left(\sum_{n=1}^{\infty} \frac{e^{2inrz}}{2n \text{sh} nr a} \right), \quad \text{Im} z > -a/2. \quad (3.141)$$

We continue by studying the trigonometric G -function just defined.

Proposition III.14 (defining AΔE): *The function $G(r, a; z)$ is the upward iteration solution to the AΔE*

$$\frac{G(z + ia/2)}{G(z - ia/2)} = 1 - e^{2irz}. \quad (3.142)$$

Proof: This is clear from the product representation (3.140) [recall (2.1)–(2.4)]. \square

Notice that the AΔE (3.142) is not regular. However, a shift $z \rightarrow z + ia/2$ (say) gives rise to a regular AΔE. Indeed, the function

$$\phi(z) = \ln(1 - \exp(2ir(z + ia/2))) = - \sum_{n=1}^{\infty} n^{-1} q^n e^{2inrz} \quad (3.143)$$

satisfies the assumptions of Theorem II.4, and $G(r, a; z + ia/2)$ is a minimal solution to the associated multiplicative AΔE. [Compare the logarithm of (3.141) with (2.106) to see this.] Observe also that (3.142) agrees with the $A \rightarrow \infty$ limit of the elliptic AΔE

$$\frac{G(r, a, A; z - iA/2 + ia/2)}{G(r, a, A; z - iA/2 - ia/2)} = -2ir \prod_{n=1}^{\infty} (1 - e^{-2nAr})^2 \cdot e^{irz} s(r, A; z), \quad (3.144)$$

cf. (3.100), (3.96)–(3.99), (3.139) and (2.90).

Proposition III.15 (automorphy properties): *The function G is periodic with primitive period π/r . It obeys the multiplication formula*

$$G\left(r, \frac{a}{M}; z\right) = \prod_{j=1}^M G\left(r, a; z + \frac{ia}{2M}(M+1-2j)\right), \quad (3.145)$$

the period doubling formula

$$G(2r, a; z) = G(r, a; z)G(r, a; z - \pi/2r), \quad (3.146)$$

the scaling relation

$$G(\lambda^{-1}r, \lambda a; \lambda z) = G(r, a; z), \quad \lambda \in (0, \infty), \quad (3.147)$$

and the duplication formula

$$G(r, a; 2z) = \prod_{\sigma=+, -} G(r, a; z - i\sigma a/4)G(r, a; z - i\sigma a/4 - \pi/2r). \quad (3.148)$$

Proof: These properties follow from the series representation (3.141) in the same way as in the two previous cases. \square

Proposition III.16 (zeros, poles, residues): *The function $G(z)$ has no zeros and simple poles given by*

$$z_{jk} \equiv j\pi/r - ia(k + 1/2), \quad j \in \mathbb{Z}, \quad k \in \mathbb{N} \quad (\text{poles}). \quad (3.149)$$

The residues at the poles $z_{j0}, j \in \mathbb{Z}$, are given by

$$r_0 = i \left(2r \prod_{n=1}^{\infty} (1 - q^{2n}) \right)^{-1} = \frac{i}{2r} G(ia/2), \quad (3.150)$$

and the residues at the remaining poles $z_{jk}, j \in \mathbb{Z}, k \in \mathbb{N}^*$, are given by

$$r_k = r_0 / \prod_{m=1}^k (1 - q^{-2m}). \quad (3.151)$$

Proof: The first assertion is immediate from (3.140). The residues (3.150) follow either from (3.109) by taking a limit, or directly from (3.140). Using

$$G(z + z_{0k}) = \prod_{m=1}^k (1 - q^{-2m} e^{2irz})^{-1} G(z + z_{00}), \quad (3.152)$$

the residues at the remaining poles can now be obtained, yielding (3.151). \square

Proposition III.17 (asymptotics): *The function G satisfies the bound*

$$G(r, a; z) = 1 + O(\exp(-2r\operatorname{Im}z)), \quad \operatorname{Im}z \rightarrow \infty, \quad (3.153)$$

uniformly for $\operatorname{Re}z \in \mathbb{R}$.

Proof: This estimate readily follows from the series representation (3.141). \square

Proposition III.18 (meromorphic continuation): *The function G can be analytically continued to a function that is meromorphic in r, a and z , provided ar stays in the right half plane. Fixing r, a with $\operatorname{Re}(ar) > 0$, one obtains a meromorphic function without zeros and with simple poles located at (3.149).*

Proof: This can be read off from the product representation (3.140). \square

The propositions derived thus far have elliptic and/or hyperbolic analogs. In the previous two cases, however, the G -function satisfies $G(z)G(-z) = 1$, a relation that does not hold in the trigonometric case. Instead, we have the following result.

Proposition III.19 (functional equation): *The trigonometric G -function satisfies*

$$G(r, a; z)G(r, a; -z) = R(r, a; z)^{-1}, \quad (3.154)$$

where the rhs is given by (3.93).

Proof: This is obvious from the series representations (3.141) and (3.95). \square

We point out that this functional equation may be seen as a footprint left by the second A Δ E satisfied by the elliptic G -function: Taking $a \rightarrow a_+$, the rhs can be written $R_+(z)^{-1}$, so (3.154) can be deduced from (3.100) with $\delta = -$ and the limit (3.139).

Next, we introduce the function

$$T(r; z) \equiv \frac{G(r, 1; 0)}{G(r, 1; -z)} \exp\left(\frac{rz^2}{2} + iz \ln(2r) - \frac{1}{2} \ln \pi\right), \quad \operatorname{Re} r > 0. \quad (3.155)$$

This renormalized version of $G(r, a; z)$ satisfies the A Δ E

$$\frac{T(r; z + i/2)}{T(r; z - i/2)} = \frac{i \sin rz}{r} \quad (3.156)$$

and functional equation

$$T(r; z)T(r; -z) = \pi^{-1} \exp(rz^2 + irz) \frac{s(r, 1; z + i/2)}{s(r, 1; i/2)}. \quad (3.157)$$

Taking $r \downarrow 0$, the right-hand sides of (3.156) and (3.157) obviously converge to iz and $\pi^{-1} \operatorname{ch} \pi z$ [recall (2.92)], resp., in accordance with the next proposition.

Proposition III.20 (relation to gamma function): *One has*

$$\lim_{r \downarrow 0} T(r; z) = 1/\Gamma(iz + 1/2) \quad (3.158)$$

uniformly for z in \mathbb{C} -compacts.

Proof: We begin by noting that it suffices to show that (3.158) holds uniformly on compacts of the lower half plane (LHP). (Indeed, from (3.156) we have

$$T(r; z + ik) = \frac{i}{r} \sin r(z + i(k - 1/2)) \cdots \frac{i}{r} \sin r(z + i/2) T(r; z), \quad (3.159)$$

so if (3.158) holds uniformly on LHP-compacts, then the rhs of (3.159) converges in the same sense to

$$(iz - k + 1/2) \cdots (iz - 1/2) \frac{1}{\Gamma(iz + 1/2)} = \frac{1}{\Gamma(iz + ik + 1/2)}. \quad (3.160)$$

Hence, (3.158) follows for compacts of $\operatorname{Im} z \leq k$. To this end we use the formula

$$e(z) = e(0) + ze'(0) + \int_0^z dw \int_0^w ds e''(s) \quad (3.161)$$

to rewrite the logarithms of $T(r; z)$ and $1/\Gamma(iz + 1/2)$. This yields

$$T(r; z) = \exp\left(-\frac{1}{2} \ln \pi + izK(r) + \int_0^z dw \int_0^w ds h(r; s)\right) \quad (3.162)$$

with

$$h(r; z) \equiv 2r \sum_{n=1}^{\infty} \frac{nre^{-2inrz}}{\operatorname{sh} nr} + r, \quad \operatorname{Im} z < 1/2, \quad (3.163)$$

$$K(r) \equiv \ln(2r) + \sum_{n=1}^{\infty} \frac{r}{\operatorname{sh} nr} \quad (3.164)$$

[cf. (3.155) and (3.141)] and

$$\frac{1}{\Gamma(i z + 1/2)} = \exp\left(-\frac{1}{2} \ln \pi - i z \psi\left(\frac{1}{2}\right) + \int_0^z dw \int_0^w ds h(s)\right) \quad (3.165)$$

with

$$h(z) \equiv 2 \int_0^{\infty} dy \frac{y e^{-2iyz}}{\operatorname{sh} y}, \quad \operatorname{Im} z < 1/2, \quad (3.166)$$

cf. (A37), (A12), and (A33), (A34).

Comparing (3.163) and (3.166), we deduce

$$\lim_{r \downarrow 0} h(r; z) = h(z) \quad (3.167)$$

uniformly on LHP-compacts. Comparing then (3.162) with (3.165), we see that it remains to show

$$\lim_{r \downarrow 0} K(r) = -\psi\left(\frac{1}{2}\right). \quad (3.168)$$

To prove this, we use the AΔEs (3.156) and (A24) to write

$$\frac{T(r; -i)}{T(r; 0)} \frac{\Gamma(3/2)}{\Gamma(1/2)} = \frac{r}{2 \operatorname{sh}(r/2)}. \quad (3.169)$$

Due to (3.162) and (3.165), the lhs can be rewritten

$$\exp\left(K(r) + \psi\left(\frac{1}{2}\right) + \int_0^{-i} dw \int_0^w ds [h(r; s) - h(s)]\right), \quad (3.170)$$

and since the integral converges to 0 for $r \downarrow 0$ we now obtain (3.168). Therefore, the proof of the proposition is complete. \square

Comparing the AΔEs (3.156) and (3.70), we deduce that the quotient

$$Q(r; z) \equiv T(r; z)/H(ir/\pi; z), \quad \operatorname{Re} r > 0, \quad (3.171)$$

of the trigonometric and hyperbolic functions is i -periodic. Moreover, comparing poles and zeros of T and H , we deduce that Q is entire in z and has simple zeros at

$$z = -k\pi/r + i(l + 1/2), \quad k \in \mathbb{N}^*, \quad l \in \mathbb{Z}. \quad (3.172)$$

Furthermore, recalling Prop. III.6, we infer

$$\lim_{r \downarrow 0} Q(r; z) = 1 \quad (\text{uniformly on compacts}). \quad (3.173)$$

Our last proposition concerns two zero step size limits that may be tied in with (3.136) and (3.137) via (3.139). We set

$$\mathbb{C}_-(r, d) \equiv \mathbb{C} \setminus \{-i[d/2, \infty) + k\pi/r | k \in \mathbb{Z}\}, \quad r > 0, d \geq 0, \quad (3.174)$$

and define a branch $g(r, a; z)$ of $-i \ln G$ in $\mathbb{C}_-(r, a)$ by requiring

$$g(r, a; z) \equiv -i \sum_{n=1}^{\infty} \frac{e^{2inrz}}{2n \operatorname{sh} nra}, \quad \operatorname{Im} z > -a/2, \quad (3.175)$$

cf. (3.141). Now we put

$$d_a(r, \lambda, \mu; z) \equiv g(r, a; z + i\lambda a) - g(r, a; z + i\mu a), \quad z \in \mathbb{C}_-(r, a), \quad \lambda, \mu \in \mathbb{R}, \quad (3.176)$$

$$D_a(r; z) \equiv a g(r, a; z), \quad z \in \mathbb{C}_-(r, a). \quad (3.177)$$

(Compare this to (3.133)–(3.135).)

Proposition III.21 (zero step size limits): *One has*

$$\lim_{a \downarrow 0} d_a(r, \lambda, \mu; z) = -i(\lambda - \mu) \ln(1 - e^{2irz}), \quad \lambda, \mu \in \mathbb{R}, \quad (3.178)$$

$$\lim_{a \downarrow 0} D_a(r; z) = - \int_{i\infty}^z dw \ln(1 - e^{2irw}), \quad (3.179)$$

uniformly on compact subsets of the cut plane $\mathbb{C}_-(r, 0)$ (3.174). Here, \ln is real valued for $z, w \in i(0, \infty)$, and the integration path in (3.179) belongs to $\mathbb{C}_-(r, 0)$.

Proof: From (3.175) it readily follows that the proposition is valid when the cut plane $\mathbb{C}_-(r, 0)$ is replaced by its upper half plane subset. Applying Theorem II.7 to the functions $f_a(z) \equiv d_a(z + i)$ and $f_a(z) \equiv D_a(z + i)$ (which satisfy the hypotheses of that theorem for a_0 small enough), one obtains validity for all of the cut plane. \square

Translated to G , the limit (3.178) becomes

$$\lim_{a \downarrow 0} \frac{G(r, a; z + i\lambda a)}{G(r, a; z + i\mu a)} = \exp((\lambda - \mu) \ln(1 - e^{2irz})), \quad \lambda, \mu \in \mathbb{R}, \quad (3.180)$$

uniformly on compact subsets of the cut plane $\mathbb{C}_-(r, 0)$. Just as in the previous two cases [cf. (3.91) and (3.138)], this formula is evident from the defining AΔE when $\lambda - \mu$ is an integer. For $\lambda - \mu \notin \mathbb{Z}$, the branch cuts in the lower half plane arise from the coalescence of poles and zeros that can be read off from (3.149).

We conclude this subsection by detailing the relation of the trigonometric G -function $G(r, a; z)$ to the \tilde{q} -gamma function $\Gamma_{\tilde{q}}(z)$. Recall the latter is given by (cf., e.g., Ref. 4, p. 16)

$$\Gamma_{\tilde{q}}(z) = (1 - \tilde{q})^{1-z} \prod_{n=1}^{\infty} \frac{(1 - \tilde{q}^n)}{(1 - \tilde{q}^{z+n-1})}. \quad (3.181)$$

Comparing this to the product formula (3.140) for G , we see that when we take

$$\tilde{q} \equiv q^2 = e^{-2ar} \quad (3.182)$$

we may write G as

$$G(r, a; az) = \Gamma_{\tilde{q}}(-iz + 1/2) (1 - \tilde{q})^{-iz - 1/2} \prod_{n=1}^{\infty} (1 - \tilde{q}^n)^{-1}. \quad (3.183)$$

From this we readily obtain [recall (3.155)]

$$T(r; z) = \frac{\Gamma_{\tilde{q}}(1/2)}{\Gamma_{\tilde{q}}(iz + 1/2)} \exp\left(-\frac{1}{2} \ln \pi + \frac{rz^2}{2} - iz \ln\left(\frac{1 - e^{-2r}}{2r}\right)\right). \quad (3.184)$$

Using these relations, some of the above results can be translated in terms of $\Gamma_{\tilde{q}}$, recovering results that have been obtained by several authors, cf. Ref. 4 and references given there.

IV. SCATTERING FUNCTIONS

A. The hyperbolic case

We present our results on the hyperbolic scattering function $u(a_+, a_-, b; z)$ in a form that anticipates our account of the elliptic case. First of all, we define u by

$$u(z) \equiv \frac{G(z - ib + i(a_+ + a_-)/2) G(z + ib - i(a_+ + a_-)/2)}{G(z - i(a_+ - a_-)/2) G(z + i(a_+ - a_-)/2)}, \quad (4.1)$$

where $G(z) = G(a_+, a_-; z)$ is the hyperbolic G -function from Subsection III A. In (4.1) and in many later formulas, the dependence on a_+ and a_- is suppressed. This should cause no confusion, since u —just like G —satisfies

$$u(a_+, a_-; z) = u(a_-, a_+; z), \quad (4.2)$$

cf. (3.23). Similarly, the automorphy properties (3.22) and (3.24) yield

$$u(-z) = 1/u(z), \quad (4.3)$$

$$u(\lambda a_+, \lambda a_-, \lambda b; \lambda z) = u(a_+, a_-, b; z), \quad \lambda \in (0, \infty). \quad (4.4)$$

By virtue of Prop. III.5 the u -function is meromorphic in a_+, a_-, b and z , provided the quotient a_+/a_- stays away from the negative real axis. As a rule, however, we restrict our considerations to parameters in the set

$$\mathcal{H} \equiv \{(a_+, a_-, b) | a_+, a_- > 0, b \in \mathbb{R}\}. \quad (4.5)$$

This choice corresponds to physical applications; in particular, it guarantees $|u(x)| = 1$ for real x .

Next, we observe that the AΔEs (3.4) entail that u solves the AΔEs

$$\frac{u(z + ia_{\delta}/2)}{u(z - ia_{\delta}/2)} = \frac{s_{-\delta}(z - ib + ia_{\delta}/2) s_{-\delta}(z + ib - ia_{\delta}/2)}{s_{-\delta}(z + ia_{\delta}/2) s_{-\delta}(z - ia_{\delta}/2)}, \quad (4.6)$$

where we have introduced

$$s_{\delta}(z) \equiv \frac{\text{sh}(\pi z/a_{\delta})}{\pi/a_{\delta}}, \quad \delta = +, -. \quad (4.7)$$

(This definition mimicks the elliptic definition (3.98), cf. (2.92).) Fixing $\delta \in \{+, -\}$, the AΔE (4.6) is regular unless the parameters (a_+, a_-, b) belong to the planes

$$a_{\delta} = 2na_{-\delta}, \quad n \in \mathbb{N}^*, \quad (4.8)$$

or

$$b = ka_{-\delta} + a_{\delta}/2, \quad k \in \mathbb{Z}. \quad (4.9)$$

These planes separate the region \mathcal{H} (4.5) into infinitely many connected components, one of which reads

$$\mathcal{R}_{\delta} \equiv \{(a_+, a_-, b) \in \mathcal{H} \mid a_{\delta} \in (0, 2a_{-\delta}), b \in (a_{\delta}/2, a_{-\delta} + a_{\delta}/2)\}. \quad (4.10)$$

Choosing parameters in \mathcal{R}_{δ} , the u -function may now be characterized as the unique minimal solution to the AΔE (4.6) that satisfies

$$u(0) = 1, \quad |u(x)| = 1, \quad x \in \mathbb{R}. \quad (4.11)$$

Indeed, the pole/zero properties of the G -function (cf. Prop. III.3) entail that u (4.1) is a regular solution to (4.6) if and only if $(a_+, a_-, b) \in \mathcal{R}_{\delta}$. Moreover, for all $(a_+, a_-, b) \in \mathcal{H}$ one has

$$u(z) = \exp\left(\pm \frac{i\pi}{a_+ a_-} (b - a_+)(b - a_-)\right) + O(\exp(\pm(\epsilon - 2\pi/a_m)z)), \quad \operatorname{Re} z \rightarrow \pm\infty, \quad (4.12)$$

uniformly for $\operatorname{Im} z$ in \mathbb{R} -compacts, cf. Prop. III.4. Therefore, u is indeed a minimal solution to (4.6) for parameters in \mathcal{R}_{δ} (4.10). From Theorem II.1 and (4.11) one now easily deduces the above uniqueness assertion.

It should be remarked at this point that the AΔE (4.6) does admit minimal solutions whenever the parameters do not belong to the planes (4.8) and (4.9). Indeed, this readily follows from Section II. More concretely, a minimal solution can be constructed by multiplying $u(z)$ by finitely many factors of the form $s_{\delta}(z-p)/s_{\delta}(z+p)$ that cancel the poles and zeros of $u(z)$ in the strip $|\operatorname{Im} z| < a_{\delta}/2$. (Observe that $u(z)$ has no poles and zeros for $|\operatorname{Im} z| = a_{\delta}/2$ unless (4.8) or (4.9) holds true.)

Since the rhs of (4.6) is $a_{-\delta}$ -periodic in b , the quotient $u(b+a_{-\delta}; z)/u(b; z)$ is ia_{δ} -periodic in z . Specifically, one obtains from (4.1) and (3.4)

$$\frac{u(b+a_{-\delta}; z)}{u(b; z)} = -\frac{s_{\delta}(z+ib)}{s_{\delta}(z-ib)}. \quad (4.13)$$

Therefore, iteration yields (taking $k_+, k_- \in \mathbb{Z}$)

$$\frac{u(b+k_+a_++k_-a_-; z)}{u(b; z)} = \prod_{\delta=+,-} \prod_{j_{\delta}=1}^{|k_{\delta}|} \frac{s_{-\delta}(z+i(k_{\delta}/|k_{\delta}|)(b-a_{\delta}/2)+ia_{\delta}(j_{\delta}-1/2))}{(z \rightarrow -z)}. \quad (4.14)$$

Next, we introduce the parameter subset

$$\mathcal{D} \equiv \{(a_+, a_-, b) \in \mathcal{H} \mid b = k_+a_+ + k_-a_-, k_+, k_- \in \mathbb{Z}\} \quad (4.15)$$

of \mathcal{H} (4.5). Since the numbers $k_+a_+ + k_-a_-$, $k_+, k_- \in \mathbb{Z}$, are dense in \mathbb{R} whenever $a_+/a_- \notin \mathbb{Q}$, the subset \mathcal{D} is dense in \mathcal{H} . Now from (4.1) we read off

$$u(a_+, a_-, a_+; z) = u(a_+, a_-, a_-; z) = 1 \quad (4.16)$$

and also, using (3.4),

$$u(a_+, a_-, 0; z) = -1. \quad (4.17)$$

Hence, (4.14) yields

$$u(a_+, a_-, k_+ a_+ + k_- a_-; z) = c_{k_+, k_-} \prod_{\delta=+, -} \prod_{j \delta=1}^{|k_\delta|} \frac{s_{-\delta}(z + i a_\delta(j_\delta - \theta(k_\delta)))}{(i \rightarrow -i)} \quad (4.18)$$

with

$$\theta(j) \equiv \begin{cases} 0, & j < 0, \\ 1, & j > 0, \end{cases} \quad (4.19)$$

and

$$c_{k,l} \equiv (-)^{k+l+1}, \quad k, l \in \mathbb{Z}. \quad (4.20)$$

In words, the u -function is an elementary function for parameters in the dense subset \mathcal{D} of \mathcal{H} . (Of course, whenever a_-/a_+ is a rational number, there exist infinitely many distinct pairs $(k, l) \in \mathbb{Z}^2$ for which the number $ka_+ + la_-$ is the same; this yields different representations for the same function.)

We continue by noting the symmetry property

$$u(b; z) = u(a_+ + a_- - b; z), \quad (4.21)$$

which can be read off from (4.1). Combining this with (4.14) (taking $k_+, k_- = 1$), we deduce

$$\frac{u(-b; z)}{u(b; z)} = \frac{s_+(z + ib)}{s_+(z - ib)} \frac{s_-(z + ib)}{s_-(z - ib)}. \quad (4.22)$$

Since this parameter transformation leaves \mathcal{D} (4.15) invariant, it does not give rise to additional elementary representations for u .

Next, we derive analogs of the multiplication formula (3.25). First, we use (4.1) to get

$$\begin{aligned} u\left(\frac{a_+}{M}, a_-, b; z\right) &= \prod_{j=1}^M \frac{G(z - ib + i(a_+/2) + i(a_-/2) + i(a_+/M)(1-j))}{G(z - i(a_+/2) + i(a_-/2) + i(a_+/M)(M-j))} \\ &\quad \times \frac{G(z + ib - i(a_+/2) - i(a_-/2) + i(a_+/M)(M-j))}{G(z + i(a_+/2) - i(a_-/2) + i(a_+/M)(1-j))}. \end{aligned} \quad (4.23)$$

with $G(z) = G(a_+, a_-; z)$. Rearranging and using (4.1) once more, we deduce

$$\begin{aligned} u\left(\frac{a_+}{M}, a_-, b; z\right) &= u(a_+, a_-, b; z) \prod_{k=1}^{M-1} u\left(a_+, a_-, b; z + ik \frac{a_+}{M}\right) \\ &\quad \times \frac{G(z + ik(a_+/M) - ib + i(a_-/2) - i(a_+/2))}{G(z + ik(a_+/M) - ib + i(a_-/2) + i(a_+/2))} \\ &\quad \times \frac{G(z + ik(a_+/M) - i(a_-/2) + i(a_+/2))}{G(z + ik(a_+/M) - i(a_-/2) - i(a_+/2))}. \end{aligned} \quad (4.24)$$

This can be simplified by using the AΔE (3.4), which yields

$$u\left(\frac{a_+}{M}, a_-, b; z\right) = (-)^{M-1} \prod_{k=0}^{M-1} u\left(a_+, a_-, b; z + ik \frac{a_+}{M}\right) \prod_{j=1}^{M-1} \frac{s_-(z + ija_+/M)}{s_-(z - ib + ija_+/M)}. \quad (4.25)$$

Equivalently, we may also rearrange (4.23) to get

$$u\left(\frac{a_+}{M}, a_-, b; z\right) = (-)^{M-1} \prod_{k=0}^{M-1} u\left(a_+, a_-, b; z - ik \frac{a_+}{M}\right) \prod_{j=1}^{M-1} \frac{s_-(z + ib - ija_+/M)}{s_-(z - ija_+/M)}. \quad (4.26)$$

Substituting $a_- \rightarrow a_-/N$ in the formulas (4.25) and (4.26), and using first (4.2) and then one of these formulas again, one obtains four representations for $u(a_+/M, a_-/N, b; z)$ in terms of $u(a_+, a_-, b; z)$ and sh-quotients.

The choices $b = a_+/2$ or $b = a_-/2$ yield the sine-Gordon soliton-soliton S -matrix. Taking $b = a_+/2$, it follows from (4.18) that there exists a dense set of a_- -choices yielding an elementary u . Specifically, choosing $a_- = a_+(1 + 2j)/2l$ with $j \in \mathbb{N}, l \in \mathbb{N}^*$, we have $b = a_+/2 = la_- - ja_+$. Thus, setting

$$\alpha_{jl} \equiv \frac{\pi}{2l}(1 + 2j), \quad j \in \mathbb{N}, \quad l \in \mathbb{N}^*, \quad (4.27)$$

we deduce from (4.18)

$$u(\pi, \alpha_{jl}, \pi/2; z) = \prod_{m=1}^j \frac{\text{sh} \pi \alpha_{jl}^{-1}(z + im\pi)}{(z \rightarrow -z)} \prod_{k=1}^{l-1} \frac{\text{sh}(z + ik\alpha_{jl})}{(z \rightarrow -z)} \quad (\text{sG}). \quad (4.28)$$

We proceed by obtaining and studying integral representations. In view of (3.1) and (3.3), we may rewrite u (4.1) as

$$u(z) = \exp(E(z)) \quad (4.29)$$

with

$$E(z) \equiv 2i \int_0^\infty \frac{dy}{y} \frac{\text{sh}(a_+ - b)y \text{sh}(a_- - b)y}{\text{sha}_+ y \text{sha}_- y} \sin 2yz. \quad (4.30)$$

Clearly, the integral converges absolutely provided

$$|\text{Im} z| < d(a_+, a_-, b)/2, \quad (4.31)$$

where

$$d(a_+, a_-, b) \equiv a_+ + a_- - |a_+ - b| - |a_- - b|. \quad (4.32)$$

In particular, one has

$$d(a_+, a_-, b) > a_\delta \Leftrightarrow (a_+, a_-, b) \in \mathcal{R}_\delta, \quad (4.33)$$

cf. (4.10). This bound amounts to the regularity of $u(z)$ in \mathcal{R}_δ , viewed as a solution to (4.6): u has no poles and zeros in the strip $|\text{Im} z| \leq a_\delta/2$ when $(a_+, a_-, b) \in \mathcal{R}_\delta$.

More generally, setting

$$\mathcal{C} \equiv \{(a_+, a_-, b) \in \mathcal{A} \mid b \in (0, a_+ + a_-)\}, \quad (4.34)$$

the representation (4.29) makes sense and holds true in a strip around the real z -axis if and only if the parameters belong to \mathcal{C} . Indeed, one easily verifies

$$d(a_+, a_-, b) > 0 \Leftrightarrow (a_+, a_-, b) \in \mathcal{C}. \quad (4.35)$$

Observe that $\mathcal{R}_+ \cup \mathcal{R}_-$ is a proper subset of \mathcal{C} .

Letting $|\operatorname{Im} z| < a_\delta/2$ and choosing parameters in \mathcal{R}_δ , we can derive a second integral representation from Theorems II.3 and II.2, as applied to the AΔE (4.6). From (4.29) and (4.30) we read off that the minimum integer k in Theorem II.3 equals 1. Setting

$$\phi_\delta(z) \equiv \ln \left(\frac{s_\delta(z - ib + ia_\delta/2) s_\delta(z + ib - ia_\delta/2)}{s_\delta(z + ia_\delta/2) s_\delta(z - ia_\delta/2)} \right) \quad (4.36)$$

with \ln real for z real, we now deduce

$$E(z) = \frac{1}{2ia_\delta} \int_{-\infty}^{\infty} dx \phi_\delta(x) \operatorname{th} \frac{\pi}{a_\delta}(z - x), \quad (a_+, a_-, b) \in \mathcal{R}_\delta, \quad |\operatorname{Im} z| < a_\delta/2. \quad (4.37)$$

(Indeed, both lhs and rhs vanish for $z=0$, and equality of derivatives is easily derived via (2.27) with $a \rightarrow a_\delta$ and $\phi(u) \rightarrow \phi'_\delta(u)$.) Notice that the integral on the rhs converges absolutely for real z and any $(a_+, a_-, b) \in \mathcal{H}$; even so, (4.37) is in general false for parameters not belonging to \mathcal{R}_δ . Note also that for parameters in $\mathcal{R}_+ \cap \mathcal{R}_-$ one gets two different representations without manifest $a_+ \leftrightarrow a_-$ symmetry.

Using the identity (A42) we can rewrite (4.37) as

$$E(z) = \frac{\operatorname{sh}(2\pi z/a_\delta)}{ia_\delta} \int_0^\infty \frac{\phi_\delta(x) dx}{\operatorname{ch}(2\pi z/a_\delta) + \operatorname{ch}(2\pi x/a_\delta)}, \quad (a_+, a_-, b) \in \mathcal{R}_\delta, \quad |\operatorname{Im} z| < a_\delta/2. \quad (4.38)$$

Combining this with (A43), (A44) and the Plancherel relation for the cosine transform, one recovers the symmetric representation (4.30).

We proceed by deriving yet another asymmetric representation for the u -function, in terms of an infinite product of gamma functions. (Somewhat surprisingly, this representation is not an easy consequence of (3.63), (3.64) and (3.67).) First, we introduce

$$\gamma_l(\rho, g, s) \equiv \Gamma(s+1+l/\rho) \Gamma(-s+g+l/\rho) \Gamma(s+l/\rho) \Gamma(-s+1-g+l/\rho) / (s \rightarrow -s), \quad (4.39)$$

where $l \in \mathbb{N}$, $\rho \in \mathbb{C}^-$, $g, s \in \mathbb{C}$. Fixing l, g, s and taking $\rho > 0$ and small enough, we may invoke (A45) to deduce

$$\gamma_l(\rho, g, s) = \exp \left(4 \int_0^\infty \frac{dy}{y} \frac{\operatorname{sh}(g-1)y \operatorname{sh} 2sy \operatorname{sh} gy}{\operatorname{sh} y} e^{-2ly/\rho} \right). \quad (4.40)$$

This representation is well defined and valid for

$$l \operatorname{Re}(\rho^{-1}) > |\operatorname{Re} g| + |\operatorname{Re} s|. \quad (4.41)$$

By virtue of (B18) it can be rewritten

$$\gamma_l(\rho, g, s) = \exp \left(4 \int_0^\infty e^{-2lt/\rho} f_3(g-1, 2s, g, t) dt \right). \quad (4.42)$$

Next, we assert that the function

$$P(\rho, g, s) \equiv \lim_{N \rightarrow \infty} \prod_{l=1}^N \gamma_l(\rho, g, s) \quad (4.43)$$

is well defined and meromorphic in $\mathbb{C}^- \times \mathbb{C}^2$. To prove this, we fix a compact $K \subset \mathbb{C}^- \times \mathbb{C}^2$ and put $w \equiv 2l/\rho$. Letting (ρ, g, s) vary over K , we can ensure (by taking $l \geq L$ with L large enough) that the bound (B21) applies for a suitable $\chi \in (0, \pi/2)$ and R (depending on K). Thus we deduce that γ_l is analytic on K and satisfies

$$|\gamma_l(\rho, g, s) - 1| \leq C_K/l^2, \quad \forall (\rho, g, s) \in K, \quad \forall l \geq L. \quad (4.44)$$

Consequently, the function $\prod_{l=L}^N \gamma_l$ converges uniformly on K to an analytic function for $N \rightarrow \infty$, and the assertion easily follows.

We claim that u can be written

$$u(a_+, a_-, b; z) = \frac{\Gamma\left(\frac{iz}{a_-} + 1\right) \Gamma\left(-\frac{iz}{a_-} + \frac{b}{a_-}\right)}{(z \rightarrow -z)} P\left(\frac{a_-}{a_+}, \frac{b}{a_-}, \frac{iz}{a_-}\right). \quad (4.45)$$

Since we already know that u is meromorphic for $(a_-/a_+, b, z) \in \mathbb{C}^- \times \mathbb{C}^2$, we need only prove this for $z = x \in \mathbb{R}$ and parameters in \mathcal{C} (4.34). To this end we show that the rhs is given by $\exp(E(x))$ (with $E(x)$ defined by (4.30)): Using (A45) and (4.40) we have (with $g \equiv b/a_-$)

$$\begin{aligned} & \frac{\Gamma\left(\frac{ix}{a_-} + 1\right) \Gamma\left(-\frac{ix}{a_-} + g\right)}{(x \rightarrow -x)} \prod_{l=1}^N \gamma_l\left(\frac{a_-}{a_+}, g, \frac{ix}{a_-}\right) \\ &= \exp\left(2i \int_0^\infty \frac{dy}{y} \frac{\operatorname{sh}(1-g)y \sin(2xy/a_-)}{\operatorname{sh}y} \left(e^{-gy} - 2 \operatorname{sh}g y \sum_{l=1}^N \exp(-2ly a_+/a_-)\right)\right) \\ &= \exp\left(i \int_0^\infty \frac{dy}{y} \frac{\operatorname{sh}(a_- - b)y \sin(2xy)}{\operatorname{sh}a_- y \operatorname{sh}a_+ y} (e^{-by}(e^{a_+y} - e^{-a_+y}) \right. \\ & \quad \left. + (e^{-by} - e^{by})e^{-a_+y}(1 - e^{-2a_+Ny}))\right). \end{aligned} \quad (4.46)$$

A dominated convergence argument now shows that we may take $N \rightarrow \infty$ under the integral sign, yielding the limit $\exp(E(x))$, as claimed.

We conclude this subsection by deriving four distinct limits of the u -function, using parameters

$$a_+ \equiv \pi, a_- \equiv \beta\nu, b \equiv \beta\nu g, \quad \beta, \nu > 0, \quad g \in \mathbb{R}. \quad (4.47)$$

First, we assert that

$$\lim_{\beta \downarrow 0} u(\pi, \beta\nu, \beta\nu g; \beta p) = \frac{\Gamma\left(\frac{ip}{\nu} + 1\right) \Gamma\left(-\frac{ip}{\nu} + g\right)}{(p \rightarrow -p)} \quad (\Pi_{\text{nr}} \text{ limit}), \quad (4.48)$$

where the limit is mero-uniform in p . To show this, we use (4.1), (3.22), (3.24) and (3.69) to write

$$u(\pi, \beta\nu, \beta\nu g; \beta p) = \frac{H(\rho; p/\nu - ig + i/2)H(\rho; -p/\nu - i/2)}{(p \rightarrow -p)}, \quad \rho \equiv \beta\nu/\pi. \quad (4.49)$$

Then the assertion follows from (3.72).

The formula (4.48) can be interpreted as the (nonrelativistic) Π_{nr} limit of the (relativistic) Π_{rel} S -matrix, cf. Ref. 1, Eq. (3.45). It can also be derived from the product representation (4.45). Indeed, one has

$$\lim_{\rho \downarrow 0} P(\rho, g, s) = 1 \quad (4.50)$$

uniformly for g, s in a fixed compact $B \subset \mathbb{C}^2$. To verify this, note first that $\gamma_l(\rho, g, s)$ (4.39) is analytic in B for $\rho > 0$ small enough, and given by (4.40). From this representation it follows that $\gamma_l(\rho, g, s)$ converges to 1 as $\rho \downarrow 0$, uniformly for $(g, s) \in B$. Next, observe that for $\rho \leq \epsilon$ (with ϵ depending only on B) one may use (4.42) and the bound (B21) with $w \equiv 2l/\rho$ to deduce

$$|\gamma_l(\rho, g, s) - 1| \leq C_B \rho^2 / l^2 \leq C_B \epsilon^2 / l^2, \quad \forall (g, s) \in B, \quad \forall l \in \mathbb{N}^*. \quad (4.51)$$

Clearly, this bound suffices to dominate the l -dependence, so one infers $P \rightarrow 1$, uniformly on B .

The next limit amounts to taking the I_{rel} limit of the dual Π_{rel} S -matrix, cf. Ref. 1: We claim

$$\lim_{\beta \downarrow 0} u(\pi, \beta\nu, \beta\nu g; \nu x) = \exp(i\pi(1-g)), \quad x \in \mathcal{R}_0 \quad (\text{I}_{\text{rel}} \text{ limit}), \quad (4.52)$$

where the limit is uniform on compacts of \mathcal{R}_0 (2.73). Before proving this, let us note that the restriction on x is essential: for $\text{Re} x < 0$ one obtains the complex conjugate phase factor by virtue of (4.3). (For $g \notin \mathbb{Z}$, the poles and zeros of u become dense on the imaginary axis as $\beta \downarrow 0$, cf. (4.1) and Prop. III.3.) Observe also that the phase amounts to a limit of the phase in (4.12).

To prove (4.52), we use the product representation (4.45) and several results from Appendix B. First, we handle the prefactor

$$Q_\beta(g, x) \equiv \frac{\Gamma(ix/\beta + 1)\Gamma(-ix/\beta + g)}{(x \rightarrow -x)}. \quad (4.53)$$

It can be rewritten

$$Q_\beta(g, x) = e^{i\pi(1-g)} \left(\frac{\Gamma(w_+ + 1)}{\Gamma(w_+ + g)} e^{(g-1)\text{In} w_+} \right) \left(\frac{\Gamma(w_- + g)}{\Gamma(w_- + 1)} e^{(1-g)\text{In} w_-} \right), \quad w_\pm \equiv \pm \frac{ix}{\beta}. \quad (4.54)$$

Using (B23) to rewrite the functions in brackets, and letting x vary over a fixed compact $K \subset \mathcal{R}_0$, we now exploit the bound (B20). First, taking $R = 1 + |g|$ and $\chi = \pi/4$ (say), one can ensure $w_+, w_- \in S_{R, \chi}$ for all $x \in K$ by choosing β small enough. Then it follows from (B20) that

$$\lim_{\beta \downarrow 0} Q_\beta(g, x) = \exp(i\pi(1-g)) \quad (4.55)$$

uniformly for $x \in K$. (This may be viewed as the $\Pi_{\text{nr}} \rightarrow \text{I}_{\text{nr}}$ S -matrix limit, cf. Ref. 1, Eq. (3.45).)

It remains to prove

$$\lim_{\rho \downarrow 0} P(\rho, g, iy/\rho) = 1 \quad (4.56)$$

uniformly on compacts of $\{\text{Re} y > 0\}$. To this end we first use (4.39) and (B23) to write

$$\begin{aligned}\gamma_l(\rho, g, iy/\rho) = & \exp(\mathcal{L}_2((iy+l)/\rho, 1, g) + \mathcal{L}_2((-iy+l)/\rho, g, 1) + \mathcal{L}_2((iy+l)/\rho, 0, 1-g) \\ & + \mathcal{L}_2((-iy+l)/\rho, 1-g, 0)).\end{aligned}\quad (4.57)$$

Next, we let y vary over a compact $K \subset \mathcal{R}_0$, and use the bound (B20) in the same way as before to infer that $\gamma_l \rightarrow 1$ for $\rho \downarrow 0$, uniformly on K .

As a consequence, (4.56) will follow provided we can supply a bound controlling the interchange of limits $N \rightarrow \infty$ and $\rho \downarrow 0$. Now the estimate (B20) is not sufficiently strong, since it only leads to $1/l$ -decrease of $|\gamma_l - 1|$, and the sequence $(1, 1/2, 1/3, \dots)$ is not in l^1 . But we can obtain a suitable bound by combining the representation (4.42) with the estimates (B21) and (B26), as follows.

We begin by observing that (4.42) and (B15) entail

$$\gamma_l(\rho, g, iy/\rho) = \exp(4\mathcal{L}_3(2l/\rho, g-1, g, 2iy/\rho)). \quad (4.58)$$

Letting y vary over K and choosing $\rho \in (0, \epsilon]$ with ϵ small enough, we may take $r_3 = c_K/\rho$ in the bound (B21) on \mathcal{L}_3 . Choosing now $\chi = 0, R = (c_K + 1)/\rho$ and $L > (c_K + 1)/2$, we deduce

$$\left| \mathcal{L}_3\left(\frac{2l}{\rho}, g-1, g, \frac{2iy}{\rho}\right) - \frac{ipyg(g-1)}{l^2} \right| \leq \frac{\rho^3}{4l^2} C_3, \quad \rho \in (0, \epsilon], \quad l \geq L, \quad y \in K. \quad (4.59)$$

Next, we use the bound (B26) to majorize the rhs of (4.59) by $C\rho/l^2$. By dominated convergence, this suffices to conclude that the function $\Pi_{l=L}^\infty \gamma_l$ converges to 1 as $\rho \downarrow 0$, uniformly on K . Since we have already shown that $\gamma_l \rightarrow 1$ uniformly on K for all $l \geq 1$, we may now deduce (4.56). (Notice that (4.58) and (B21) are not adequate for showing $\gamma_l \rightarrow 1$ for small l ; this is why we used (4.57) and (B19).)

Alternatively, (4.52) can be derived as a corollary of Prop. III.7. Indeed, from (4.1) we have

$$u(\pi, a, ag; z) = \frac{G(\pi, a; z + i\pi/2 + ia(1/2 - g))}{G(\pi, a; z + i\pi/2 - ia/2)} \cdot \frac{G(\pi, a; z - i\pi/2 + ia(g - 1/2))}{G(\pi, a; z - i\pi/2 + ia/2)}. \quad (4.60)$$

Thus, we may use (3.91) with $\operatorname{Re} z > 0$ to deduce the limit (4.52).

It is of interest to reconsider this limit in the setting of Theorem II.4. Choosing, e.g., $g \in (1/2, 1)$, one can take $f_a(z)$ equal to $\partial_z \ln u(\pi, a, ag; z)$; letting $a \rightarrow 0$, one gets $s_m(a) \rightarrow 0$ and $f_a(z) \rightarrow 0$ uniformly on compacts in the left and right half planes. Even so, $f_a(z)$ does not remain bounded near the origin, since $u(z)$ has distinct limits in the left and right half planes.

We continue by obtaining a third limit of the u -function, keeping the parameters (4.47), but now taking b fixed while letting $\beta \downarrow 0$. Specifically, we claim

$$\begin{aligned}\lim_{\beta \downarrow 0} \exp\left(-\frac{2ip}{\nu} \ln\left(\frac{\beta}{2\sin b}\right)\right) u(\pi, \beta\nu, b; \beta p) &= \frac{\Gamma\left(\frac{ip}{\nu} + 1\right)}{(p \rightarrow -p)} \exp\left(\frac{2ip}{\nu} \ln(2\nu)\right), \\ &b \in (0, \pi) \quad (\text{VI}_{\text{nr}} \text{ limit}),\end{aligned}\quad (4.61)$$

where the limit is mero-uniform. The function on the rhs may be viewed as the (nonrelativistic Toda) VI_{nr} S -matrix, cf. Ref. 1, Eq. (3.45). The limiting transition $\Pi_{\text{rel}} \rightarrow \text{VI}_{\text{nr}}$ is readily controlled at the level of the Poisson commuting classical Hamiltonians, cf. the paragraph containing Eq. (3.87) in Ref. 2. Formally, it also holds true for the corresponding quantum Hamiltonians. The S -matrix limit (4.61) agrees with the obvious conjecture that the limit holds true for the suitably normalized (reduced $N=2$) eigenfunctions; the plane wave factor on the lhs reflects the diverging position shift (3.87) in Ref. 2.

To prove (4.61), we begin by observing that

$$\lim_{g \uparrow \infty} \exp\left(\frac{2ip}{\nu} \lg\right) \frac{\Gamma\left(-\frac{ip}{\nu} + g\right)}{(p \rightarrow -p)} = 1 \quad (4.62)$$

uniformly on p -compacts. (This limit amounts to the $\Pi_{\text{nr}} \rightarrow \text{VI}_{\text{nr}}$ S -matrix limit, cf. Ref. 1, Eq. (3.45), and the paragraph containing Eq. (2.116) in Ref. 2.) Indeed, this follows from (B23) and (B20) (taking $w=g$) in a by now familiar way. As a result, (4.61) will follow once we show

$$\lim_{\rho \downarrow 0} P(\rho, b/\pi\rho, s) = \exp\left(2s \ln\left(\frac{b}{\sin b}\right)\right), \quad b \in (0, \pi), \quad (4.63)$$

uniformly on s -compacts.

To prove (4.63), we write

$$\begin{aligned} \gamma_l(\rho, b/\pi\rho, s) &= \exp(\mathcal{L}_2(l/\rho, s+1, -s+1)) \exp(\mathcal{L}_2(l/\rho, s, -s)) \\ &\quad \times \exp(\mathcal{L}_2((l\pi+b)/\pi\rho, -s, s)) \\ &\quad \times \exp(\mathcal{L}_2((l\pi-b)/\pi\rho, -s+1, s+1)) \exp(-2s \ln(1-b^2/l^2\pi^2)). \end{aligned} \quad (4.64)$$

Since $b \in (0, \pi)$, we have $l\pi \pm b > 0$, and so we conclude using (B20)

$$\lim_{\rho \downarrow 0} \gamma_l(\rho, b/\pi\rho, s) = \exp(-2s \ln(1-b^2/l^2\pi^2)) \quad (4.65)$$

uniformly on s -compacts. Now from (A23)–(A25) [with $\alpha=0$, cf. (A28)] one derives the well-known identity

$$\frac{\sin b}{b} = \prod_{l=1}^{\infty} \left(1 - \frac{b^2}{l^2\pi^2}\right). \quad (4.66)$$

Using this on the rhs of (4.63) and comparing with (4.65), we infer that we need only supply a bound that is sufficiently strong to render the interchange of limits legitimate.

The bound (B20) leads to an $O(l^{-1})$ -majorization, so it is not strong enough. Just as in the previous case, we will now derive an $O(l^{-2})$ estimate (for l sufficiently large) by combining (B21) and (B26). To this purpose we observe that we may write

$$\gamma_l(\rho, b/\pi\rho, s) = \exp(4\mathcal{L}_3(2l/\rho, -1+b/\pi\rho, 2s, b/\pi\rho)), \quad (4.67)$$

cf. (4.42) and (B15). For s in a compact $B \subset \mathbb{C}$ and $\rho \in (0, \epsilon]$ with ϵ small enough, we can take $r_3 = c_B/\rho$ in (B21). Choosing then $\chi=0, R=(c_B+1)/\rho$ and $L > (c_B+1)/2$, we obtain

$$\left| \mathcal{L}_3\left(\frac{2l}{\rho}, -1 + \frac{b}{\pi\rho}, 2s, \frac{b}{\pi\rho}\right) - \frac{sb(b-\pi\rho)}{2l^2\pi^2} \right| \leq \frac{\rho^3}{4l^2} C_3, \quad \rho \in (0, \epsilon], \quad l \geq L, \quad s \in B. \quad (4.68)$$

Using now (B26), we obtain an upper bound C/l^2 on the rhs. As before, this suffices to conclude that (4.63) holds true. The upshot is that the proof of (4.61) is now complete.

As a corollary of (4.61), we can obtain the integral

$$\frac{\Gamma(1+iz)}{\Gamma(1-iz)} = \exp\left(\frac{\text{sh}2\pi z}{2i} \int_0^\infty \frac{dt}{\text{ch}2\pi z + \text{ch}\pi t} \ln\left(\frac{4}{t^2+1}\right)\right), \quad |\text{Im}z| < \frac{1}{2}. \quad (4.69)$$

Indeed, combining the integral

$$\operatorname{sh} 2\pi z \int_0^\infty \frac{dt}{\operatorname{ch} 2\pi z + \operatorname{ch} \pi t} = 2z, \quad |\operatorname{Im} z| < \frac{1}{2} \quad (4.70)$$

(which results from (A43), e.g.), with (4.29), (4.38) and (4.36), we obtain

$$\begin{aligned} & \exp\left(-2iz \ln\left(\frac{\beta}{\sin b}\right)\right) u(\pi, \beta, b; \beta z) \\ &= \exp\left(\frac{\operatorname{sh} 2\pi z}{2i} \int_0^\infty \frac{dt}{\operatorname{ch} 2\pi z + \operatorname{ch} \pi t} \ln\left(\frac{\operatorname{sh}^2 \beta t/2 + \sin^2(b - \beta/2)}{\operatorname{sh}^2 \beta t/2 + \sin^2 \beta/2} \cdot \frac{\beta^2}{\sin^2 b}\right)\right), \end{aligned} \quad (4.71)$$

where $\beta \in (0, b/2)$, $b \in (0, \pi)$, $|\operatorname{Im} z| < 1/2$. A straightforward dominated convergence argument now shows that the rhs of (4.71) converges to the rhs of (4.69) for $\beta \downarrow 0$. From (4.61) we see that the lhs converges to the lhs of (4.69), so (4.69) results.

Finally, we obtain a limit that may be viewed as the classical limit of the quantum Π_{rel} S -matrix. To this end we introduce

$$L_{\hbar}(p) \equiv i\hbar \ln u(\pi, \hbar/\lambda, b; p), \quad (\lambda, b, p) \in (0, \infty) \times (0, \pi) \times \mathcal{R}_0, \quad (4.72)$$

with $\ln u \rightarrow 0$ for $p \rightarrow 0$, $\hbar > 0$ denoting Planck's constant. We now claim that

$$\lim_{\hbar \rightarrow 0} \partial_p L_{\hbar}(p) = \lambda \ln\left(\frac{\operatorname{sh}(p+ib)\operatorname{sh}(p-ib)}{\operatorname{sh}^2 p}\right) \quad (\text{classical limit}) \quad (4.73)$$

uniformly on compact subsets of the right half plane \mathcal{R}_0 , with \ln real valued for $p > 0$. (The rhs amounts to the classical Π_{rel} phase shift, cf. Ref. 1, Eq. (2.75) with $\beta = 1$.)

To prove this claim, we substitute $ag \rightarrow b$ in (4.60) and use (3.83) and (3.84) to write

$$\begin{aligned} i\lambda \ln u(\pi, a, b; z) &= -D_a(z + i\pi/2 - ib) - D_a(z - i\pi/2 + ib) + D_a(z + i\pi/2) + D_a(z - i\pi/2) \\ &\quad - ad_a(1/2, 0; z + i\pi/2 - ib) - ad_a(-1/2, 0; z - i\pi/2 + ib) \\ &\quad + ad_a(-1/2, 0; z + i\pi/2) + ad_a(1/2, 0; z - i\pi/2). \end{aligned} \quad (4.74)$$

Taking $a \rightarrow 0$, the limit of (4.74) exists uniformly on compacts in \mathcal{R}_0 by virtue of (3.85) and (3.86). Taking z -derivatives, one readily obtains a limit that amounts to (4.73).

B. The elliptic case

The elliptic scattering function is defined in terms of the elliptic G -function from Subsection III B via (4.1). In view of Prop. III.11, this yields a function that is meromorphic in r, a_+, a_-, b and z , as long as $a_+ r$ and $a_- r$ stay in the right half plane. We shall from now on restrict the parameters to

$$\mathcal{E} \equiv \{(r, a_+, a_-, b) | r > 0, (a_+, a_-, b) \in \mathcal{R}\}, \quad (4.75)$$

cf. (4.5). By virtue of Prop. III.9 the elliptic u -function is periodic in z with primitive period π/r ; moreover, it satisfies (4.2), (4.3), and

$$u(2r, a_+, a_-, b; z) = u(r, a_+, a_-, b; z) u(r, a_+, a_-, b; z - \pi/2r), \quad (4.76)$$

$$u(\lambda^{-1}r, \lambda a_+, \lambda a_-, \lambda b; \lambda z) = u(r, a_+, a_-, b; z), \quad \lambda \in (0, \infty). \quad (4.77)$$

Recalling (3.96)–(3.100), and using also (2.91), we see that u solves the AΔEs

$$\frac{u(z+ia_{\delta}/2)}{u(z-ia_{\delta}/2)} = \exp(2r(a_{\delta}-b)) \frac{s_{-\delta}(z-ib+ia_{\delta}/2)s_{-\delta}(z+ib-ia_{\delta}/2)}{s_{-\delta}(z+ia_{\delta}/2)s_{-\delta}(z-ia_{\delta}/2)}. \quad (4.78)$$

It now follows just as in the hyperbolic case that u is a regular solution to (4.78) if and only if $(a_+, a_-, b) \in \mathcal{R}_{\delta}$. Since u is π/r -periodic in z , the latter restriction also ensures that u is the unique minimal solution satisfying (4.11). Furthermore, with (4.6) replaced by (4.78), the remark below (4.12) applies verbatim to the elliptic case.

Using (3.100) and (2.91) we now obtain the analog of (4.13):

$$\frac{u(b+a_{-\delta};z)}{u(b;z)} = -e^{2irz} \frac{s_{\delta}(z+ib)}{s_{\delta}(z-ib)}. \quad (4.79)$$

To simplify the iterations of these AΔEs, we use the formula

$$\frac{s(r, a; z_+ + ina)}{s(r, a; z_- - ina)} = e^{-2irn(z_+ + z_-)} \frac{s(r, a; z_+)}{s(r, a; z_-)}, \quad n \in \mathbb{N}, \quad (4.80)$$

which follows from (2.91). Then we obtain once more the relation (4.14), but now with an extra factor $\exp(2irz(k_+ + k_- - 2k_+k_-))$ on the rhs. Noting the elliptic analog

$$u(r, a_+, a_-, a_+; z) = u(r, a_+, a_-, a_-; z) = 1 \quad (4.81)$$

of (4.16), we deduce the elliptic analog

$$u(r, a_+, a_-, 0; z) = -e^{-2irz} \quad (4.82)$$

of (4.17) and, more generally, the explicit formula (4.18), with (4.20) replaced by

$$c_{k,l} = (-)^{k+l+1} \exp(2irz(k+l-2kl-1)), \quad k, l \in \mathbb{Z} \quad (4.83)$$

It is clear that the symmetry property (4.21) continues to hold in the elliptic case. Moreover, it leads again to the relation (4.22) between $u(-b; z)$ and $u(b; z)$. Next, we note that (4.23) still holds true, since the elliptic G -function satisfies the multiplication formula (3.25). Hence, (4.24) follows as before. Using the AΔEs (3.100) and (2.91) we then obtain as the analogs of (4.25) and (4.26)

$$\begin{aligned} u\left(r, \frac{a_+}{M}, a_-, b; z\right) &= (-)^{M-1} \exp(ir(M-1)(2Mz + ia_+ - ib)) \\ &\cdot \prod_{k=0}^{M-1} u\left(r, a_+, a_-, b; z + ik \frac{a_+}{M}\right) \prod_{j=1}^{M-1} \frac{s_{-}(z + ija_+/M)}{s_{-}(z - ib + ija_+/M)} \end{aligned} \quad (4.84)$$

and

$$\begin{aligned} u\left(r, \frac{a_+}{M}, a_-, b; z\right) &= (-)^{M-1} \exp(ir(M-1)(2Mz - ia_+ + ib)) \\ &\cdot \prod_{k=0}^{M-1} u\left(r, a_+, a_-, b; z - ik \frac{a_+}{M}\right) \prod_{j=1}^{M-1} \frac{s_{-}(z + ib - ija_+/M)}{s_{-}(z - ija_+/M)}. \end{aligned} \quad (4.85)$$

Once more, $a_+ \leftrightarrow a_-$ symmetry can now be used to obtain four distinct representations for $u(r, a_+/M, a_-/N, b; z)$ in terms of $u(r, a_+, a_-, b; z)$ and s -quotients.

The choices $b = a_+/2$ or $b = a_-/2$ yield the XYZ soliton-soliton S -matrix. Thus it follows from (4.18) and (4.83) that the counterpart of (4.28) reads

$$u(r, \pi, \alpha_{jl}, \pi/2; z) = \exp(2irz(l-j+2lj-1)) \cdot \prod_{m=1}^j \frac{s(r, \alpha_{jl}; z + im\pi)}{(z \rightarrow -z)} \prod_{k=1}^{l-1} \frac{s(r, \pi; z + ik\alpha_{jl})}{(z \rightarrow -z)} \quad (\text{XYZ}). \quad (4.86)$$

Next, we use (4.1), (3.92) and (3.3) to obtain

$$u(z) = \exp(E(z)) = \exp\left(2i \sum_{n=1}^{\infty} \frac{\text{sh}(a_+ - b)nr \text{sh}(a_- - b)nr}{n \text{sha}_+ nr \text{sha}_- nr} \sin 2nrz\right). \quad (4.87)$$

The series converges absolutely if and only if (4.31) holds true. As before, regularity of $u(z)$ for parameters in \mathcal{R}_δ can be read off from (4.33). Furthermore, the series representation (4.87) is valid for real z iff the parameters belong to the convergence region (4.34).

Choosing $(a_+, a_-, b) \in \mathcal{R}_\delta$ and introducing

$$\phi_\delta(z) \equiv \ln \left(\frac{s_{-\delta}(z - ib + ia_\delta/2) s_{-\delta}(z + ib - ia_\delta/2)}{s_{-\delta}(z + ia_\delta/2) s_{-\delta}(z - ia_\delta/2)} \right) + 2r(a_\delta - b) \quad (4.88)$$

with \ln real for z real, we can combine (4.78) and (4.87) to deduce that $\phi_\delta(z)$ satisfies the assumptions (2.100)–(2.102) of Theorem II.5. Therefore, (2.107) yields

$$E(z) = \frac{1}{2i\pi} \int_{-\pi/2r}^{\pi/2r} dy \phi_\delta(y) K(r, a_\delta; z - y), \quad (a_+, a_-, b) \in \mathcal{R}_\delta, \quad |\text{Im}z| < \frac{a_\delta}{2}. \quad (4.89)$$

This representation amounts to the elliptic counterpart of (4.37). Once more, the restriction on the parameters is essential (though boundary points of \mathcal{R}_δ belonging to \mathcal{H} (4.5) can be allowed, of course).

The product representation (3.117) for the elliptic G -function can be combined with (4.1) to yield

$$u(r, a_+, a_-, b; z) = \prod_{m,n=1}^{\infty} \frac{(1 - 2q_+^{2m-1} q_-^{2n-1} e^{-2irz} \text{ch}(b - (a_+ + a_-)/2) + q_+^{4m-2} q_-^{4n-2} e^{-4irz})}{(z \rightarrow -z)} \cdot \frac{(1 - 2q_+^{2m-1} q_-^{2n-1} e^{2irz} \text{ch}(a_+ - a_-)/2 + q_+^{4m-2} q_-^{4n-2} e^{4irz})}{(z \rightarrow -z)}, \quad q_\delta \equiv e^{-a_\delta r}. \quad (4.90)$$

From this product representation one can read off meromorphy and pole/zero properties of $u(z)$. Notice that it is manifestly symmetric in a_+, a_- , in contradistinction to the product representation (4.45) for the hyperbolic u -function.

We proceed by deriving four limits of the u -function. First, we observe that

$$\lim_{r \downarrow 0} u(r, a_+, a_-, b; z) = u_{\text{hyp}}(a_+, a_-, b; z) \quad (\text{II}_{\text{rel}} \text{ limit}), \quad (4.91)$$

where the limit is mero-uniform. (Here, u_{hyp} denotes the u -function from Subsection IV A.) Indeed, in the definition (4.1) of the elliptic u -function we may replace the elliptic G -functions by G_{ren} -functions, cf. (3.123). Then (4.91) is a consequence of Prop. III.12.

Second, we assert that the limit

$$\lim_{A \uparrow \infty} u(r, a, A, b; z) = u_{\text{trig}}(r, a, b; z) \quad (\text{III}_{\text{rel}} \text{ limit}) \quad (4.92)$$

exists mero-uniformly. (Here, u_{trig} denotes the u -function studied in the next subsection.) To prove this, we use (4.1) and (3.22) to write

$$u(r, a, A, b; z) = \frac{G(r, a, A; z + ib - ia/2 - iA/2)G(r, a, A; -z + ia/2 - iA/2)}{G(r, a, A; -z + ib - ia/2 - iA/2)G(r, a, A; z + ia/2 - iA/2)}. \quad (4.93)$$

Invoking now (3.139), we obtain the mero-uniform limit

$$\lim_{A \uparrow \infty} u(r, a, A, b; z) = \frac{G(r, a; z + ib - ia/2)G(r, a; -z + ia/2)}{G(r, a; -z + ib - ia/2)G(r, a; z + ia/2)}, \quad (4.94)$$

which amounts to (4.92), cf. (4.100) below.

Third, fixing $g \in \mathbb{R}$, we claim that

$$\lim_{a \downarrow 0} u(r, A, a, ag; z) = \exp((1-g)(i\pi - 2irz)), \quad z \in \mathcal{R}_r \quad (\text{IV}_{\text{nr}} \text{ limit}), \quad (4.95)$$

uniformly on compacts in the period strip \mathcal{R}_r (2.113). Indeed, from (4.93) and (3.138) we obtain

$$\lim_{a \downarrow 0} u(r, A, a, ag; z) = \exp((1-g)\ln(R(r, A; -z - iA/2)/R(r, A; z - iA/2))) \quad (4.96)$$

uniformly on compacts of \mathcal{R}_r . Now the limit (4.95) easily results from (3.93).

We continue by examining this result in the setting of Subsection II C. Taking $g \in [1, 2]$ and $a \in (0, A/4]$, it entails that Theorem II.7 applies to $f_a(z) \equiv \ln u(r, A, a, ag; z)$. In this case $f'_a(z)$ converges to the constant $2ir(g-1)$, uniformly on compacts $K \subset \mathcal{R}_r$, but $f'_a(z)$ diverges near $z=0$ as $a \rightarrow 0$. Indeed, the π/r -periodic function $f_a(x)$, $x \in \mathbb{R}$, converges pointwise to a π/r -periodic function $f(x)$ that has unequal limits for $x \downarrow 0$ and $x \uparrow \pi/r$ (unless $g=1$, of course). Notice in this connection that it does not follow from the above that $f_a(z)$ remains bounded in the strip $|\text{Im}z| \leq a/2$ as $a \rightarrow 0$; we do not know whether this holds true.

We conclude this subsection by deriving the generalization of the classical limit (4.73). Thus we define

$$L_{\hbar}(z) \equiv i\hbar \ln u(r, A, \hbar/\lambda, b; z), \quad (r, \lambda, b, z) \in (0, \infty)^2 \times (0, A) \times \mathcal{R}_r, \quad (4.97)$$

with $\ln u \rightarrow 0$ for $z \rightarrow 0$ and $\hbar > 0$ Planck's constant. Then we have

$$\lim_{\hbar \rightarrow 0} \partial_z L_{\hbar}(z) = \lambda \ln \left(e^{-2rb} \frac{s(r, A; z + ib)s(r, A; z - ib)}{s(r, A; z)^2} \right) \quad (\text{classical limit}) \quad (4.98)$$

uniformly on an arbitrary compact $K \subset \mathcal{R}_r$, with \ln real for $z \in (0, \pi/r)$.

To prove this assertion, we exploit the obvious generalization of (4.74) and Prop. III.13 to infer

$$\lim_{a \downarrow 0} ia \partial_z \ln u(r, A, a, b; z) = \ln \left(\frac{R(r, A; z + iA/2 - ib) R(r, A; z - iA/2 + ib)}{R(r, A; z + iA/2) R(r, A; z - iA/2)} \right) \quad (4.99)$$

uniformly on K . Using (3.93) and (2.91), we see that this limit amounts to (4.98). Notice that the limit can be understood from Theorem II.7 and (4.78), with $a \ln u(z)$ playing the role of $f_a(z)$.

C. The trigonometric case

The trigonometric scattering function is defined by

$$u(r, a, b; z) \equiv \frac{G(z + ib - ia/2) G(-z + ia/2)}{G(-z + ib - ia/2) G(z + ia/2)} \quad (4.100)$$

with $G(z) \equiv G(r, a; z)$ denoting the trigonometric G -function (3.140). From the corresponding product representation

$$u(r, a, b; z) = \prod_{m=1}^{\infty} \frac{(1 - q^{2m-2} e^{-2rb-2irz})(1 - q^{2m} e^{2irz})}{(1 - q^{2m-2} e^{-2rb+2irz})(1 - q^{2m} e^{-2irz})}, \quad q \equiv e^{-ar}, \quad (4.101)$$

we read off that u admits analytic continuation to a function that is meromorphic in r, a, b and z , provided ar stays in the right half plane. However, in the sequel we restrict the parameters to

$$\mathcal{T} \equiv \{(r, a, b) \mid r > 0, a > 0, b \in \mathbb{R}\}. \quad (4.102)$$

As before, this restriction entails $|u(z)| = 1$ for real z .

Obviously, u is periodic in z with primitive period π/r ; it also satisfies (4.3) and the relations

$$u(2r, a, b; z) = u(r, a, b; z) u(r, a, b; z - \pi/2r), \quad (4.103)$$

$$u(\lambda^{-1}r, \lambda a, \lambda b; \lambda z) = u(r, a, b; z), \quad \lambda \in (0, \infty). \quad (4.104)$$

From (2.90) and (4.78) [or directly from (4.100) and (3.142)] we deduce that u satisfies the AΔE

$$\frac{u(z + ia/2)}{u(z - ia/2)} = \exp(2r(a - b)) \frac{\sin r(z - ib + ia/2) \sin r(z + ib - ia/2)}{\sin r(z + ia/2) \sin r(z - ia/2)}. \quad (4.105)$$

Clearly, this AΔE is regular unless $b = a/2$. Now from the product representation (4.101) we see that $u(r, a, b; z)$ may be viewed as the unique minimal solution to (4.105) that obeys (4.11), provided the parameters belong to the regularity region

$$\mathcal{R} \equiv \{(r, a, b) \in \mathcal{T} \mid b \in (a/2, \infty)\}. \quad (4.106)$$

Next, we use (4.101) to conclude

$$\frac{u(b + a; z)}{u(b; z)} = -e^{2irz} \frac{\sin r(z + ib)}{\sin r(z - ib)}. \quad (4.107)$$

(Alternatively, this follows from (4.79) by taking a limit.) By iteration this gives rise to (taking $k \in \mathbb{Z}$)

$$\frac{u(b + ka; z)}{u(b; z)} = e^{2irkz} \prod_{j=1}^{|k|} \frac{\sin r(z + i(k/|k|)(b - a/2) + ia(j - 1/2))}{(z \rightarrow -z)}. \quad (4.108)$$

Now from the product representation (4.101) we read off

$$u(r, a, a; z) = 1, \quad (4.109)$$

$$u(r, a, 0; z) = -e^{-2irz}, \quad (4.110)$$

and so (4.108) entails

$$u(r, a, ka; z) = (-)^{k+1} e^{2ir(k-1)z} \prod_{j=1}^{|k|} \frac{\sin r(z + ia(j - \theta(k)))}{(i \rightarrow -i)}, \quad k \in \mathbb{Z}, \quad (4.111)$$

with $\theta(k)$ defined by (4.19).

The trigonometric specializations of the relations (4.84) and (4.85) read

$$\begin{aligned} u\left(r, \frac{a}{M}, b; z\right) &= (-)^{M-1} \exp(ir(M-1)(2Mz + ia - ib)) \\ &\cdot \prod_{k=0}^{M-1} u\left(r, a, b; z + ik \frac{a}{M}\right) \prod_{j=1}^{M-1} \frac{\sin r(z + ija/M)}{\sin r(z - ib + ija/M)} \end{aligned} \quad (4.112)$$

and

$$\begin{aligned} u\left(r, \frac{a}{M}, b; z\right) &= (-)^{M-1} \exp(ir(M-1)(2Mz - ia + ib)) \\ &\cdot \prod_{k=0}^{M-1} u\left(r, a, b; z - ik \frac{a}{M}\right) \prod_{j=1}^{M-1} \frac{\sin r(z + ib - ija/M)}{\sin r(z - ija/M)}. \end{aligned} \quad (4.113)$$

Of course, these formulas can also be verified directly from (4.100) and the multiplication formula (3.145).

We proceed by obtaining series and integral representations for the (logarithm of the) u -function. From (4.100) and (3.141) we obtain (formally at first)

$$u(z) = \exp(E(z)) = \exp\left(2i \sum_{n=1}^{\infty} \frac{e^{-bnr} \operatorname{sh}(a-b)nr}{n \operatorname{sh} nr} \sin 2nrz\right). \quad (4.114)$$

(Alternatively, this can be deduced from (4.87) and (4.92).) The series converges absolutely provided

$$|\operatorname{Im} z| < d(a, b)/2, \quad (4.115)$$

with

$$d(a, b) \equiv a + b - |a - b|. \quad (4.116)$$

Thus one has

$$d(a, b) > a \Leftrightarrow b > a/2 \quad (4.117)$$

in agreement with the fact that u is a minimal solution to the AΔE (4.105) for parameters in \mathcal{R} (4.106). More generally, the series representation (4.114) makes sense and holds true in a strip around the real z -axis iff the parameter b is positive.

Next, we take $(r, a, b) \in \mathcal{R}$ and set

$$\phi(z) \equiv \ln \left(\frac{\sin r(z - ib + ia/2) \sin r(z + ib - ia/2)}{\sin r(z + ia/2) \sin r(z - ia/2)} \right) + 2r(a - b) \quad (4.118)$$

with \ln real-valued for $z \in \mathbb{R}$. Obviously, ϕ satisfies the assumptions (2.100) and (2.101) of Theorem II.5, and comparing (4.105) and (4.114) it follows that ϕ satisfies (2.102), too. Thus, (2.107) applies, yielding the integral representation

$$E(z) = \frac{1}{2i\pi} \int_{-\pi/2r}^{\pi/2r} dy \phi(y) K(r, a; z - y), \quad (r, a, b) \in \mathcal{R}, \quad |\operatorname{Im} z| < \frac{a}{2}. \quad (4.119)$$

By continuity, the representation still holds for $b = a/2$, but it is false in general for $b < a/2$.

To conclude this subsection, we obtain three limits of the trigonometric scattering function. First, we use (3.155) to write

$$u(r, 1, b; z) = \frac{T(r; z - ib + i/2) T(r; -z - i/2)}{T(r; -z - ib + i/2) T(r; z - i/2)} \exp(2ir(b - 1)z). \quad (4.120)$$

Then it follows from Prop. III.20 that we have

$$\lim_{r \downarrow 0} u(r, 1, g; z) = \frac{\Gamma(-iz + g) \Gamma(iz + 1)}{(z \rightarrow -z)} \quad (\Pi_{\text{nr}} \text{ limit}) \quad (4.121)$$

mero-uniformly in z . (Compare this to (4.48).)

Second, we observe that

$$\lim_{a \downarrow 0} u(r, a, ag; z) = \exp((1 - g)(i\pi - 2irz)), \quad z \in \mathcal{R}_r \quad (\hat{\Pi}_{\text{nr}} \text{ limit}), \quad (4.122)$$

uniformly on compact subsets of the period strip \mathcal{R}_r (2.113). Indeed, this readily follows from (3.180), cf. also (4.95) and (4.96). The remark below (4.96) applies to the case at hand as well.

Third, we introduce

$$L_{\hbar}(z) \equiv i\hbar \ln u(r, \hbar/\lambda, b; z), \quad (r, \lambda, b, z) \in (0, \infty)^3 \times \mathcal{R}_r, \quad (4.123)$$

with $\ln u \rightarrow 0$ for $z \rightarrow 0$ and $\hbar > 0$ Planck's constant. Then we claim that

$$\lim_{\hbar \rightarrow 0} \partial_z L_{\hbar}(z) = \lambda \ln \left(e^{-2rb} \frac{\sin r(z + ib) \sin r(z - ib)}{\sin^2 rz} \right) \quad (\text{classical limit}) \quad (4.124)$$

uniformly on compacts of \mathcal{R}_r , with \ln real-valued for $z \in (0, \pi/r)$. To prove this claim, we use (4.100) and (3.176), (3.177) to write

$$\begin{aligned} i\lambda \ln u(r, a, b; z) &= -D_a(z + ib) + D_a(-z + ib) - D_a(-z) + D_a(z) - ad_a(r, -1/2, 0; z + ib) \\ &\quad + ad_a(r, -1/2, 0; -z + ib) - ad_a(r, 1/2, 0; -z) + ad_a(r, 1/2, 0; z), \end{aligned} \quad (4.125)$$

where we take $z \in \mathcal{R}_r$. Invoking now Prop. III.21, the limit (4.124) readily follows.

Comparing the rhs of (4.124) to the classical phase shift obtained in Ref. 17, p. 336, we get agreement when we take $\lambda \rightarrow \beta^{-1}$, $r \rightarrow |\mu|/2$, $b \rightarrow |\beta g|$, save for a constant shift $-2\lambda r b \rightarrow -|\mu g|$. The latter shift can be understood from the fact that the distance between the classical actions of the III_{rel} system is bounded below by $|\mu g|$ (cf. Ref. 17, p. 256); by contrast, the minimal distance between successive indices n_i, n_{i+1} of the multivariable polynomials occurring at the quantum level equals 0. (See also Ref. 2, Subsection 6.2.)

V. WEIGHT FUNCTIONS

A. The hyperbolic case

Our study of the hyperbolic weight function $w(a_+, a_-, b; z)$ runs largely parallel to our study of the u -function in Subsection IV A. The w -function is defined by

$$w(z) \equiv \frac{G(z+ib-i(a_++a_-)/2)G(z+i(a_++a_-)/2)}{G(z-ib+i(a_++a_-)/2)G(z-i(a_++a_-)/2)}, \quad (5.1)$$

so it satisfies

$$w(a_+, a_-; z) = w(a_-, a_+; z) \quad (5.2)$$

just as $G(z)$ and $u(z)$, cf. (4.1) and (4.2). The analogs of (4.3) and (4.4) are

$$w(-z) = w(z), \quad (5.3)$$

$$w(\lambda a_+, \lambda a_-, \lambda b; \lambda z) = w(a_+, a_-, b; z), \quad \lambda \in (0, \infty). \quad (5.4)$$

For several purposes it is convenient to introduce a reduced weight function

$$w_r(z) \equiv \frac{G(z+ib-i(a_++a_-)/2)}{G(z-ib+i(a_++a_-)/2)}. \quad (5.5)$$

Using the AΔEs (3.4), one infers that w and w_r are related by

$$w(z) = 4 \operatorname{sh}(\pi z/a_+) \operatorname{sh}(\pi z/a_-) w_r(z). \quad (5.6)$$

Obviously, w_r also satisfies (5.2)–(5.4).

Just as the u -function, the functions w and w_r are meromorphic in a_+, a_-, b and z , as long as a_-/a_+ stays away from $(-\infty, 0]$, cf. Prop. III.5. In particular, both u and w_r are well defined for $b, z \in \mathbb{C}$. Using (4.1) and (3.4), one readily verifies that the latter functions are related by

$$u(iz; ib) = w_r(b; z) \frac{4 \operatorname{sh} \frac{\pi}{a_+}(z+ib) \operatorname{sh} \frac{\pi}{a_-}(z+ib)}{G(ib-i(a_+-a_-)/2)G(ib+i(a_+-a_-)/2)}. \quad (5.7)$$

This relation can be used to translate various features of w_r in terms of u and vice versa.

From now on we take $(a_+, a_-, b) \in \mathcal{H}$ (4.5). We proceed by studying w and w_r with regard to the AΔEs they satisfy, namely

$$\frac{w(z+ia\delta/2)}{w(z-ia\delta/2)} = \frac{s_{-\delta}(z+ib-ia\delta/2)}{s_{-\delta}(z-ib+ia\delta/2)} \cdot \frac{s_{-\delta}(z+ia\delta/2)}{s_{-\delta}(z-ia\delta/2)} \quad (5.8)$$

and

$$\frac{w_r(z+ia\delta/2)}{w_r(z-ia\delta/2)} = - \frac{s_{-\delta}(z+ib-ia\delta/2)}{s_{-\delta}(z-ib+ia\delta/2)}, \quad (5.9)$$

resp. (To check this, recall the definition (4.7) and the AΔEs (3.4).)

Consider first w_r . The planes (4.9) separate the region \mathcal{H} (4.5) into infinitely many strip-like components, one of which reads

$$\mathcal{S}_\delta \equiv \{(a_+, a_-, b) \in \mathcal{H} \mid b \in (a\delta/2, a_{-\delta} + a\delta/2)\}. \quad (5.10)$$

The pole/zero properties of $G(z)$ given by Prop. III.3 entail that w_r is free of zeros and poles in the strip $|\operatorname{Im} z| \leq a_\delta/2$ if and only if $(a_+, a_-, b) \in \mathcal{S}_\delta$. Now from Prop. III.4 we deduce that for all $(a_+, a_-, b) \in \mathcal{H}$ one has

$$w_r(z) = \exp\left(\pm \frac{\pi z}{a_+ a_-} (2b - a_+ - a_-)\right) (1 + O(\exp(\pm(\epsilon - 2\pi/a_m)z))), \quad \operatorname{Re} z \rightarrow \pm\infty, \quad (5.11)$$

uniformly for $\operatorname{Im} z$ in \mathbb{R} -compacts. Thus, choosing parameters in \mathcal{S}_δ , one may characterize w_r as a minimal solution to the AΔE (5.9) that is even and positive for $z \in \mathbb{R}$; these properties determine the solution up to a positive constant, cf. Theorem II.1. Next, we note that the rhs of (5.9) is $a_{-\delta}$ -periodic in b , and identically equal to -1 for parameters satisfying (4.9). (As such, the AΔE is regular for all $(a_+, a_-, b) \in \mathcal{H}$, by contrast to (4.6).) But w_r is neither $a_{-\delta}$ -periodic in b , nor an exponential when (4.9) holds true. We shall presently obtain the corresponding ia_δ -periodic multiplier, after considering w in relation to the AΔE (5.8) it obeys.

We begin by noting that the w -function has asymptotics

$$w(z) = \exp\left(\pm \frac{2\pi bz}{a_+ a_-}\right) (1 + O(\exp(\pm(\epsilon - 2\pi/a_m)z))), \quad \operatorname{Re} z \rightarrow \pm\infty. \quad (5.12)$$

Thus, it is a minimal solution to (5.8) whenever it has no poles and zeros for $|\operatorname{Im} z| \leq a_\delta/2$. In view of (5.6), for this to happen it is necessary that $w_r(z)$ have a double pole at $z=0$. For a_+, a_- fixed, this necessary condition is satisfied only for a discrete set of b , so w is generically not a regular solution—in contrast to w_r , which is regular for parameters in \mathcal{S}_δ .

It should be pointed out, though, that both of the AΔEs (5.8) do admit minimal solutions for all $(a_+, a_-, b) \in \mathcal{H}$. (Indeed, this readily follows from Theorem II.3.) In particular, let us introduce the asymmetric weight function

$$w_\delta(a_+, a_-, b; z) \equiv \frac{G(z + ib - i(a_+ + a_-)/2) G(z + i(a_\delta - a_-)/2)}{G(z - ib + i(a_+ + a_-)/2) G(z - i(a_\delta - a_-)/2)}. \quad (5.13)$$

This function is related to w_r and w via

$$w_\delta(z) = w_r(z) \frac{\operatorname{sh}(\pi z/a_{-\delta})}{\operatorname{sh}(\pi z/a_\delta)} = w(z)/4\operatorname{sh}^2(\pi z/a_\delta) \quad (5.14)$$

on account of (3.5), (5.8) and (5.6). Since w solves (5.8), so does w_δ . Choosing the parameters in \mathcal{H}_δ (4.10), w_δ is a minimal solution, as is easily verified. Multiplying and/or dividing w_δ by finitely many factors of the form $s_\delta(z-c)$, one can construct explicit minimal solutions for arbitrary parameters.

We continue by obtaining analogs of the formulas (4.13)–(4.20). First, we use the AΔEs (3.4) to obtain

$$\frac{W(b + a_{-\delta}; z)}{W(b; z)} = 4\operatorname{sh} \frac{\pi}{a_\delta} (z + ib) \operatorname{sh} \frac{\pi}{a_\delta} (z - ib), \quad W = w, w_r, w_+, w_-. \quad (5.15)$$

Taking $k_+, k_- \in \mathbb{Z}$, these AΔEs can be iterated to yield

$$\begin{aligned} & \frac{W(b+k_+a_++k_-a_-;z)}{W(b;z)} \\ &= \prod_{\delta=+,-} \prod_{j_\delta=1}^{|k_\delta|} \left(4 \left(\operatorname{sh} \frac{\pi}{a_- \delta} \left(z + i \frac{k_\delta}{|k_\delta|} \left(b - \frac{a_\delta}{2} \right) \right. \right. \right. \\ & \quad \left. \left. \left. + i a_\delta \left(j_\delta - \frac{1}{2} \right) \right) \right) (i \rightarrow -i) \right)^{k_\delta/|k_\delta|}. \end{aligned} \quad (5.16)$$

Next, we note that (5.5) and (3.4) entail

$$w(a_+, a_-, 0; z) = 1, \quad (5.17)$$

$$w(a_+, a_-, a_\delta/2; z) = 2 \operatorname{th}(\pi z/a_\delta) \operatorname{sh}(\pi z/a_- \delta), \quad (5.18)$$

$$w(a_+, a_-, (a_+ + a_-)/2; z) = 4 \operatorname{sh}(\pi z/a_+) \operatorname{sh}(\pi z/a_-). \quad (5.19)$$

Therefore, the weight functions are elementary functions for parameters in the dense subset

$$\mathcal{D}_w \equiv \{(a_+, a_-, b) \in \mathcal{H} \mid b = l_+ a_+ + l_- a_-, l_+, l_- \in \mathbb{Z}/2\} \quad (5.20)$$

of \mathcal{H} (4.5). Specifically, one readily obtains from (5.16)–(5.19) (using the notation (4.19) and taking $k_+, k_- \in \mathbb{Z}$)

$$w(a_+, a_-, k_+ a_+ + k_- a_-; z) = \prod_{\delta=+,-} \prod_{j_\delta=1}^{|k_\delta|} \left(4 \left(\operatorname{sh} \frac{\pi}{a_- \delta} (z + i a_\delta (j_\delta - \theta(k_\delta))) \right) (i \rightarrow -i) \right)^{k_\delta/|k_\delta|}, \quad (5.21)$$

$$\begin{aligned} & w(a_+, a_-, a_\delta/2 + k_+ a_+ + k_- a_-; z) \\ &= 2 \operatorname{th} \left(\frac{\pi z}{a_\delta} \right) \operatorname{sh} \left(\frac{\pi z}{a_- \delta} \right) \prod_{j_\delta=1}^{|k_\delta|} \left(4 \left(\operatorname{sh} \frac{\pi}{a_- \delta} \left(z + i a_\delta \left(j_\delta - \frac{1}{2} \right) \right) \right) (i \rightarrow -i) \right)^{k_\delta/|k_\delta|} \\ & \quad \cdot \prod_{j_{-\delta}=1}^{|k_{-\delta}|} \left(4 \left(\operatorname{ch} \frac{\pi}{a_\delta} (z + i a_{-\delta} (j_{-\delta} - \theta(k_{-\delta}))) \right) (i \rightarrow -i) \right)^{k_{-\delta}/|k_{-\delta}|}, \end{aligned} \quad (5.22)$$

$$\begin{aligned} & w(a_+, a_-, (a_+ + a_-)/2 + k_+ a_+ + k_- a_-; z) \\ &= 4 \operatorname{sh} \left(\frac{\pi z}{a_+} \right) \operatorname{sh} \left(\frac{\pi z}{a_-} \right) \prod_{\delta=+,-} \prod_{j_\delta=1}^{|k_\delta|} \left(4 \left(\operatorname{ch} \frac{\pi}{a_- \delta} \left(z + i a_\delta \left(j_\delta - \frac{1}{2} \right) \right) \right) (i \rightarrow -i) \right)^{k_\delta/|k_\delta|}. \end{aligned} \quad (5.23)$$

We proceed by noting that none of the weight functions has the reflection symmetry (4.21) of the scattering function. Instead, one gets from (5.5) the relation

$$w_r(a_+ + a_- - b; z) = 1/w_r(b; z). \quad (5.24)$$

Combining this with (5.16), one obtains

$$w_r(-b; z) w_r(b; z) = \prod_{\delta=+,-} \left(4 \operatorname{sh} \frac{\pi}{a_\delta} (z + i b) \operatorname{sh} \frac{\pi}{a_\delta} (z - i b) \right)^{-1}. \quad (5.25)$$

Using the multiplication formula (3.25), one can work out analogs of the relations (4.23)–(4.26) for the weight functions. We shall not do so, however. We do point out that w_r satisfies an additional relation involving shifts of b —as opposed to shifts of z :

$$w_r\left(\frac{a_+}{M}, \frac{a_-}{N}, b; z\right) = \prod_{j=0}^{M-1} \prod_{k=0}^{N-1} w_r\left(a_+, a_-, b + \frac{a_+}{M}j + \frac{a_-}{N}k; z\right). \quad (5.26)$$

(Indeed, this formula readily follows from (5.5) and (3.25).)

By contrast to the scattering function, the weight functions are elementary functions on all of the sine-Gordon lines. In particular, from (5.6) and (5.18) we have

$$w(\pi, \alpha, \pi/2; z) = 2\text{th}z \text{sh}(\pi\alpha^{-1}z) \quad (\text{sG}) \quad (5.27)$$

for all $\alpha > 0$. (Compare this to (4.28).)

Next, we obtain an integral representation for w_r : From (3.1), (3.3) and (5.5) we have

$$w_r(z) = \exp(I(z)), \quad (5.28)$$

where

$$I(z) \equiv \int_0^\infty \frac{dy}{y} \left(\frac{\text{sh}(a_+ + a_- - 2b)y}{\text{sh}a_+ y \text{sh}a_- y} \cos 2yz - \frac{a_+ + a_- - 2b}{a_+ a_- y} \right). \quad (5.29)$$

This integral converges absolutely provided

$$|\text{Im}z| < e(a_+, a_-, b)/2, \quad (5.30)$$

where

$$e(a_+, a_-, b) \equiv a_+ + a_- - |2b - a_+ - a_-|. \quad (5.31)$$

Thus we have in particular

$$e(a_+, a_-, b) > a_\delta \Leftrightarrow (a_+, a_-, b) \in \mathcal{S}_\delta, \quad (5.32)$$

which says once more that w_r is regular for parameters in \mathcal{S}_δ .

More generally, the integral representation (5.28) sense and holds true in a strip around the real z -axis iff the parameters belong to \mathcal{C} (4.34). Indeed, one clearly has

$$e(a_+, a_-, b) > 0 \Leftrightarrow (a_+, a_-, b) \in \mathcal{C}. \quad (5.33)$$

Combining the representation with (5.6), (5.14) and (5.15), we obtain the positivity property

$$W(a_+, a_-, b; x) > 0, \quad \forall (a_+, a_-, b, x) \in \mathcal{H} \times \mathbb{R}^*, \quad W = w, w_r, w_+, w_-. \quad (5.34)$$

From (3.1) and (3.3) we also obtain an integral representation for the asymmetric weight function w_δ (5.13), viz.,

$$w_\delta(z) = \exp(I_\delta(z)) \quad (5.35)$$

with

$$I_\delta(z) \equiv 2 \int_0^\infty \frac{dy}{y} \left(\frac{\text{sh}(a_- - b)y \text{ch}(a_\delta - b)y}{\text{sh}a_+ y \text{sh}a_- y} \cos 2yz - \frac{a_- - b}{a_+ a_- y} \right). \quad (5.36)$$

Obviously, this integral has the same convergence properties as the integral (4.30), so the analysis embodied in (4.31)–(4.35) applies once again.

We have not found illuminating analogs of the representations (4.38) and (4.45), so we conclude this subsection by deriving two limits of the weight function w . (Corresponding limits for w_r, w_+ and w_- readily follow, so they will not be spelled out.) Once again, we switch to parameters (4.47).

First, we use (5.1), (3.22), (3.24) and (3.69) to obtain

$$w(\pi, \beta\nu, \beta\nu g; \beta p) = \exp(2g \ln(2\beta\nu)) \frac{H(\rho; p/\nu + i/2)H(\rho; -p/\nu + i/2)}{H(\rho; p/\nu - ig + i/2)H(\rho; -p/\nu - ig + i/2)}, \quad \rho \equiv \beta\nu/\pi. \quad (5.37)$$

Therefore, Prop. III.6 entails

$$\lim_{\beta \downarrow 0} (2\beta\nu)^{-2g} w(\pi, \beta\nu, \beta\nu g; \beta p) = \frac{\Gamma(ip/\nu + g)\Gamma(-ip/\nu + g)}{\Gamma(ip/\nu)\Gamma(-ip/\nu)} \quad (\text{I}_{\text{rel}} \text{ limit}), \quad (5.38)$$

where the limit is mero-uniform. (The limiting weight function is associated to the analytic difference operators of the I_{rel} regime, cf. Refs. 1 and 2.)

Second, we may write

$$w(\pi, a, ag; z) = \frac{G(\pi, a; z - i\pi/2 + ia(g - 1/2))}{G(\pi, a; z - i\pi/2 + ia(1/2))} \cdot \frac{G(\pi, a; z + i\pi/2 + ia(1/2))}{G(\pi, a; z + i\pi/2 + ia(1/2 - g))}. \quad (5.39)$$

Therefore, we deduce from (3.91)

$$\lim_{\beta \downarrow 0} w(\pi, \beta\nu, \beta\nu g; \nu x) = \exp(2g \ln(2 \operatorname{sh} \nu x)), \quad x \in \mathcal{R}_0 \quad (\text{II}_{\text{nr}} \text{ limit}) \quad (5.40)$$

(with \ln real-valued for $x > 0$), uniformly on compacts of \mathcal{R}_0 . (The limit is the weight function of the II_{nr} regime, cf. Refs. 1 and 2)

B. The elliptic case

The elliptic w -function is defined by replacing in (5.1) the hyperbolic G -functions by their elliptic counterparts. Obviously, this yields a function that is periodic in z with primitive period π/r , and which satisfies (5.2), (5.3), and (4.76), (4.77) with u replaced by w .

Just as in the hyperbolic case, we introduce a reduced weight function by (5.5). Then we obtain via (3.100) and (3.96)–(3.99)

$$w(z) = 4r^2 \prod_{k=1}^{\infty} (1 - q_+^{2k})^2 (1 - q_-^{2k})^2 \cdot s_+(z) s_-(z) w_r(z). \quad (5.41)$$

Evidently, w_r shares the automorphy properties of w mentioned above.

From Prop. III.11 we deduce that w and w_r are meromorphic in r, a_+, a_-, b and z , provided $a_+ r$ and $a_- r$ stay in the right half plane. As the analog of (5.7) we then obtain

$$u(iz; ib) = w_r(b; z) \frac{4r^2 \prod_{k=1}^{\infty} (1 - q_+^{2k})^2 (1 - q_-^{2k})^2 \cdot s_+(z + ib) s_-(z + ib)}{G(ib - i(a_+ - a_-)/2) G(ib + i(a_+ - a_-)/2)}. \quad (5.42)$$

From now on we take the parameters in \mathcal{E} (4.75). Turning to the AΔEs satisfied by w and w_r , we obtain once more

$$\frac{w(z+ia_\delta/2)}{w(z-ia_\delta/2)} = \frac{s_\delta(z+ib-ia_\delta/2)}{s_\delta(z-ib+ia_\delta/2)} \cdot \frac{s_\delta(z+ia_\delta/2)}{s_\delta(z-ia_\delta/2)}, \quad (5.43)$$

whereas (5.9) is replaced by

$$\frac{w_r(z+ia_\delta/2)}{w_r(z-ia_\delta/2)} = -\exp(2irz) \frac{s_\delta(z+ib-ia_\delta/2)}{s_\delta(z-ib+ia_\delta/2)}. \quad (5.44)$$

Considering first w_r , we reach the same conclusion as in the hyperbolic case—Prop. III.10 and π/r -periodicity in z play the role of Prop. III.3 and the asymptotics (5.11). Turning to $w(z)$, one readily sees that it generically has double zeros at $z=k\pi/r, k \in \mathbb{Z}$, and hence is not regular. The asymmetric function w_δ defined by (5.13) is now related to w_r and w via

$$w_\delta(z) = w_r(z) \prod_{k=1}^{\infty} \left(\frac{1-q_\delta^{2k}}{1-q_\delta^{-2k}} \right)^2 \cdot \frac{s_\delta(z)}{s_\delta(z)} = \frac{w(z)}{4r^2 \prod_{k=1}^{\infty} (1-q_\delta^{2k})^4 \cdot s_\delta(z)^2}. \quad (5.45)$$

Since $s_\delta(z)^2$ is not ia_δ -periodic, w_δ does not satisfy the AΔE (5.43), however. To obtain minimal periodic solutions to (5.43), one should rather multiply $w(z)$ by an elliptic function with periods π/r and ia_δ . We shall neither embark on this nor on a study of the AΔEs solved by the functions w_+ and w_- .

We continue by obtaining the counterparts of (5.15)–(5.19). First, from (5.1), (5.45) and (3.100) we readily get

$$\frac{W(b+a_\delta; z)}{W(b; z)} = 4r^2 e^{-2rb} \prod_{k=1}^{\infty} (1-q_\delta^{2k})^4 \cdot s_\delta(z+ib) s_\delta(z-ib), \quad W = w, w_r, w_+, w_-. \quad (5.46)$$

To obtain the analog of (5.16), we employ the relation

$$s(r, a; z_+ + ina) s(r, a; z_- - ina) = e^{-2irn(z_+ - z_-)} e^{2arn^2} s(r, a; z_+) s(r, a; z_-), \quad n \in \mathbb{N}, \quad (5.47)$$

which is easily derived from (2.91). (This formula plays the same role as (4.80) in simplifying the iterated AΔEs.) A straightforward calculation now yields (with $k_+, k_- \in \mathbb{Z}$)

$$\begin{aligned} \frac{W(b+k_+a_++k_-a_-; z)}{W(b; z)} &= \exp(2rb(2k_+k_- - k_+ - k_-)) \prod_{\delta=+, -} \exp(ra_\delta k_\delta (k_\delta - 1)(2k_-_\delta \\ &\quad - 1)) \prod_{j_\delta=1}^{|k_\delta|} \left(4r^2 \prod_{k=1}^{\infty} (1-q_\delta^{2k})^4 \left(s_\delta \left(z + i \frac{k_\delta}{|k_\delta|} \left(b - \frac{a_\delta}{2} \right) \right. \right. \right. \\ &\quad \left. \left. \left. + ia_\delta \left(j_\delta - \frac{1}{2} \right) \right) \right) (i \rightarrow -i) \right)^{k_\delta/|k_\delta|}. \end{aligned} \quad (5.48)$$

Next, we use (5.1) and (3.100) to obtain

$$w(r, a_+, a_-, 0; z) = 1, \quad (5.49)$$

$$w(r, a_+, a_-, a_\delta/2; z) = 4r^2 \prod_{k=1}^{\infty} (1-q_+^{2k})^2 (1-q_-^{2k})^2 \cdot \frac{s_\delta(z)}{R_\delta(z)} s_\delta(z), \quad (5.50)$$

$$w(r, a_+, a_-, (a_+ + a_-)/2; z) = 4r^2 \prod_{\delta=+,-} \prod_{k=1}^{\infty} (1 - q_{\delta}^{2k})^2 \cdot s_{\delta}(z). \quad (5.51)$$

If we now combine these formulas with the quotient formula (5.48), we obtain obvious analogs of (5.21)–(5.23)—which we do not spell out.

We proceed by observing that (5.24) holds true for the elliptic w_r , too. In tandem with (5.48), this entails

$$w_r(-b; z)w_r(b; z) = \prod_{\delta=+,-} \left(4r^2 \prod_{k=1}^{\infty} (1 - q_{\delta}^{2k})^4 \cdot s_{\delta}(z + ib)s_{\delta}(z - ib) \right)^{-1}. \quad (5.52)$$

Analogues of (4.23)–(4.26) for the elliptic weight functions are readily derived from the multiplication formula (3.25), so they will be skipped. The latter formula also entails that the elliptic w_r -function obeys (5.26).

As the elliptic counterpart of (5.27) we obtain from (5.50) and (5.41)

$$w(r, \pi, \alpha, \pi/2; z) = 4r^2 \prod_{k=1}^{\infty} (1 - e^{-2k\pi r})^2 (1 - e^{-2k\alpha r})^2 \cdot \frac{s(r, \pi; z)}{R(r, \pi; z)} s(r, \alpha; z) \quad (\text{XYZ}). \quad (5.53)$$

This holds true for all $\alpha > 0$, as opposed to the explicit formula (4.86), which holds for the dense set (4.27).

We now turn to deriving and studying a series representation for w_r . Recalling (3.3) and (3.92), the definition (5.5) entails

$$w_r(z) = \exp(S(z)) = \exp\left(\sum_{n=1}^{\infty} \frac{\text{sh}(a_+ + a_- - 2b)nr}{n \text{sha}_+ n r \text{sha}_- n r} \cos 2nrz \right). \quad (5.54)$$

The convergence properties of the infinite series $S(z)$ occurring here are the same as those of the integral $I(z)$ (5.29), so the analysis encoded in (5.30)–(5.33) applies verbatim. Using this representation, (5.46) and (5.45), we now deduce the positivity property

$$W(r, a_+, a_-, b; x) > 0, \quad \forall (r, a_+, a_-, b, x) \in \mathcal{E} \times (0, \pi/r), \quad W = w, w_r, w_+, w_-. \quad (5.55)$$

It is of interest to compare the series representation (5.54) to Theorem II.5. Choosing parameters in \mathcal{S}_{δ} , one deduces that Theorem II.5 applies to the additive version of (5.44), and that w_r corresponds to the unique minimal solution (2.106). Via (2.107) one can now obtain an integral representation for w_r —as an analog of the representation (4.89) for the elliptic u -function.

To conclude this subsection, we derive three limits of the w -function. First, we use Prop. III.12 to infer

$$\lim_{r \downarrow 0} \exp\left(\frac{\pi^2 b}{3ra_+ a_-} \right) w(r, a_+, a_-, b; z) = w_{\text{hyp}}(a_+, a_-, b; z) \quad (\text{II}_{\text{rel}} \text{ limit}), \quad (5.56)$$

where the limit is mero-uniform. (Here, w_{hyp} denotes the w -function from Subsection V A.) Note that the renormalizing exponential is necessary, and that no such factor occurs in the u -function counterpart (4.91).

Next, we claim that the limit

$$\lim_{A \uparrow \infty} w(r, a, A, b; z) = w_{\text{trig}}(r, a, b; z) \quad (\text{III}_{\text{rel}} \text{ limit}) \quad (5.57)$$

exists mero-uniformly. (Here, w_{trig} denotes the w -function studied in the next subsection.) Indeed, we may rewrite (5.1) as

$$w(r, a, A, b; z) = \frac{G(r, a, A; z + ib - ia/2 - iA/2) G(r, a, A; -z + ib - ia/2 - iA/2)}{G(r, a, A; z - ia/2 - iA/2) G(r, a, A; -z - ia/2 - iA/2)}, \quad (5.58)$$

so (3.139) yields the mero-uniform limit

$$\lim_{A \uparrow \infty} w(r, a, A, b; z) = \frac{G(r, a; z + ib - ia/2) G(r, a; -z + ib - ia/2)}{G(r, a; z - ia/2) G(r, a; -z - ia/2)}. \quad (5.59)$$

In view of (5.61) below, this entails (5.57).

Finally, fixing $g \in \mathbb{R}$, one has

$$\lim_{a \downarrow 0} w(r, A, a, ag; z) = \exp \left(2g \ln \left(2r \prod_{k=1}^{\infty} (1 - e^{-2kAr})^2 \cdot s(r, A; z) \right) \right), \quad z \in \mathcal{R}_r \quad (\text{IV}_{\text{nr}} \text{ limit}) \quad (5.60)$$

(with \ln real for $z \in (0, \pi/r)$), uniformly on compacts of \mathcal{R}_r (2.113). To check this, one need only substitute $b = ag$ in (5.58), invoke the limit (3.138), and recall (3.96)–(3.99).

C. The trigonometric case

The trigonometric w -function is defined by

$$w(r, a, b; z) \equiv \frac{G(z + ib - ia/2) G(-z + ib - ia/2)}{G(z - ia/2) G(-z - ia/2)} \quad (5.61)$$

with G given by (3.140). Thus, it can be written

$$w(r, a, b; z) = \prod_{n=0}^{\infty} \left(\frac{1 - q^{2n} e^{2irz}}{1 - q^{2n} e^{-2rb + 2irz}} \right) (z \rightarrow -z), \quad q \equiv e^{-ar}. \quad (5.62)$$

We note that w is π/r -periodic and even in z , and satisfies (4.103) and (4.104) with u replaced by w .

Next, we introduce the reduced weight function

$$w_r(z) \equiv G(z + ib - ia/2) G(-z + ib - ia/2), \quad (5.63)$$

which has the same automorphy properties as w . Recalling the functional equation (3.154) and AΔE (3.142) satisfied by the trigonometric G -function, one readily verifies that w_r and w are related by

$$w(z) = 4r \prod_{l=1}^{\infty} (1 - q^{2l})^2 \cdot s(r, a; z) \sin(rz) w_r(z). \quad (5.64)$$

Obviously, w_r and w are meromorphic in r, a, b and z , as long as ar stays in the right half plane. As the counterpart of (5.42) one easily gets

$$u(iz; ib) = 4r \prod_{l=1}^{\infty} (1 - q^{2l})^2 \cdot s(r, a; z + ib) \sin r(z + ib) \frac{G(-ib + ia/2)}{G(ib + ia/2)} w_r(b; z). \quad (5.65)$$

Taking from now on parameters in \mathcal{T} (4.102), we turn to the AΔEs solved by w and w_r , viz.,

$$\frac{w(z+ia/2)}{w(z-ia/2)} = \frac{\sin r(z+ib-ia/2)}{\sin r(z-ib+ia/2)} \cdot \frac{\sin r(z+ia/2)}{\sin r(z-ia/2)} \quad (5.66)$$

and

$$\frac{w_r(z+ia/2)}{w_r(z-ia/2)} = -\exp(2irz) \frac{\sin r(z+ib-ia/2)}{\sin r(z-ib+ia/2)}. \quad (5.67)$$

Clearly, both AΔEs are regular for arbitrary parameters. Choosing parameters in \mathcal{R} (4.106), one readily verifies that the reduced weight function is a minimal solution to (5.67) that is even and positive for $z \in \mathbb{R}$. As such, it is uniquely determined up to a positive constant, cf. Theorem II.1. For $b \leq a/2$, however, it has poles in the strip $|\operatorname{Im} z| \leq a/2$, so it is not regular. The weight function $w(z)$ has double zeros for $z = k\pi/r, k \in \mathbb{Z}$, unless $b = -na, n \in \mathbb{N}$; in the latter case one easily sees that w is a minimal solution to (5.66).

To proceed, we note that w and w_r satisfy the b -AΔE

$$\frac{W(b+a; z)}{W(b; z)} = 4e^{-2rb} \sin r(z+ib) \sin r(z-ib), \quad W = w, w_r. \quad (5.68)$$

Hence, iteration yields (with $k \in \mathbb{Z}$)

$$\frac{W(b+ka; z)}{W(b; z)} = e^{-2rbk - ark(k-1)} \prod_{j=1}^{|k|} \left(4 \left(\sin r \left(z + i \frac{k}{|k|} \left(b - \frac{a}{2} \right) + ia \left(j - \frac{1}{2} \right) \right) \right) (i \rightarrow -i) \right)^{k/|k|}. \quad (5.69)$$

Now from (5.61) we read off

$$w(r, a, 0; z) = 1, \quad (5.70)$$

so we deduce

$$w(r, a, ka; z) = e^{-ark(k-1)} \prod_{j=1}^{|k|} (4 [\sin r(z + ia(j - \theta(k)))]) [i \rightarrow -i]^{k/|k|}, \quad (5.71)$$

where $k \in \mathbb{Z}$ and the notation (4.19) is used. Moreover, from (3.154) we have

$$w_r(r, a, a/2; z) = R(r, a; z)^{-1}, \quad (5.72)$$

so recalling (5.64) we obtain (with $k \in \mathbb{Z}$)

$$\begin{aligned} w(r, a, a/2 + ka; z) &= 4r \prod_{l=1}^{\infty} (1 - q^{2l})^2 \cdot \frac{s(r, a; z)}{R(r, a; z)} \sin rz \\ &\quad \cdot \prod_{j=1}^{|k|} \left(4 \left(\sin r \left(z + ia \left(j - \frac{1}{2} \right) \right) \right) (i \rightarrow -i) \right)^{k/|k|}. \end{aligned} \quad (5.73)$$

Using the multiplication formula (3.145), one easily derives analogs of (4.112) and (4.113) for the weight functions. In addition, (3.145) entails that w_r satisfies

$$w_r\left(r, \frac{a}{M}, b; z\right) = \prod_{k=0}^{M-1} w_r\left(r, a, b + \frac{a}{M}k; z\right). \quad (5.74)$$

Next, we use (3.141) to obtain a series representation for w_r , namely

$$w_r(z) = \exp\left(\sum_{n=1}^{\infty} \frac{e^{nr(a-2b)}}{n \operatorname{sh} nra} \cos 2nrz\right). \quad (5.75)$$

Provided $b > 0$, this representation makes sense and holds true for $|\operatorname{Im} z| < b$. In particular, this entails once more that w_r is a minimal solution to (5.67) when the parameters belong to \mathcal{R} (4.106). (More specifically, w_r amounts to the unique minimal solution given by (2.106).) Furthermore, using (5.68) and (5.64) one deduces

$$W(r, a, b; x) > 0, \quad \forall (r, a, b, x) \in \mathcal{T} \times (0, \pi/r), \quad W = w, w_r. \quad (5.76)$$

We finish this subsection by obtaining two limits of the trigonometric weight function w . Recalling (3.155), we rewrite (5.61) with $a = 1$ as

$$w(r, 1, b; z) = \frac{T(r; -z + i/2)T(r; z + i/2)}{T(r; -z - ib + i/2)T(r; z - ib + i/2)} \exp(rb(1-b) + 2b \ln(2r)). \quad (5.77)$$

From Prop. III.20 we now infer

$$\lim_{r \downarrow 0} (2r)^{-2g} w(r, 1, g; z) = \frac{\Gamma(-iz + g)\Gamma(iz + g)}{\Gamma(-iz)\Gamma(iz)} \quad (\text{I}_{\text{rel}} \text{ limit}), \quad (5.78)$$

where the limit is mero-uniform. (Compare this to (5.38).)

Next, we substitute $b = ag$, with $g \in \mathbb{R}$ fixed, in (5.61). Recalling then the limit (3.180), we deduce

$$\lim_{a \downarrow 0} w(r, a, ag; z) = \exp(2g \ln(2 \sin rz)), \quad z \in \mathcal{R}_r \quad (\text{III}_{\text{nr}} \text{ limit}) \quad (5.79)$$

(with \ln real-valued for $z \in (0, \pi/r)$), where the limit is uniform on compact subsets of the period strip \mathcal{R}_r (2.113).

ACKNOWLEDGMENT

This work was partially supported by the Netherlands Organisation for the Advancement of Research (NWO).

APPENDIX A: THE GAMMA FUNCTION AND RELATED INTEGRALS

This appendix serves a twofold purpose. First of all, it is included to render this paper more self-contained. Indeed, most of the Laplace, sine and cosine transforms we derive below can be found—without proof—in standard sources such as Refs. 18 and 19; moreover, all of the properties of the psi and gamma functions we need can be found—with detailed proofs—in various sources, for instance Ref. 16. Our second purpose, however, is to demonstrate how these properties can be very quickly derived via the minimal solution (2.26) to a suitable AΔE (2.22); this yields a paradigm for the study of generalized psi and gamma functions undertaken in Section III.

Specifically, our starting point is the AΔE

$$F(z+i/2)-F(z-i/2)=\frac{i}{z-i/2}\equiv\chi(z). \quad (\text{A1})$$

A contour integration yields

$$\hat{\chi}(y)=\frac{1}{2\pi}\int_{-\infty}^{\infty}dx\frac{i}{x-i/2}e^{ixy}=-e^{-y/2}\theta(y), \quad (\text{A2})$$

so this AΔE is of the type considered in the proof of Theorem II.3. Indeed, (A2) entails

$$\hat{\phi}(y)=iy e^{-y/2}\theta(y), \quad \phi(z)\equiv\chi'(z)=-i(z-i/2)^{-2}, \quad (\text{A3})$$

and therefore $\phi(z)$ has all of the properties (2.18)–(2.21). From Theorem II.2 we now obtain a solution

$$f(z)=2i\int_0^{\infty}dy\frac{ye^{-y}}{\text{sh}y}e^{-2iyz}, \quad \text{Im}z<1, \quad (\text{A4})$$

to the AΔE (2.22), which is the uniquely determined solution with properties (2.23)–(2.25).

As a consequence, the function

$$F_1(z)=F_1(0)+c_1z+\int_0^{\infty}dy\frac{e^{-y}}{\text{sh}y}(1-e^{-2iyz}) \quad (\text{A5})$$

is a solution to (A1) for a certain $c_1 \in \mathbb{C}$. Now we have

$$F_1(i/2)-F_1(-i/2)=ic_1+\int_0^{\infty}dy\frac{e^{-y}}{\text{sh}y}(-e^y+e^{-y})=ic_1-2. \quad (\text{A6})$$

Hence, noting $\chi(0)=-2$, we need $c_1=0$ to solve (A1). Of course, we are free to choose $F_1(0)$, and we shall set

$$F_1(0)=\int_0^{\infty}dy\left(\frac{e^{-2y}}{y}-\frac{e^{-y}}{\text{sh}y}\right)\equiv-\gamma. \quad (\text{A7})$$

(As will soon become clear, γ is Euler's constant.) The upshot is that we obtain a solution

$$F_1(z)\equiv\int_0^{\infty}dy\left(\frac{e^{-2y}}{y}-\frac{e^{-y(1+2iz)}}{\text{sh}y}\right), \quad \text{Im}z<1, \quad (\text{A8})$$

to the AΔE (A1). Note that the function $F_2(z)\equiv F_1(-z+i)$ yields a second solution to (A1), so that $F_1(z)-F_1(-z+i)$ is an i -periodic meromorphic function (determined explicitly below).

Next, we observe that the AΔE (2.22), with $\phi(z)$ given by (A3), can also be solved by downward iteration, yielding the solution

$$\tilde{f}(z)=-i\sum_{k=1}^{\infty}(z-ik)^{-2}. \quad (\text{A9})$$

Now this solution clearly has the properties (2.23)–(2.25), so we must have $\tilde{f}(z)=f(z)$. From this we readily deduce

$$F_1(z) = -\gamma + i \sum_{k=1}^{\infty} \left(\frac{1}{z-ik} + \frac{1}{ik} \right). \quad (\text{A10})$$

(Indeed, the function on the rhs has derivative $\tilde{f}(z) = f(z)$ and value $-\gamma$ for $z=0$, just as $F_1(z)$, (A7) and (A8).) As a consequence, we obtain the functional equation

$$F_1(z+i/2) - F_1(-z+i/2) = - \sum_{n=0}^{\infty} \left(\frac{1}{iz+n+1/2} + \frac{1}{iz-n-1/2} \right) = i\pi \operatorname{th} \pi z. \quad (\text{A11})$$

Note that the rhs amounts to the i -periodic meromorphic function mentioned below (A8).

We are now prepared to make contact with the psi and gamma functions. First, we introduce

$$\psi(z) \equiv F_1(-iz+i) = \int_0^{\infty} dy \left(\frac{e^{-2y}}{y} - \frac{e^{y(1-2z)}}{\operatorname{sh} y} \right), \quad \operatorname{Re} z > 0. \quad (\text{A12})$$

Then we obtain from (A1) and (A11) the $\Delta\Delta E$

$$\psi(z+1) - \psi(z) = 1/z \quad (\text{A13})$$

and functional equation

$$\psi(z+1/2) - \psi(-z+1/2) = \pi \operatorname{tg} \pi z. \quad (\text{A14})$$

Moreover, we have $\psi(1) = -\gamma$ and $\psi(z)$ has simple poles at $z=0, -1, -2, \dots$, cf. (A10).

Next, consider any primitive $\Psi(z)$ of $\psi(z)$, restricted to the cut plane

$$\mathbb{C}^- \equiv \{z \in \mathbb{C} \mid z \notin (-\infty, 0]\}. \quad (\text{A15})$$

Clearly, $\Psi(z)$ is analytic in \mathbb{C}^- and satisfies

$$\Psi(z+1) - \Psi(z) = \ln z + c_1, \quad z \in \mathbb{C}^-, \quad (\text{A16})$$

$$\Psi(z+1/2) + \Psi(-z+1/2) = -\ln(\cos \pi z) + c_2, \quad \pm z \notin [1/2, \infty), \quad (\text{A17})$$

in view of (A13) and (A14). Now from (A12) we have

$$\Psi(2) - \Psi(1) = \int_1^2 dw \psi(w) = \int_0^{\infty} dy \left(\frac{e^{-2y}}{y} + \frac{e^y}{2y \operatorname{sh} y} (e^{-4y} - e^{-2y}) \right) = 0, \quad (\text{A18})$$

so that $c_1 = 0$ in (A16). Clearly, c_2 in (A17) depends on the arbitrary constant in $\Psi(z)$; we render Ψ unique by requiring $2\Psi(1/2) = \ln \pi$ and then we get $c_2 = \ln \pi$ by taking $z=0$ in (A17).

The upshot is that we obtain a primitive $\Psi(z)$ of $\psi(z)$ satisfying

$$\Psi(z+1) - \Psi(z) = \ln z, \quad (\text{A19})$$

$$\Psi(z+1/2) + \Psi(-z+1/2) = \ln(\pi / \cos \pi z). \quad (\text{A20})$$

Introducing the function

$$\Gamma(z) \equiv \exp(\Psi(z)) \quad (\text{A21})$$

(defined at first in \mathbb{C}^-), it readily follows that $\Gamma(z)$ extends to a meromorphic function without zeros and with simple poles at $z=0, -1, -2, \dots$. Indeed, from (A10) and (A12) we deduce that we have

$$\Psi(z) = \alpha - \gamma z - \ln z - \sum_{n=1}^{\infty} \left(\ln \left(1 + \frac{z}{n} \right) - \frac{z}{n} \right) \quad (\text{A22})$$

for some $\alpha \in \mathbb{C}$ (with $\ln z$ real for $z > 0$, of course). Therefore, we obtain

$$\frac{1}{\Gamma(z)} = e^{-\alpha + \gamma z} z \prod_{n=1}^{\infty} \left(1 + \frac{z}{n} \right) e^{-z/n} \quad (\text{A23})$$

and from this the assertion is clear. From (A19) and (A20) we also obtain the AΔE

$$\Gamma(1+z) = z\Gamma(z) \quad (\text{A24})$$

and functional equation

$$\Gamma(z+1/2)\Gamma(-z+1/2) = \pi / \cos \pi z. \quad (\text{A25})$$

In order to determine α , we note that (A23) and (A24) entail

$$e^{-\alpha} = \lim_{z \rightarrow 0} \frac{1}{z\Gamma(z)} = \frac{1}{\Gamma(1)}. \quad (\text{A26})$$

Now from (A24) and (A25) we have

$$\Gamma(z+1/2)\Gamma(-z+3/2) = \frac{\pi(-z+1/2)}{\cos \pi z}, \quad (\text{A27})$$

which yields $\Gamma(1)^2 = 1$ for $z \rightarrow 1/2$. Thus we conclude

$$\Gamma(1) = 1, \quad \alpha = 0, \quad (\text{A28})$$

since $\Gamma(z)$ is positive for $z > 0$. (To see this, note that (A12) entails $\psi(z)$ is real for $z > 0$. As $\Psi(1/2)$ is real, it follows that $\Psi(z)$ is real for $z > 0$, so positivity is clear from (A21).)

Combining (A23) and (A28), we see that $\Gamma(z)$ is the customary gamma function in Weierstrass product form, as anticipated by our notation. Similarly, $\psi(z)$ is the usual psi function (the logarithmic derivative of the gamma function), and (A12) amounts to Gauss' formula, cf., e.g., Ref. 16.

We now derive a number of definite integrals by exploiting the properties of $\psi(z)$ and $\Gamma(z)$ established above. The order in which this is done is determined by the order in which these integrals are needed in the main text, except when logical necessity requires otherwise.

First, we use the well-known integral

$$\int_0^{\infty} \frac{dy}{y} (e^{-qy} - e^{-py}) = \int_0^{\infty} dy \int_q^p ds e^{-sy} = \int_q^p \frac{ds}{s} = \ln(p/q) \quad (\text{A29})$$

and (A12) to obtain

$$\psi(z+1/2) - \ln z = \int_0^{\infty} dy \left(\frac{1}{y} - \frac{1}{\sinh y} \right) e^{-2yz}, \quad \operatorname{Re} z > 0. \quad (\text{A30})$$

Integrating this from 0 to z and using $2\Psi(1/2) = \ln \pi$, we arrive at

$$\Psi\left(z + \frac{1}{2}\right) - \frac{1}{2} \ln \pi - z \ln z + z = \frac{1}{2} \int_0^\infty \frac{dy}{y} \left(\frac{1}{y} - \frac{1}{\operatorname{sh} y} \right) (1 - e^{-2yz}), \quad \operatorname{Re} z > 0. \quad (\text{A31})$$

Now the function on the lhs is analytic in \mathbb{C}^- and the integral on the rhs converges absolutely for $\operatorname{Re} z \geq 0$. Thus, (A31) holds true for $\operatorname{Re} z = 0$, too. Putting $z = ix$ and $z = -ix$, $x \in \mathbb{R}$, in (A31), and taking the sum of the resulting equations, we obtain using (A20)

$$\ln(\pi / \operatorname{ch} \pi x) - \ln \pi + \pi x = C - \int_0^\infty \frac{dy}{y} \left(\frac{1}{y} - \frac{1}{\operatorname{sh} y} \right) \cos 2yx, \quad (\text{A32})$$

where we have set

$$C \equiv \int_0^\infty \frac{dy}{y} \left(\frac{1}{y} - \frac{1}{\operatorname{sh} y} \right). \quad (\text{A33})$$

If we now take $x \rightarrow \infty$ in (A32), then the integral has limit 0 (by virtue of the Riemann-Lebesgue lemma), so we must have

$$C = \ln 2. \quad (\text{A34})$$

Combining this with (A31) and (A21), we obtain the integral representation

$$\Gamma(z + 1/2) = (2\pi)^{1/2} \exp\left(z \ln z - z - \frac{1}{2} \int_0^\infty \frac{dy}{y} \left(\frac{1}{y} - \frac{1}{\operatorname{sh} y} \right) e^{-2yz}\right), \quad (\text{A35})$$

which holds true for $\operatorname{Re} z \geq 0$.

Next, we put $q = 2, p = 2w$ in (A29) and integrate w.r.t. w from 0 to z to obtain the identity

$$z \ln z - z = \int_0^\infty \frac{dy}{y} \left(e^{-2yz} + \frac{e^{-2yz} - 1}{2y} \right). \quad (\text{A36})$$

Inserting this in (A35), we get the representation

$$\Gamma(z + 1/2) = (2\pi)^{1/2} \exp\left(\int_0^\infty \frac{dy}{y} \left(z e^{-2y} - \frac{1}{2y} + \frac{e^{-2yz}}{2 \operatorname{sh} y} \right)\right), \quad (\text{A37})$$

which is valid for $\operatorname{Re} z > -1/2$. A routine calculation using (A37) and (A29) (with $q = 2, p = 2w$) now yields

$$\frac{\Gamma(w + \lambda)}{\Gamma(w + \mu)} e^{(\mu - \lambda) \ln w} = \exp\left(\int_0^\infty \frac{dt}{t} e^{-wt} \left(\lambda - \mu + \frac{e^{-\lambda t} - e^{-\mu t}}{1 - e^{-t}} \right)\right), \quad (\text{A38})$$

which holds true for $\operatorname{Re} w > \max(0, -\operatorname{Re} \lambda, -\operatorname{Re} \mu)$. Therefore, the function

$$F(w, \lambda) \equiv \frac{w + \lambda}{w - \lambda} \left(\frac{\Gamma(w + \lambda)}{\Gamma(w - \lambda)} e^{-2\lambda \ln w} \right)^2 \quad (\text{A39})$$

admits the representation

$$F(w, \lambda) = \exp \left(2 \int_0^\infty \frac{dt}{t} e^{-wt} (2\lambda - \operatorname{sh} \lambda t \operatorname{cth} t/2) \right), \quad (\text{A40})$$

provided $\operatorname{Re} w > |\operatorname{Re} \lambda|$. (To check this, use (A38) and (A29) with $q = w - \lambda$ and $p = w + \lambda$.)

The function $F(w, \lambda)$ will reappear in Appendix B; it is crucial for obtaining Prop. III.5 in Subsection III A. We conclude by deriving some formulas that are used towards the end of Subsection IV A. First, (A12) entails the cosine transform

$$\begin{aligned} & \psi((p+1+ix)/2) - \psi((q+1+ix)/2) + (x \rightarrow -x) \\ &= 2 \int_0^\infty \frac{dy}{\operatorname{sh} y} (e^{-qy} - e^{-py}) \cos xy, \quad \operatorname{Re} p, \operatorname{Re} q > -1, \quad x \in \mathbb{R} \end{aligned} \quad (\text{A41})$$

Now we take $\operatorname{Re} p \in (-1, 1)$ and put $q = -p$. Using (A14) and the elementary identity

$$\operatorname{tg}(\sigma + i\tau) + \operatorname{tg}(\sigma - i\tau) = \frac{2 \sin 2\sigma}{\cos 2\sigma + \operatorname{ch} 2\tau}, \quad \sigma, \tau \in \mathbb{C}, \quad (\text{A42})$$

we obtain

$$\int_0^\infty dy \frac{\operatorname{sh} py}{\operatorname{sh} y} \cos xy = \frac{\pi}{2} \frac{\sin \pi p}{\cos \pi p + \operatorname{ch} \pi x}, \quad |\operatorname{Re} p| < 1, \quad x \in \mathbb{R}. \quad (\text{A43})$$

Integrating this with respect to p from s to t yields

$$2 \int_0^\infty \frac{dy}{y} \frac{(\operatorname{ch} ty - \operatorname{ch} sy)}{\operatorname{sh} y} \cos xy = \ln \left(\frac{\operatorname{ch} \pi x + \cos \pi s}{\operatorname{ch} \pi x + \cos \pi t} \right), \quad |\operatorname{Re} s|, |\operatorname{Re} t| < 1. \quad (\text{A44})$$

Finally, we integrate (A41) w.r.t. x from 0 to $-2is$ and put $p = t + \lambda, q = t - \lambda$. The resulting formula entails the identity

$$\begin{aligned} & \frac{\Gamma(s + (1 + \lambda + t)/2) \Gamma(-s + (1 - \lambda + t)/2)}{(s \rightarrow -s)} = \exp \left(2 \int_0^\infty \frac{dy}{y} \frac{\operatorname{sh} \lambda y \operatorname{sh} 2sy}{\operatorname{sh} y} e^{-ty} \right), \\ & \operatorname{Re} t - |\operatorname{Re} \lambda| > -1, \quad s \in i\mathbb{R}. \end{aligned} \quad (\text{A45})$$

APPENDIX B: UNIFORM ESTIMATES

The main goal of this appendix consists in deriving bounds that are sufficiently strong to control the convergence and meromorphy properties of infinite products involving gamma functions, which occur in the main text. Our tool for doing so is Theorem B.1, which deals with Laplace transforms $L(w), w \in \mathbb{C}$, of a certain type. More generally, this theorem can be used to obtain estimates on remainders in asymptotic expansions that hold uniformly in sectorial regions $|\operatorname{Arg} w| \leq \pi - \epsilon, |w| \geq K = K(\epsilon)$ for any $\epsilon > 0$. As such, it is inspired by, but simpler than, the methods that can be found in Ref. 20, Sections 21–25, and Ref. 16, Section 13.6.

Assume $h(z)$ is a function that is analytic in the right half plane $\operatorname{Re} z > 0$ and at $z = 0$. Moreover, assume $h(z)$ satisfies the bound

$$|h(te^{i\phi})| \leq C(\chi) e^{rt}, \quad \forall (t, \phi) \in [0, \infty) \times [-\chi, \chi], \quad (\text{B1})$$

where $\chi \in [0, \pi/2)$ and $r \in [0, \infty)$, and where $C(\chi)$ is a positive non-decreasing function on $[0, \pi/2)$.

Theorem B.1: *The function*

$$L(w) \equiv \int_0^\infty e^{-wt} h(t) dt \quad (\text{B2})$$

is well defined and analytic in $\{\operatorname{Re} w > r\}$. Furthermore, $L(w)$ can be continued to a function that is analytic in

$$U_r \equiv \{\operatorname{Re} w \geq 0, |w| > r\} \cup \{\operatorname{Re} w < 0, |\operatorname{Im} w| > r\}. \quad (\text{B3})$$

Finally, fixing $\chi \in [0, \pi/2)$ and $R > r$ one has

$$|L(w)| \leq C(\chi)(R-r)^{-1}, \quad \forall w \in S_{R,\chi} \quad (\text{B4})$$

where

$$S_{R,\chi} \equiv \bigcup_{|\phi| \leq \chi} \{\operatorname{Re}(e^{i\phi} w) \geq R\}. \quad (\text{B5})$$

Proof: The first assertion is obvious. To prove the second one, consider the integral

$$e^{i\chi} \int_0^\infty \exp(-wte^{i\chi}) h(te^{i\chi}) dt, \quad \chi \in (-\pi/2, \pi/2). \quad (\text{B6})$$

Due to the bound (B1) this defines a function $L_\chi(w)$ that is analytic in the region

$$U_{r,\chi} \equiv \{\operatorname{Re}(e^{i\chi} w) > r\}. \quad (\text{B7})$$

We claim that $L_\chi(w)$ equals $L(w)$ in $U_{r,0} \cap U_{r,\chi}$. Taking this for granted, the second assertion follows, since we have

$$U_r = \bigcup_{|\chi| < \pi/2} U_{r,\chi}. \quad (\text{B8})$$

To prove the claim we first take $\chi \in [0, \pi/2)$. Fixing $w \in U_{r,0} \cap U_{r,\chi}$, we then have

$$\inf_{\phi \in [0, \chi]} \{\operatorname{Re}(e^{i\phi} w)\} = \min(\operatorname{Re} w, \operatorname{Re}(e^{i\chi} w)) = r + \epsilon \quad (\text{B9})$$

with $\epsilon = \epsilon(w) > 0$. Using (B1) we now obtain

$$|\exp(-wte^{i\phi}) h(te^{i\phi})| \leq C(\chi) e^{-\epsilon t}, \quad \forall (\phi, t) \in [0, \chi] \times [0, \infty). \quad (\text{B10})$$

This bound entails that the integral of $e^{-wz} h(z)$ over the contour $z = Ke^{i\phi}$, $\phi \in [0, \chi]$, vanishes for $K \rightarrow \infty$. Thus we may replace the contour $te^{i\chi}$, $t \in [0, \infty)$, in the z -plane by the positive real axis, yielding $L_\chi(w) = L(w)$. This proves our claim for non-negative χ , and the same reasoning applies to negative χ .

It remains to prove (B4). To this end we fix $w \in S_{R,\chi}$. In view of (B5) we can find $\phi \in [-\chi, \chi]$ such that $\operatorname{Re}(we^{i\phi}) \geq R$. Then we get

$$|L(w)| = |L_\phi(w)| \leq \int_0^\infty |\exp(-wte^{i\phi}) h(te^{i\phi})| dt \leq C(\chi) \int_0^\infty e^{-Rt} e^{rt} dt = C(\chi)(R-r)^{-1}, \quad (\text{B11})$$

where we used (B1). Thus (B4) holds true. \square

To illustrate how this result can be applied, we consider the Laplace transform

$$\mathcal{L}(w) \equiv \int_0^\infty e^{-wt} f(t) dt, \quad (\text{B12})$$

$$f(t) \equiv \frac{1}{t} - \frac{1}{\text{sh}t}, \quad (\text{B13})$$

occurring on the rhs of (A30). Integrating by parts n times, we obtain

$$\mathcal{L}(w) - \sum_{l=1}^n w^{-l} f^{(l-1)}(0) = w^{-n} \int_0^\infty e^{-wt} f^{(n)}(t) dt. \quad (\text{B14})$$

Now the function $h(t) \equiv f^{(n)}(t)$ satisfies the assumptions of Theorem B.1 with $r=0$, so (B4) yields a bound on the remainder integral that is uniform in $S_{R,\chi}$; fixing $\delta > 0$, the sectorial region $|\text{Arg} w| \leq \pi/2 + \chi - \delta, |w| \geq K$, belongs to $S_{R,\chi}$ for $K = K(\delta, R, \chi)$ large enough, cf. Fig. 1.

The Laplace transform in (A35) can be handled in the same way. This yields an asymptotic expansion that is substantially equivalent to the Stirling series, valid uniformly in sectorial regions of the above type.

For applications in the main text, however, we shall exploit Theorem B.1 to obtain uniform estimates pertaining to the Laplace transforms

$$\mathcal{L}_j(w) = \int_0^\infty e^{-wt} f_j(t) dt, \quad j = 1, 2, 3, \quad (\text{B15})$$

with

$$f_1 \equiv \frac{1}{t} (2\lambda - \text{sh}\lambda t \text{cth}t/2) \Rightarrow f_1(0) = 0, \quad f_1'(0) = -\lambda(2\lambda^2 + 1)/6, \quad (\text{B16})$$

$$f_2 \equiv \frac{1}{t} \left(\lambda - \mu + \frac{e^{-\lambda t} - e^{-\mu t}}{1 - e^{-t}} \right) \Rightarrow f_2(0) = (\lambda - \mu)(\lambda + \mu - 1)/2, \quad (\text{B17})$$

$$f_3 \equiv \frac{\text{sh}\lambda t \text{sh}\mu t \text{sh}\kappa t}{t \text{sh}t} \Rightarrow f_3(0) = 0, \quad f_3'(0) = \lambda \mu \kappa. \quad (\text{B18})$$

Then the functions $h_1 \equiv f_1'', h_2 \equiv f_2'$ and $h_3 \equiv f_3''$ satisfy the hypotheses of Theorem B.1. Correspondingly, we deduce the bounds

$$\left| \mathcal{L}_1(w, \lambda) + \frac{\lambda(2\lambda^2 + 1)}{6w^2} \right| \leq \frac{C_1(\chi, \lambda)}{|w^2|(R - r_1)}, \quad r_1 \equiv |\lambda|, \quad (\text{B19})$$


$$\left| \mathcal{L}_2(w, \lambda, \mu) - \frac{(\lambda - \mu)(\lambda + \mu + 1)}{2w} \right| \leq \frac{C_2(\chi, \lambda, \mu)}{|w|(R - r_2)}, \quad r_2 \equiv \max(|\lambda|, |\mu|), \quad (\text{B20})$$

$$\left| \mathcal{L}_3(w, \lambda, \mu, \kappa) - \frac{\lambda \mu \kappa}{w^2} \right| \leq \frac{C_3(\chi, \lambda, \mu, \kappa)}{|w^2|(R - r_3)}, \quad r_3 \equiv |\lambda| + |\mu| + |\kappa|, \quad (\text{B21})$$

which hold true for $R > r_j$ and all $w \in S_{R,\chi}$. The functions C_j are positive and non-decreasing in χ for fixed values of the parameters, and they are continuous in the parameters for fixed χ .

Recalling (A40), one easily obtains a corresponding bound on

$$F(w, \lambda) = \exp(2\mathcal{L}_1(w, \lambda)). \quad (\text{B22})$$

FIG. 1. The region $S_{R, \chi}$ and the complement of the region U_r .

We will need this bound in Subsection III A. Similarly, from (A38) one has

$$\frac{\Gamma(w+\lambda)}{\Gamma(w+\mu)} e^{(\mu-\lambda)\ln w} = \exp(\mathcal{L}_2(w, \lambda, \mu)), \quad (\text{B23})$$

and the bound on the lhs following from (B20) will be used several times in Subsection IV A. For the applications of (B19) and (B20) we do not need a bound on the parameter dependence of C_1 and C_2 ; continuity in the parameters suffices. As concerns (B21), however, it is important to

have more information on C_3 . Indeed, in Subsection IV A we shall use (B21) on four occasions; in two cases the parameters vary over \mathbb{C} -compacts, but in the remaining applications one or two parameters go to infinity.

In order to control this divergence, we first note that the function

$$h(t, p) \equiv \frac{\operatorname{sh} pt}{t} \quad (\text{B24})$$

satisfies the bounds

$$|\partial_t^j h(te^{i\phi}, p)| \leq d_j(\chi) |p|^{j+1} \exp(|p|t), \quad \forall (t, \phi, p) \in [0, \infty) \times [-\chi, \chi] \times \mathbb{C}, \quad (\text{B25})$$

with d_j positive non-decreasing functions on $[0, \pi/2)$, and $j=0, 1, 2$. (Write h as $pf(pt)$, $f(x) \equiv \operatorname{sh} x/x$, to verify this.) Factorizing f_3 accordingly, we deduce that the function C_3 in the bound (B1) on f_3'' satisfies

$$|C_3(\chi, \lambda, \mu, \kappa)| \leq d(\chi) |\lambda \mu| (|\lambda|^2 + |\mu|^2 + |\kappa|^2 + |\lambda \mu| + |\lambda \kappa| + |\mu \kappa|) \quad (\text{B26})$$

with d positive and non-decreasing on $[0, \pi/2)$. This bound on the parameter dependence is sufficient for our purposes.

¹S. N. M. Ruijsenaars, "Finite-dimensional soliton systems," in *Integrable and superintegrable systems*, edited by B. Kupershmidt (World Scientific, Singapore, 1990), pp. 165–206.

²S. N. M. Ruijsenaars, "Systems of Calogero-Moser type," to be published in *Proceedings of the 1994 Banff summer school 'Particles and fields'*, edited by G. Semenov and L. Vinet.

³J. Thomae, "Beiträge zur Theorie der durch die Heinesche Reihe darstellbaren Funktionen," *J. Reine Angew. Math.* **70**, 258–281 (1869).

⁴G. Gasper and M. Rahman, *Basic hypergeometric series* (Cambridge U.P., Cambridge, 1990).

⁵N. E. Nörlund, *Vorlesungen über Differenzenrechnung* (Springer, Berlin, 1924).

⁶L. M. Milne-Thomson, *The calculus of finite differences* (Macmillan, London, 1933).

⁷H. Meschkowski, *Differenzgleichungen* (Vandenhoeck & Ruprecht, Göttingen, 1959).

⁸P. Schroth, "Zur Definition der Nörlundschen Hauptlösung von Differenzgleichungen," *Manuscr. Math.* **24**, 239–251 (1978).

⁹L. Hörmander, *An introduction to complex analysis in several variables* (Van Nostrand, London, 1966).

¹⁰S. N. M. Ruijsenaars and H. Schneider, "A new class of integrable systems and its relation to solitons," *Ann. Phys. (N.Y.)* **170**, 370–405 (1986).

¹¹S. N. M. Ruijsenaars, "Complete integrability of relativistic Calogero-Moser systems and elliptic function identities," *Commun. Math. Phys.* **110**, 191–213 (1987).

¹²M. A. Olshanetsky and A. M. Perelomov, "Quantum integrable systems related to Lie algebras," *Phys. Repts.* **94**, 313–404 (1983).

¹³S. N. M. Ruijsenaars, "Generalized Lamé functions," to be published.

¹⁴A. B. Zamolodchikov and A. B. Zamolodchikov, "Factorized S -matrices in two dimensions as the exact solutions of certain relativistic quantum field theory models," *Ann. Phys. (N.Y.)* **120**, 253–291 (1979).

¹⁵R. J. Baxter, *Exactly solved models in statistical mechanics* (Academic, New York, 1982).

¹⁶E. T. Whittaker and G. N. Watson, *A course of modern analysis* (Cambridge U.P., Cambridge, 1933).

¹⁷S. N. M. Ruijsenaars, "Action-angle maps and scattering theory for some finite-dimensional integrable systems. III. Sutherland type systems and their duals," *Publ. RIMS, Kyoto Univ.* **31**, 247–353 (1995).

¹⁸A. Erdélyi, *Tables of integral transforms, Vol. 1* (McGraw-Hill, New York, 1954).

¹⁹I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products* (Academic, New York, 1965).

²⁰E. T. Copson, *Asymptotic expansions* (Cambridge U.P., Cambridge, 1965).