

Dispelling the myth of robotic efficiency


Ian Crawford explains why human space exploration will tell us more about the solar system than robotic exploration alone.

There is a widely held view in the astronomical community that unmanned robotic space vehicles are, and always will be, more efficient explorers of planetary surfaces than astronauts (e.g. Coates 2001, Clements 2009, Rees 2011). Partly this comes from a common assumption that robotic exploration is *cheaper* than human exploration (although this isn't necessarily true if like is compared with like) and partly from the expectation that developments in technology will relentlessly increase the capability and reduce the size and cost of robotic missions to the point that human exploration will not be able to compete. I argue below that the experience of human exploration during the Apollo missions, more recent field analogue studies and trends in robotic space exploration all point to exactly the opposite conclusion.

Benefits of human space exploration

As demonstrated by the Apollo missions 40 years ago – and leaving the question of cost to a separate examination below – human space exploration has a number of advantages over robotic operations on planetary surfaces. These have been discussed in detail elsewhere (e.g. Spudis 2001, Crawford 2001, 2010, Garvin 2004, Cockell 2004, Snook *et al.* 2007) and were endorsed by the independent Commission on the Scientific Case for Human Space Exploration commissioned by the RAS in 2005 (Close *et al.* 2005, hereinafter “the RAS Report”). These advantages can be summarized as follows:

- On-the-spot decision making and flexibility, with increased opportunities for serendipitous discoveries.
- Greatly enhanced mobility and attendant opportunities for geological exploration and instrument deployment. Compare the 35.7km traversed in three days by the Apollo 17 astronauts in December 1972 with the almost identical distance (34.4km) traversed by the Mars Exploration Rover Opportunity in eight years.
- Greatly increased efficiency in sample collection and sample return capacity. Compare the 382kg of samples returned by Apollo with the



1: The increasing size of Mars rovers, from Pathfinder (front left), a Mars Exploration Rover (left), to Mars Science Laboratory (right). This increase in size (and cost), contrary to predictions that improved technology will result in smaller and cheaper robots, is mandated by the nature of the martian surface and complexity of exploration objectives. Human missions would be even larger and more expensive, but, crucially, much more capable. (NASA/JPL-Caltech)

0.32 kg from the Russian robotic sample return missions Lunas 16, 20 and 24, and the zero kg returned so far by any robotic mission to Mars.

- Increased potential for large-scale exploratory activities (e.g. drilling) and the deployment and maintenance of complex equipment.
 - The development of a space-based infrastructure to support space-based astronomy and other scientific applications (e.g. the construction and maintenance of large space telescopes).
- With the exception of the final point, for which the best demonstration is provided by the five successful space shuttle servicing missions to the Hubble Space Telescope (e.g. NRC 2005), demonstration of the benefits of human spaceflight for planetary exploration must be sought in a comparison of the relative efficiencies of the Apollo missions and robotic missions to the Moon and Mars, supported where appropriate with terrestrial analogue studies.

The relative efficiency of human over robotic exploration of planetary surfaces is well recognized by scientists directly involved with the latter. For example, regarding the exploration of Mars, the RAS Report (paragraph 70) noted:

“The expert evidence we have heard strongly suggests that the use of autonomous robots alone will very significantly limit what can be learned about our nearest potentially habitable planet.”

Putting it more bluntly, Steve Squyres, the PI

for the Mars Exploration Rovers (MERs) Spirit and Opportunity, has written:

“[t]he unfortunate truth is that most things our rovers can do in a perfect sol [i.e. a martian day] a human explorer could do in less than a minute” (Squyres 2005 p234–5).

This is of course only a qualitative assessment, albeit by someone well placed to make an informed judgement. Nevertheless, at face value it implies a human/robot efficiency ratio of about 1500, which is far larger than the likely ratio of cost between a human mission to Mars and the cost of the MERs (see below). Even this, however, does not fairly compare human exploration efficiency with robotic exploration. This is because much of the scientific benefit of human missions will consist of samples returned, drill cores drilled and geophysical instruments deployed, all of which were demonstrated by Apollo on the Moon, but none of which have been achieved by the MERs nor will be achieved by the more capable (and vastly more expensive) Mars Science Laboratory (MSL) that is due to land on Mars in 2012.

More objective estimates of the relative efficiency of robots and humans as field geologists have been given by Garvin *et al.* (2004) and Snook *et al.* (2007). Garvin (2004) summarized the results of a NASA survey of several dozen planetary scientists and engineers on the relative efficiency of human and robotic capabilities

Table 1: Humans vs robots in space

skill	objective measurement	humans	advantage	robots
strength	Y	high strength/high torque; sometimes too strong		load/torque can be varied over very wide range with precise control
endurance	Y	limited by available consumable and physical tolerances		limited by design, environmental decay
precision	Y	high degree of training is required to ensure repeated performance in humans		once programmed, robot precision is limited only by electromechanical design
cognition	N	creative and limited only by prior training		execution of preprogrammed routines
perception	N	highly integrated sensory suite of limited use in space environment, visual acuity is very high		can detect minute environmental changes; can sense trace elements in low concentration
detection	Y	high detection sensitivity, though sensory paths limited during exploration		extremely high detection ability if preprogrammed and equipped with proper sensors
sensory acuity	Y	highly integrated sensory suite of limited use in space environment, visual acuity is very high		capable of detecting minute environmental changes if so equipped
speed	Y	able to cover great distances quickly		able to work very slowly and steadily
response time	Y	spot decisions and rapid response is customary, sometimes a disadvantage		rapid to programmed events, latency delay for "hold" events
decision making	N	flexible, unlimited in either speed or capacity		primitive learning capacity to scripted events
reliability	Y	high in terms of meeting mission objectives, but require support systems of high complexity		high reliability, but relatively short lives in space exploration environments
adaptability	N	highly adaptable to new and changing situations		reprogrammable to a limited extent, otherwise limited by design and system redundancy
agility	N	agility limited only by design of exoshell		computation requirements dictate slow movements with limited agility
versatility	N	readily self-programmable to provide multipurpose services and functions		generally designed to perform specific functions and poorly equipped to new applications
dexterity	Y	ability to manipulate large and very small objects with high flexibility		can exhibit very high DoF and fast reaction times
fragility	N	generally robust but total system failure can be caused by small effects		exploration robots generally very fragile, especially attendant instrument suites
expendibility	N	human life is precious and we place ever higher value on it		robot also high value – today we send robots to less interesting scientific sights on Mars because value is high – return unnecessary
maintainability	N	low cycle time between periodic consumable replenishments, requires expert skills to maintain		limited only by design – can be maintained by low-skill personnel

Tabular summary of exploration skills after Garvin (2004). The fourth column indicates relative advantage of humans or robots on a sliding scale (green: the advantage lies with humans; red: with robots; white: equal). In most cases humans have a clear advantage. Since 2004 the extreme endurance of the MERs has moved this entry more in favour of robots, but they remain slow and inflexible. (Courtesy of Jim Garvin/NASA/Springer)

in 18 different skill sets relevant to planetary exploration. The results are summarized in table 1, and show a clear balance in favour of human capabilities (with the implicit recognition that the most efficient exploration strategies of all will be those consisting of human-robotic partnerships where each complements the other).


This conclusion is corroborated by direct field comparisons of human and robotic exploration at planetary analogue sites on Earth. Snook *et al.* (2007) reported the results of one such study, conducted at the Haughton impact crater in the Canadian Arctic, where the efficiency of a human explorer (suitably encumbered in

a spacesuit) was compared with that of a tele-operated rover (controlled from NASA Ames Research Centre in California) in the performance of a range of exploration tasks. The rover was more sophisticated than those employed in present-day space missions, and included simulation of artificial intelligence capabilities

that are only likely to be incorporated in actual space missions from 2015 at the earliest. Nevertheless, the space-suited “astronaut” was found to be much more efficient in performing exploration tasks than the rover, and Snook *et al.* (2007 p438) concluded that “humans could be one to two orders of magnitude more productive per unit time in exploration than future terrestrially controlled robots”.

Although this estimate is an order of magnitude lower than Squyres’ off-the-cuff estimate of 1500 given above, this is mainly because the comparison was conducted between human and *tele-robotic* exploration, rather than between humans and supervised quasi-autonomous robotic exploration such as carried out by the MERs and MSL. Tele-robotic exploration is known to be more efficient than autonomous robotic operation, precisely because real-time human interaction is involved, but it cannot be employed effectively on planetary surfaces more distant than the Moon because of the inevitable communications delay (Lester and Thronson 2011). Garvin (2004, see his figure 2) has compared the efficiencies of robotic, tele-robotic, and human exploration, from which it is clear that if humans are “one to two orders of magnitude more efficient” than tele-robots then they will be even *more* efficient when compared with robotic vehicles such as the MERs or MSL, bringing the two estimates into better agreement.

Moreover, while comparisons based on the relative time taken to perform certain tasks do indeed show humans to be more efficient than robots, they nevertheless grossly underestimate the added scientific value of having humans on planetary surfaces. This is because astronauts have to come back to Earth, and can therefore bring large quantities of intelligently collected samples back with them. Robotic explorers, on the other hand, generally do not return – one reason why they are cheaper – so nothing can come back with them. Even if robotic sample return missions are implemented, neither the quantity nor the diversity of these samples will be as high as would be achievable in the context of a human mission – again compare the 382 kg of samples (collected from more than 2000 discrete locations) returned by Apollo, with the 0.32 kg (collected from three locations) brought back by the Luna sample return missions. The Apollo sample haul might also be compared with the ≤ 0.5 kg generally considered in the context of future robotic Mars sample return missions (e.g. ISAG 2011). Note that this comparison is not intended in any way to downplay the scientific importance of robotic Mars sample return, which will in any case be essential before human missions can responsibly be sent to Mars, but merely to point out the step change in sample availability (both in quantity and diversity) that may be expected when and if human missions *are* sent to the planet.



2: Cumulative number of refereed publications in the ADS database for the Apollo, Luna 16/20/24, Lunokhod, Surveyor, and Mars Exploration Rover missions. See “Bibliometric search details” on page 2.26 for more information.


Bibliometrics

If, as argued above, human exploration of planetary surfaces is really scientifically so much more efficient than robotic exploration, then we would expect to see this reflected in the scientific literature. As the only human exploration missions to date are the Apollo missions, any such bibliometric comparison must be between publications based on Apollo data and those based on various robotic missions to the Moon and Mars. With this in mind, figure 2 shows the cumulative number of refereed publications recorded in the SAO/NASA Astrophysics Data System (ADS) resulting from the six successful Apollo landings, the three Luna sample return missions (Lunas 16, 20, 24), the two tele-operated Lunokhod rovers (Lunas 17, 21), the five successful Surveyor lunar soft landers, and the two MERs (Spirit and Opportunity) on Mars. Those interested will find more details of the search parameters in the box on page 2.26.

Several things are immediately apparent. Most obvious is the sheer volume of Apollo’s scientific legacy compared to the other missions illustrated. This alone goes a long way to vindicate the points made above about human versus robotic efficiency. The second point to note is that the next most productive set of missions are the lunar sample return missions Lunas 16, 20 and 24, which highlights the importance of sample return. Indeed, a large part of the reason why Apollo has resulted in many more publications than the Luna missions is the much larger quantity and diversity of the samples – which will always be greater in the context of human missions. The third point to note is that, despite being based on data obtained and samples collected more than 40 years ago, and unlike the Luna, Lunokhod, or Surveyor publications, which have clearly levelled off, the Apollo publication rate is still rising. Indeed, it is actually rising as fast as, or

faster than, the publications rate derived from the Mars Exploration Rovers, despite the fact that data derived from the latter are much more recent. No matter how far one extrapolates into the future, it is clear that the volume of scientific activity generated by the MERs, or other robotic exploration missions, will never approach that due to Apollo.

However, to my mind, the most staggering thing about figure 2 is that this enormous scientific legacy is based on a total of only 12.5 days total contact time with the lunar surface. Note that this is the total cumulative time the Apollo astronauts were on the Moon, including downtime in the Lunar Module, not the cumulative EVA time, which was just 3.4 days (Orloff and Harland 2006). This may be compared with a total of 436 active days on the surface for the Lunokhods (Wilson 1987) and 5162 days for the Mars Exploration Rovers (to the end of 2011, allowing for the fact that contact was lost with Spirit on 22 March 2010). This comparison is illustrated in figure 3, which shows the cumulative number of publications divided by days of fieldwork on the surface (adopting 12.5 days for Apollo to allow for a fair comparison with the rovers). This is the same as dividing the cumulative number of publications by the number of sites studied up to a given date and the average days of fieldwork per site. Dips in the cumulative curves occur when a new mission arrives, instantaneously increasing the days of fieldwork before this has fed through to increased publications. Note the logarithmic scale – by this metric, Apollo was over three orders of magnitude more efficient in producing scientific papers per day of fieldwork than are the MERs. This is essentially the same as Squyres’ (2005) intuitive estimate given above, and is consistent with the more quantitative analogue fieldwork tests reported by Snook *et al.* (2007).



3: Refereed publications per day exploring the surface of the Moon or Mars. This is the same as dividing the cumulative number of publications by the number of sites visited by a given date and the average time spent there. The plot covers those missions plotted in figure 2 which had, or have, surface mobility, and which can therefore be considered as having performed “fieldwork”.

Human exploration costs

Although it is generally taken for granted that human exploration is more expensive than robotic exploration, and this is certainly true if the aggregate costs are the only ones considered, the situation is not as clear cut as it is sometimes made out to be. For one thing, the ratio of costs between human and robotic missions, while large, may nevertheless be smaller than the corresponding ratio in scientific productivity. The Apollo missions are instructive in this respect. Wilhelms (1993) and Beattie (2001) estimated a total cost of Apollo as \$2.5bn “in 1960s money”. This is rather more than the Congressional appropriations for Apollo (\$19.4bn from 1961 to 1973; tabulated by Orloff and Harland 2006). Taking the higher estimate (to be conservative) and taking “1960s” to be 1966 when Apollo expenditure peaked, this corresponds to about \$175bn today (where I have made use of the US Bureau of Statistics Inflation Calculator at http://www.bls.gov/data/inflation_calculator.htm).

It is interesting to compare this with the cost of a modern state-of-the-art robotic mission such as Mars Science Laboratory. MSL, which at this writing is en route to Mars, has cost an estimated \$2.5bn (Leone 2011). Thus, in real terms, Apollo cost 70 times as much as MSL. However, Apollo visited six sites, whereas MSL will visit one. In terms of cost-per-site Apollo was only 12 times dearer than MSL, yet each Apollo mission was vastly more capable. It is true that this comparison only strictly holds in the context of lunar exploration, where we can compare Apollo with a hypothetical future MSL-like lunar rover. In the context of Mars exploration, human missions seem likely to be more expensive than Apollo in real terms, although not necessarily by a large factor. The estimated total costs of some human Mars mission architectures are comparable to that of Apollo, or even lower (Turner 2004). The main point is that human

missions like Apollo are between two and three orders of magnitude more efficient in performing exploration tasks than robotic missions, while being only one to two orders of magnitude more expensive. In addition, human missions can accomplish scientific objectives that are unlikely to be achieved robotically at all (deep drilling and properly representative sample collection and return are obvious examples, as well as the increased opportunities for serendipitous discoveries). Looked at this way, human space exploration doesn’t seem so expensive after all!

That said, there is a more sophisticated and productive way to view the relative costs of human and robotic spaceflight. The fact is that while robotic planetary missions are science-focused, and essentially their whole costs are therefore borne by scientific budgets, human spaceflight is not wholly, or even mainly, science-driven. Rather, the ultimate drivers of human spaceflight tend to be geopolitical concerns, industrial development and innovation, and employment in key industries. Thus, science can be a beneficiary of human missions instituted and (largely) paid for by other constituencies. Apollo again provides an excellent example: it was instituted for geopolitical rather than scientific reasons, and to first order the US government’s expenditure of \$2.5bn (\$175bn today) would have occurred anyway, whether any science was performed or not. Fortunately, owing largely to the efforts of a relatively small number of senior scientists (Beattie 2001) scientific objectives and capabilities were incorporated into Apollo and resulted in the rich scientific legacy that is still being exploited today.


Of course, including science in Apollo did not come entirely free. It is interesting to compare this additional, strictly scientific, investment with the cost of robotic missions that were, and are, demonstrably less capable than Apollo. Beattie (2001) has studied the cost of including

science in Apollo (including the Apollo Lunar Surface Experiment Packages [ALSEPs], grants to the ALSEP investigators, construction of the Lunar Receiving Laboratory in Houston, grants to the initial tranche of lunar sample investigators, astronaut ALSEP and geological training etc), and arrived at a figure of \$350m in 1972 dollars (somewhat higher than other published estimates). Beattie does not explicitly include the additional cost of developing the Lunar Roving Vehicle (\$37m), but this should probably be added as the LRV was included in the last three Apollo missions mainly to enhance geological exploration. Thus we arrive at a total *scientific* cost in Apollo of \$387m in 1972 dollars. This corresponds to about \$2.09bn today, or 1.2% of the total Apollo budget.

Comparison with robotic costs

It is instructive to compare this with the costs of some past and planned robotic missions. It is sometimes difficult to get reliable estimates for these, but a search of various internet sources gives \$265m for Mars Pathfinder (which landed on Mars in 1997), \$820m for the two Mars Exploration Rovers (landed in 2004 and considering only the first 90 days of operations), and, as noted above, \$2.5bn for Mars Science Laboratory. Again employing the US Bureau of Statistics Inflation Calculator, these correspond to \$374m, \$982m, and \$2.5bn in 2011 dollars. For comparison, according to a NASA Planetary Sciences Decadal Survey Steering Committee report available online (Li and Hayati 2010), the estimated cost of the Mars Sample Return (MSR) mission proposed for around 2025 is about \$6.5bn. These mission costs are compared in figure 4, from which two points arise:

- The cost of the robotic exploration of Mars has *not* been decreasing as technology advances, but has been increasing steadily. There is a good reason for this: planetary surfaces are large, rough, rugged places, not at all amenable to exploration with small, cheap rovers no matter how much “intelligence” is built into them. There isn’t much point in having a hyper-intelligent rover the size of a matchbox if it can only travel 5m a day and gets stuck every time it encounters a rock the size of a brick. Nor is such a vehicle likely to carry much in the way of instrumentation. As a consequence, rovers have become bigger and more expensive (and more capable) with time (figure 1). This is exactly opposite to the trend predicted by some (e.g. Rees 2011), but it is one which will continue if we persist in trying to explore planetary surfaces with robots.
- In real terms, the cost of Mars Science Laboratory (\$2.5bn) already exceeds the additional cost of flying science on Apollo (\$2.09bn in today’s money). In fact, as Apollo visited six different sites on the Moon, the cost of science per site (\$348m in 2011 dollars) is actually less than the cost of *any* of these robotic Mars



4: Estimated costs of robotic Mars rover missions expressed in constant 2011 dollars. As currently conceived, MSR requires two rovers (possibly including ESA's ExoMars) in addition to other expensive elements. For how much longer will this be sustainable within purely scientific budgets? Note that this is not an inflationary increase (real-terms costs are plotted): we could continue to build rovers as small and (relatively) cheap as Pathfinder, but choose not to owing to their inherent scientific limitations.

missions and, as described above, the scientific efficiency was incomparably higher.

Conclusions

The lesson seems clear: if at some future date a series of Apollo-like human missions return to the Moon and/or are sent on to Mars, and if these are funded (as they will be) for a complex range of socio-political reasons, scientists will get more for our money piggy-backing science on them than we will get by relying on dedicated autonomous robotic vehicles which will, in any case, become increasingly unaffordable.

Fortunately, there is a way forward. In 2007 the world's space agencies came together to develop the Global Exploration Strategy (GES), which lays the foundations for a global human exploration programme that could provide us with just such an opportunity (GES 2007). One of the first fruits of the GES has been the development of a Global Exploration Roadmap (GER 2011), which outlines possible international contributions to human missions to the Moon, near-Earth asteroids and, eventually, Mars. The motivations for the GES are, needless-to-say, multifaceted, and include a range of geopolitical and societal motivations (many of them highly desirable in themselves) in addition to science.

Science would be a major beneficiary of participating in a human exploration programme such as envisaged by the Global Exploration Strategy. Quite simply, this will result in new knowledge, including answers to fundamental questions regarding the origin and evolution of planets, and the distribution and history of life in the solar system, that will not be obtained as efficiently, and in many cases probably not obtained at all, by reliance on robotic exploration alone. ●

I A Crawford is in the Department of Earth and Planetary Science at Birkbeck College London, UK;

i.crawford@bbk.ac.uk.

Acknowledgments. I thank Drs Peter Grindrod, Katherine Joy and Heino Falcke, who acted as sounding boards, and the LPI Librarian Mary Ann Hager for advice on the bibliometric data. This research used NASA's Astrophysics Data System.

References

- Beattie DA** 2001 *Taking Science to the Moon: Lunar Experiments and the Apollo Program* (Johns Hopkins University Press, Baltimore).
- Clements D** 2009 *Physics World* **22**(11) 16.
- Close F et al.** 2005 *Report of the RAS Commission on the Scientific Case for Human Space Exploration* <http://tinyurl.com/RAS-HSEreport>.
- Coates AJ** 2001 *Earth Moon Planets* **87** 213–219.
- Cockell CS** 2004 *Earth Moon Planets* **94** 233–243.
- Crawford IA** 2001 *Earth Moon Planets* **87** 221–231.
- Crawford IA** 2010 *Astrobiology* **10** 577–587.
- Garvin J** 2004 *Earth Moon Planets* **94** 221–232.
- GER** 2011 *The Global Exploration Roadmap* <http://tinyurl.com/global-exp>.
- GES** 2007 *The Global Exploration Strategy: Framework for Coordination* <http://tinyurl.com/GES-framework>.
- ISAG** 2011 *Planning for Mars Returned Sample Science: Final Report of the MSR End-to-End International Science Analysis Group* <http://tinyurl.com/mepag>.
- Leone D** 2011 *Space News International* 8 June 2011 <http://tinyurl.com/msl-needs>.
- Lester D and Thronson H** 2011 *Space Policy* **27** 89–93.
- Li F and Hayati S** 2010 *NASA PSDS Mars Sample Return Discussions* <http://tinyurl.com/MSL-doc>.
- NRC** 2005 *Assessment of Options for Extending the Life of the Hubble Space Telescope* (National Academies Press, Washington DC).
- Orloff RW and Harland DM** 2006 *Apollo: The Definitive Sourcebook* (Springer-Praxis, Chichester).
- Rees MJ** 2011 *Proc. IAU* **260** 16–21.
- Snook K et al.** 2007 in *The Geology of Mars*: ed. M Chapman (CUP, Cambridge) 424–455.
- Spudis P** 2001 *Earth Moon Planets* **87** 159–171.
- Squyres S** 2005 *Roving Mars: Spirit, Opportunity, and the Exploration of the Red Planet* (Hyperion, New York).
- Turner MJL** 2004 *Expedition Mars* (Springer-Praxis, Chichester).
- Wilhelms DE** 1993 *To a Rocky Moon: A Geologist's History of Lunar Exploration* (University of Arizona Press, Tucson).
- Wilson A** 1987 *Solar System Log* (Jane's, London).

Bibliometric search details

The bibliometric statistics shown in figures 2 and 3 were obtained from the SAO/NASA ADS database. The searches were restricted to refereed publications. It is likely that these statistics underestimate the total number of publications resulting from these missions, especially for Apollo, as some sample analysis work has been published in geological and petrological journals not normally included in the ADS (although, through the efforts of the library of the Lunar and Planetary Institute [LPI], the ADS should be complete up to about 1995 when the LPI Lunar Bibliography was turned over to the ADS; M A Hager, personal communication, 2011). I have made use of the ADS here principally because of its ease of use, and because it is a well-respected bibliometric database in the astronomical community.

The Apollo publications were retrieved by searching for papers containing the words “Apollo” AND “Moon” in the title or abstract (including “Moon” was necessary to exclude papers referring to the Apollo asteroids). These were then examined to check that they do indeed relate to the Apollo missions. Similarly, the Surveyor papers were retrieved by searching for “Surveyor” AND “Moon” (where this time “Moon” was included to exclude references to surveys unrelated to the Surveyor missions). The Luna 16/20/24 and Lunokhod publications were retrieved by searching for these missions by name. Finally, the MER publications were retrieved by searching for [“(“Spirit” OR “Opportunity”) AND (“rover” OR “Gusev” OR “Meridiani”)] OR “Mars Exploration Rover”]; this slightly complicated set of search criteria was rendered necessary to avoid the surprisingly large number of papers reporting work conducted in such and such a “spirit” or which provide great “opportunities” for various things unrelated to Mars exploration! A check was made against the MER Science Team's own publication list (http://marsrover.nasa.gov/science/pdf/web_publist.pdf) to ensure that these papers were correctly recovered by the ADS search criteria (although the Team papers are only a subset of the total number of papers making reference to the MERs included in figure 2, this agreement gives confidence that a large number of MER papers have not been missed in the ADS search)