

Spec and Proj - Exercises

September 30, 2021

1. A variety V is called *irreducible* if it can't be decomposed as $V = V_1 \cup V_2$ in a non-trivial way. Show that this is equivalent to (a) any two Zariski open subsets in V intersect, or (b) the ring $\mathbb{C}[V]$ is an integral domain.
2. (a) Show that Nullstellensatz fails over \mathbb{R} by finding a polynomial f such that $I_{V(f)}$ is bigger than $\text{rad}(f)$. Generalize to any non-algebraically-closed field.
(b) Over \mathbb{C} or \mathbb{R} the ideal $I_{\mathbb{A}^n}$ is zero. What happens over \mathbb{F}_p ?
3. Show that for any hypersurface $V(f) \subset \mathbb{A}^n$, the complement $\mathbb{A}^n \setminus V(f)$ is isomorphic to an affine variety.
4. Prove that every homomorphism $\mathbb{C}[W] \rightarrow \mathbb{C}[V]$ is induced from a regular map $V \rightarrow W$.
5. Let R be the ring $R = \mathbb{C}[x]/(x^2)$. Show that a homomorphism $\mathbb{C}[V] \rightarrow R$ is exactly the data of a point $p \in V$ and a vector which is tangent to V at p .
6. Let $V = V(y^2 - x^3) \subset \mathbb{A}^2$ and let $f : \mathbb{A}^1 \rightarrow V$ be the function:

$$f : t \mapsto (t^2, t^3)$$

Show that f is not an isomorphism but it is a bijection (*and even a homeomorphism in the usual complex topology*).

7. Compute the transition functions (*i.e.* the co-ordinate changes) between the standard affine charts in \mathbb{P}^2 .
8. Find an explicit method to compactify any hypersurface in \mathbb{A}^n to a hypersurface in \mathbb{P}^n .
9. (a) Let C be the projective curve $V(xy - z^2) \subset \mathbb{P}^2$. Construct a bijection $\mathbb{P}^1 \rightarrow C$.
(b) What does this have to do with traceless 2×2 matrices of rank 1?
(c) What happens if $xy - z^2$ is replaced with another quadratic form?
10. (a) Consider the complex affine curve $V_\epsilon = V(xy - \epsilon) \subset \mathbb{A}^2$ for $\epsilon \in \mathbb{C}$. Convince yourself that (i) for $\epsilon \neq 0$ the space V_ϵ is a cylinder, and (ii) as $\epsilon \rightarrow 0$ one circle in the cylinder collapses, leaving two discs glued at a point.
(b) What's the topology of the complex projective plane curve $V(xy) \subset \mathbb{P}^2$? Now consider a small perturbation $f = xy + \epsilon g$ for some quadratic $g(x, y, z)$. What's the topology of $V(f)$? *Compare this with Q9.*
(c) Now take a cubic plane curve of the form $V = V(xyz + \epsilon h(x, y, z))$. Argue that V has the topology of a torus.