

“Shear thickening” in non-shear flows: the effect of microstructure

Helen J. Wilson

Department of Mathematics, University College London, Gower Street, London WC1E 6BT, UK

(Received 12 December 2017)

The bizarre behaviour of a cornstarch suspension (sometimes called oobleck) is well known to all of us who have led public engagement events. At the right solids fraction, it flows smoothly at slow speeds, but can be shattered with a quick spoon movement; if you prepare a large enough sample you can run across the surface (but if you stand still you will sink).

In rheology circles this phenomenon is known as shear thickening, though the flows described above are not necessarily shear dominated. In recent years there has been a proliferation of research on the mechanism behind true shear thickening, using both experiments and numerical simulations of shear flows. The understanding of the underlying mechanism is improving markedly. But the paper *Microstructure and thickening of dense suspensions under extensional and shear flows* (Seto, Giusteri & Martinello, JFM 2017) is the first to consider more general flows. We have, for the first time, simulations of thickening in extensional flows, which are a far better description of oobleck with a runner on top – and can begin to quantify the difference between the idealised shear thickening and the extension thickening that happens in practice.

Keywords. Suspensions; Stokesian dynamics; Rheology.

1. Introduction

Suspension mechanics – the study of fluid containing solid particulates – has a long and distinguished history across more than a century. Einstein (1906) showed that for a dilute suspension of spheres at volume fraction ϕ in a fluid of viscosity μ , to leading order in ϕ the suspension acts as a Newtonian fluid having viscosity

$$\eta = \mu \left(1 + \frac{5}{2} \phi \right).$$

Batchelor & Green (1972) extended this result to the next order in ϕ , showing that for *steady* extensional flows,

$$\eta \approx \mu \left(1 + \frac{5}{2} \phi + 6.9 \phi^2 \right);$$

but for shear flows the appearance of Jeffrey orbits, on which a pair of particles oscillate around one another indefinitely, makes it impossible to determine the viscosity. And as soon as inter-particle forces are introduced, the rheology at order ϕ^2 becomes non-Newtonian (Brady & Morris 1997), with the suspension exhibiting first and second normal stress differences in steady simple shear flow.

These results are for dilute suspensions – suspensions having a very low solids volume fraction – and even with that strong simplification, we do not have a constitutive relation,

a governing equation for the behaviour of these materials. The best attempt so far is by Phillips *et al.* (1992), who produced a model which captures shear-induced particle migration and a viscosity based on the local concentration of solids; but this model does not incorporate any microstructure beyond the concentration, nor predict any non-Newtonian effects.

The 1997 paper of Brady & Morris was seminal in its consideration of the microstructure: the arrangement of particles within the suspension. The most recent efforts in this area still draw on that early insight: such works as Fernandez *et al.* (2013); Heussinger (2013); Seto *et al.* (2013) have used particle simulations to explore the microstructure emerging within shear flows under the influence of different forces between the particles. The most recent success in the area has been the reproduction of the phenomenon of *discontinuous shear thickening*, the abrupt increase in the steady shear viscosity of a suspension when the shear rate is increased beyond some critical rate. It is now widely accepted that a key ingredient of the phenomenon is “stress-induced friction” (Wyart & Cates 2014; Mari *et al.* 2014), and this has been confirmed by experiment (Lin *et al.* 2015; Clavaud *et al.* 2017).

However, real observations of thickening in practical flows are rarely in steady shear. We need to understand the evolution of the suspension microstructure and stress in other flows – steady and unsteady. This paper takes the first step in this process, by considering steady extensional flows.

2. Overview

The paper *Microstructure and thickening of dense suspensions under extensional and shear flows* (Seto *et al.* 2017) carries out particle dynamics simulations of concentrated suspensions in both shear and planar extensional flow. The authors use a combination of a repulsive force and frictional contact between particles, which gives similar behaviour to the stress-induced friction model of contact between particles that has been so successful of late in reproducing discontinuous shear thickening, and they investigate the microstructure of the suspension and its average stress during the flow.

In shear they are simply reproducing observations we have seen before, which nonetheless bear repeating. For slow flows below the shear thickening transition, there is substantial long-range ordering within the flow, as particles line up in the flow direction. Above the transition, frictional contacts between particles disrupt the structure, and there is global disorder. This has widely been accepted in the literature as being a fundamental part of the mechanism of shear thickening.

However, there are two key extensions here. Let us look first at the change of flow type. In steady planar extensional flow, the authors find that *there is no global ordering* even at low flow rates. This is not a huge surprise, as there are no straight streamlines along which the particles could align. We see the clear differences between shear and extensional flows in figure 1, in which we plot the pair distribution function $g(\mathbf{r})$, which describes the likelihood of finding a particle centred at a given location given that one test particle is centred at the origin. On the left are the plots for shear flow below and above the shear-thickening transition. The global ordering at low flow rates is very clear. On the right we see the corresponding plots for low and high strain rates in plane extension flow. Although there are differences between them, they are qualitatively very similar – there is no order–disorder transition here.

What is surprising is that, despite the lack of an order–disorder transition, *we see thickening in extensional flow* just as we do in shear. There are subtleties in the definition of the flow rate, and of the viscous contribution to the stress, in these different flows; but




FIGURE 1. (a) The pair distribution function $g(\mathbf{r})$ below thickening (left) and above thickening (right) for simple shear flow of a monodisperse suspension. (b) The equivalent plots for planar extensional flow. Taken from Seto *et al.* (2017), figure 6.




FIGURE 2. Plots of effective viscosity against flow rate for suspensions at different volume fractions. Solid symbols, solid lines: extensional flow; Open symbols, dashed lines: shear flow. Taken from Seto *et al.* (2017), figure 2.

using a coherent convention introduced by Giusteri & Seto (2017) they can plot viscosity against flow rate for both flows on the same graph, shown in figure 2. The thickening is not quite as extreme in extensional flow as in shear, but it is quite unarguably discontinuous thickening.

The second key extension considered in this paper is to move away from monodisperse suspensions and (as a proxy for true polydispersity) use spheres of two different sizes. Even weak polydispersity (a size ratio of 1.2) has the effect of breaking up the long-range order in slow shear flow, without making much difference to the shear thickening in the suspension. Polydispersity makes shear flow look very much like extension.

It is, therefore, clear that the mechanism of shear thickening is *not* closely tied to the disruption of a long-range order within the suspension; the long-range order itself is a symptom of the idealisations made in earlier simulations (perfect monodispersity and pure shearing flow).

The authors go on to investigate their microstructure further and find that the key to thickening is in the distribution of *contacting* particle pairs. These pairs are all (necessarily) the same distance apart, so their angular distribution is all that matters: and this angular distribution changes dramatically at the thickening transition, in both shear and extensional flows. Below thickening, the contacts are largely in a single direction (or absent in shear flow); above thickening there is a wide angular distribution. This indicates that the key factor in thickening is the development of a *contact network*, with

force chains running throughout the suspension. The microstructure also causes non-Newtonian effects in both flow types.

3. Future

A key challenge of suspension mechanics is the generation of a constitutive model that can be used for complex flows: flows with too much spatial variation for full particle simulations to be practical. Such a constitutive model would need to take into account the underlying mechanisms of shear (and indeed extension) thickening, as well as the construction and destruction of microstructure by flow.

An immediate next step from the current work may be to investigate crossover flows between shear and extension; or flows whose strength varies with time, and the processes governing the evolution of the microstructure. But there is no question that this first foray into extensional flows of suspensions is a key step towards our goals.

REFERENCES

- BATCHELOR, G K & GREEN, J T 1972 The determination of the bulk stress in a suspension of spherical particles to order c^2 . *Journal of Fluid Mechanics* **56** (2), 401–427.
- BRADY, J F & MORRIS, J F 1997 Microstructure of strongly sheared suspensions and its impact on rheology and diffusion. *Journal of Fluid Mechanics* **348**, 103–139.
- CLAUDAUD, C, BÉRUT, A, METZGER, B & FORTERRE, Y 2017 Revealing the frictional transition in shear-thickening suspensions. *Proceedings of the National Academy of Sciences* **114** (20), 5147–5152.
- EINSTEIN, A 1906 Eine neue Bestimmung der Moleküldimensionen. *Annalen der Physik* **324** (2), 289–306.
- FERNANDEZ, N, MANI, R, RINALDI, D, KADAU, D, MOSQUET, M, LOMBOIS-BURGER, H, CAYER-BARRIOZ, J, HERRMAN, H J, SPENCER, N D & ISA, L 2013 Microscopic mechanism for shear thickening of non-Brownian suspensions. *Phys. Rev. Lett.* **111**, 108301.
- GIUSTERI, G G & SETO, R 2017 A theoretical framework for steady-state rheometry in generic flow conditions. ArXiv:1702.02745.
- HEUSSINGER, C 2013 Shear thickening in granular suspensions: Interparticle friction and dynamically correlated clusters. *Phys. Rev. E* **88**, 050201.
- LIN, N Y C, GUY, B M, HERMES, M, NESS, C, SUN, J, POON, W C K & COHEN, I 2015 Hydrodynamic and contact contributions to continuous shear thickening in colloidal suspensions. *Phys. Rev. Lett.* **115**, 228304.
- MARI, R, SETO, R, MORRIS, J F & DENN, M M 2014 Shear thickening, frictionless and frictional rheologies in non-brownian suspensions. *J. Rheol.* **58**, 1693–1724.
- PHILLIPS, R J, ARMSTRONG, R C, BROWN, R A, GRAHAM, A L & ABBOTT, J R 1992 A constitutive equation for concentrated suspensions that accounts for shear-induced particle migration. *Physics of Fluids* **4**, 30–40.
- SETO, R, GIUSTERI, G G & MARTINIELLO, A 2017 Microstructure and thickening of dense suspensions under extensional and shear flows. *J. Fluid Mech.* **?**, **?**
- SETO, RYOHEI, MARI, ROMAIN, MORRIS, JEFFREY F. & DENN, MORTON M. 2013 Discontinuous shear thickening of frictional hard-sphere suspensions. *Physical Review Letters* **111** (21), 218301.
- WYART, M & CATES, M E 2014 Discontinuous shear thickening without inertia in dense non-Brownian suspensions. *Physical Review Letters* **112** (9), 098302.