

# CHEMICAL-INDUCED SURFACE STRAINS IN GOLD NANOCRYSTALS

Ian Robinson<sup>1,2</sup>

<sup>1</sup>London Centre for Nanotechnology University College, London, WC1E 6BT, UK


<sup>2</sup>Diamond Light Source, Harwell Campus, Didcot, OX11 0DE, UK

The structure and dynamics of the sulphur-gold interface are remarkably complex. A recent crystallographic study of a monodispersed Au<sub>101</sub>RS<sub>44</sub> gold nanoparticle-thiol complex revealed a crystalline core particle coated with a shell of 1nm thickness with enlarged Au-Au spacings and interpenetration of the thiol ligand species. Far from having a well-defined boundary between the metal and the organic sides of the interface, this unusual complex was found to contain a mixed compound layer as its lowest energy configuration [1]. We report the structure of faceted gold nanocrystals before and after coating with propane thiol, C<sub>3</sub>H<sub>7</sub>SH, one of the simplest SAM-forming organic molecules. We found our gold nanocrystals to be spherical with {111} facets, as expected from the theory of Equilibrium Crystal Shapes (ECS). The structure of a single 300nm-diameter particle was measured using the powerful technique of Coherent X-ray Diffraction (CXD), which is highly sensitive to the pattern of internal strains within the nanocrystal [2]. Our results show that the strain is profoundly modified by the thiol adsorption. We are able to establish the magnitude of this stress difference using finite element calculations and suggest a model involving preferential reaction on the curved regions of the crystal surface [3].

[1] "Structure of a thiol monolayer-protected gold nanoparticle at 1.1 angstrom resolution" P. D. Jadzinsky, G. Calero, C. J. Ackerson, D. A. Bushnell and R. D. Kornberg, Science 318 430-3 (2007)

[2] "Coherent Diffraction Imaging of Strains on the Nanoscale", I. K. Robinson and R. Harder, Nature Materials 8 291-298 (2009)

[3] "Shape-dependent Thiol-induced strains in gold nanocrystals", Moyu Watari, Rachel McKendry, Manuel Vögli, Gabriel Aeppli, Yeong-Ah Soh, Xiaowen Shi, Gang Xiong, Ross Harder and Ian Robinson in preparation



*Figure showing the shape of the crystal and the phase found on the surface of the particle before dosing on a colour scale from -1 (blue) to +1 radian (red). The arrows indicate the {111} directions; the substrate surface normal direction is parallel to the large facet at the top of the figure. The (111) Q-vector used for imaging is on the right hand side. The phase is the projection of the crystal's displacement field onto this direction [2].*

**Invited Oral preference:**

**Corresponding Author:** i.robinson@ucl.ac.uk

Abstracts should be submitted via e-mail to [ecsd-10@reading.ac.uk](mailto:ecsd-10@reading.ac.uk) with the subject Abstract plus the presenting authors surname. **The deadline for abstract submission will be 14<sup>th</sup> May 2010.** Notification of acceptance will be by 1<sup>st</sup> June.

**Authors are required to register before attending the conference.**