

Imaging of Nanocrystals by Inversion of Coherent X-ray Diffraction


- Ian Robinson
- Franz Pfeiffer
- Ivan Vartanians
- Ross Harder
- Meng Liang
- Mark Pfeifer
- Garth Williams

Physics Department
University of Illinois

ICTP Advanced X-ray
Imaging, June 2005

Tomato Bushy Stunt Virus

1980




I. K. Robinson ICTP, Trieste

Outline

- Coherent X-rays can Image Structure in 3D
- Solve Phase Problem by Phase Retrieval
- Internal structure of Nanocrystals
- Imaging of **Phase** Objects
- How small can we go?

Generic “Error Reduction” method






J. R. Fienup *Appl. Opt.* 21 2758 (1982)

R. W. Gerchberg and W. O. Saxton *Optik* 35 237 (1972)

I. K. Robinson ICTP, Trieste


Lensless X-ray Microscope







I. K. Robinson ICTP, Trieste

Coherent Diffraction from Crystals






Microscope Images of Gratings




5 μ m grid lines






SEMS

- Au blanket film
- Quartz substrate
- Annealed at 950°C for 70 hrs.







Symmetrized Data and two best fits

Chisq=0.0005












3D Diffraction Method







Slices through
plan view SEM:




Diffraction by Strain of Point Defect

$$A \sim \sum e^{i\mathbf{Q}\cdot(\mathbf{R}_j+\mathbf{u}_j)} \\ \approx \sum e^{i\mathbf{Q}\cdot\mathbf{R}_j} (1+i\mathbf{Q}\cdot\mathbf{u}_j)$$

Imaginary density




Modeling of 3D Phase Bump



I. K. Robinson ICTP, Trieste


Field lines of Point Charges





I. K. Robinson

ICTP, Trieste

Lensless X-ray Microscope





CXD from Silver Nanocubes





Yugang Sun and Younan Xia,
Science 298 2177 (2003)

Spherical and cylindrical waves




$\pi/4$

$\pi/2$





$\pi/4$ (x)


Reconstruction of Ag Nanocrystal




←→
200nm



Rocking
scan of Ag
cubes with
 0.01° steps






Miscentered Non-Centrosymmetric Reciprocal Space Slice




$$A_q \neq A_{\bar{q}}^*$$





Real-space phase.







What can we expect to see?



$$A(\bar{q}_o + \bar{q}_z) = \sum_i \underbrace{\sum_z \rho(x, y, z)}_{u(x, y)} e^{2\pi i \bar{q}_z \cdot \bar{z}_i} e^{2\pi i (\bar{q}_o) \cdot \bar{r}_i}$$




Direct Space Result (1000 Iterations)



Compact Light Source


Ron Ruth, Lyncean Technologies



CLS Specifications

R. J. Loewen, Ph.D. Thesis
SLAC (2003)


50MeV ($\gamma=100$) + 1 μm laser



$$\omega = \frac{4\gamma^2}{(1 + \gamma^2\theta^2)} \omega_0$$

Total Average Flux	$\gtrsim 10^{13}$ ph/s
Avg. "Monochromatic" Flux*	$\approx 10^{10}$ ph/s
Source Spot Size [†]	30 μm radius
Source Divergence [†]	≈ 3 mrad
Source Brightness	$\gtrsim 10^{11}$
X-ray Energy Range [‡]	12 keV \pm 6 keV

Molecular Movies using XFEL



I. K. Robinson ICTP, Trieste

Conclusions

- Imaging of nanocrystals by phase retrieval
- 3D imaging to see inside structure
- Phase contrast to access strains
- Off-center diffraction implies phase also
- Extension to CLS and XFEL