

# Phase Retrieval and Applications

- Ian Robinson
- Ivan Vartanyants
- Franz Pfeiffer
- Mark Pfeifer
- **Garth Williams**
- Meng Liang
- Ross Harder


Department of Physics  
University of Illinois

Euro-XFEL Workshop  
Oct 28, 2005

# Outline

- Coherence + Phase Retrieval for Structure
- The **Phase** Problem
- Nanocrystal Shapes
- Imaging of **Phase** Objects
- Extrapolation and Technical Issues for XFEL
- “First” Euro-XFEL Experiment

# Generic “Error Reduction” method




J. R. Fienup *Appl. Opt.* 21 2758 (1982)

R. W. Gerchberg and W. O. Saxton *Optik* 35 237 (1972)


# Real-space Constraints in Crystallography

R. P. Millane, J. Opt. Soc Am. A **13** 725 (1996)

- ‘Positivity’ and ‘Atomicity’ constraints (Sayre)
- Finite **support**, molecular envelope
- Solvent flattening/Molecular replacement
- Non-crystallographic symmetry
- Non-uniqueness is ‘pathologically rare’ ( $d > 1$ )
- Uses memory to avoid stagnation (Fienup HIO)




# Coherent Diffraction from Crystals






# SEMS

- Au blanket film
- Quartz substrate
- Annealed at 950°C for 70 hrs.







# Symmetrized Data and two best fits

Chisq=0.0005












# 3D Diffraction Method







Slices through  
plan view SEM:



# 'Vortices' Form Loops in 3D




# Clearing of Vortices during HIO





ER ER HIO  
HIO ER ER  
HIO HIO ER


# Progress of Phase Retrieval



# Study of Convergence with Noise



# In situ growth of Pb crystals



# Good statistics, 3D diffraction data







Figure 4.12: Center slices from 3D CXD pattern from Pb sample, on a log scale. Data file 296 from 10/03.

# Learn shape of “tight” support




# Then refine amplitude *and* phase






# Fitting of crystal shape





# Density distribution across surface





# Diffraction by Strain of Point Defect

$$A \sim \sum e^{i\mathbf{Q}\cdot(\mathbf{R}_j+\mathbf{u}_j)} \\ \approx \sum e^{i\mathbf{Q}\cdot\mathbf{R}_j} (1+i\mathbf{Q}\cdot\mathbf{u}_j)$$

Imaginary density





# Modeling of Deformation Field




# Extrapolation to Euro-XFEL

- Steep coherent flux limit to accessible size scales:  $a^6$  or  $a^3$  ?
- $10^{12}$  photons/sec will be  $10^{12}$  photons/shot
- Already at 100nm, sample stability important
- Smaller objects move faster: scales as  $a^5$ .








# Reconstruction of Ag Nanocrystal




←→  
200nm




Rocking  
scan of Ag  
cubes with  
 $0.01^\circ$  steps






# Miscentered Non-Centrosymmetric Reciprocal Space Slice



$$A_q \neq A_{\bar{q}}^*$$






**Real-space phase.**







# What can we expect to see?

$$A(\bar{q}_o + \bar{q}_z) = \sum_i \underbrace{\sum_z \rho(x, y, z)}_{u_i(x, y)} e^{2\pi i \bar{q}_z \cdot \bar{z}_i} e^{2\pi i (\bar{q}_o) \cdot \bar{r}_i}$$



# Direct Space Result (1000 Iterations)





# “First” Euro-XFEL Experiment



- Critical nucleation at solid/liquid boundary
- Fluctuations of solid in liquid near MP
- Pre-critical nuclei during crystallisation
- Smaller nuclei harder to see: **focus**
- Smaller nuclei fluctuate more: **snapshot**

# Diffuse Scattering acquires Structure using CXD

Real Space



Reciprocal Space



# Conclusions and Outlook

- Phase Retrieval of CXD demonstrated
- Internal structure of Au Nanocrystals
- Phase structure visible in Pb Nanocrystals
- Focusing for access to much smaller clusters
- Thermal motion of small clusters important
- Euro-XFEL can solve both problems